

Review

Traditional oil palm (*Elaeis guineensis* Jacq.) and its medicinal uses: A review

Bamidele Victor Owoyele^{1,*}, Gbenga Opeyemi Owolabi²

¹Department of Physiology, University of Ilorin, P.M.B. 1515, Ilorin, Nigeria; ²Department of Physiology, Ladoké Akintola University of Technology, Ogbomoso, Nigeria

ABSTRACT

The oil palm (*Elaeis guineensis* Jacq.) has been reported to originate along the gulf of the guinea in West Africa. The various parts of the tree have been used locally and traditionally for various medicinal purposes. Some of these uses have been proved by scientific experiments. Palm oil is extracted from the mesocarp of the fruit and is used traditionally for the treatment of headaches, pains, rheumatism, cardiovascular diseases, arterial thrombosis and an atherosclerosis due to its rich phytonutrients. The leaves are also used for the treatment of cancer, cardiovascular diseases, kidney diseases and wound healing. The sap also has been found to be rich in phytonutrients that can be used to treat various diseases. This review therefore seeks to explore many of the uses of the oil palm using the various parts of the oil palm.

Keywords anti-inflammatory, anti-cancer, cardiovascular diseases, *Elaeis guineensis*, medicinal uses, oil palm, rheumatism

INTRODUCTION

It is generally agreed that the oil palm (*Elaeis guineensis* Jacq.) originated from the equatorial tropical rain forest region of Africa, precisely along the gulf of guinea (Naher et al., 2013). It exists in the wild type and cultivated state. It was first introduced to Brazil and other tropical countries in the 15th century by the Portuguese (Corley, 1976). The oil palm fruit is a drupe. The American oil palm, *Elaeis oleifera* is native to tropical Central America and South America.

This review intends to provide information on the documented medicinal uses of oil palm including laboratory studies on the plant. Earlier review by Obahiagbon (2012) was limited to an aspect of the plant. This present review covers both the traditional as well as the scientific uses of the oil. The review however does not extend to the core botanical aspect of the plant. The literatures used for this review were obtained from Medline and Google Scholar search carried out from September, 2013 to April, 2014.

The traditional theory about the uses of oil palm in many parts of Nigeria is that products from oil palm are antidotes that can be used in the treatment of many ailments especially gastrointestinal disorders and poisons.

Oil palm

The African oil palm (*Elaeis guineensis* Jacquin) produces two different kinds of oil namely, palm oil and palm kernel oil (Ekwenye and Ijeomah, 2005). The pericarp consists of three layers: the exocarp (skin), mesocarp (outer pulp containing palm oil), and the endocarp (a hard shell enclosing the kernel or

endosperm, which contains oil known as kernel oil) (Naher et al., 2013). Palm oil is extracted from fleshy mesocarp of the fruit either by milling mechanically or by the traditional method (Hartley, 1977; Edem and Akpanabiatu, 2006), which contains 45 - 55% oil, but varies from light yellow to orange-red in color, and melts at 25°C (Duke, 1983). The oil colour is determined by the carotenoids (Cottrell, 1991). The major carotenoids found in palm oil are the beta-carotene (Obahiagbon, 2012). Palm kernel oil is obtained from the kernels enclosed in the endocarp (Ekwenye and Ijeomah, 2005). Palm oil contains saturated palmitic acid, oleic and linoleic acid, giving it a higher unsaturated acid content than palm kernel or coconut oils (Duke, 1983; Edem, 2002). Along with coconut oil, palm oil is one of the few highly saturated vegetable fats. It is semi-solid at room temperature and contains several saturated and unsaturated fats in the forms of glyceryl laurate (0.1%, saturated), myristate (1%, saturated), palmitate (44%, saturated), stearate (5%, saturated), oleate (39%, monounsaturated), linoleate (10%, polyunsaturated), and alpha-linolenate (0.3%, polyunsaturated) (Cottrell, 1991). Palm oil also has minor constituents including phospholipids (Goh et al., 1983). Currently, palm oil is the world largest edible oil (Naher et al., 2013) and is the main source of domestic or edible oil in Africa (Obahiagbon, 2012; Oluba et al., 2009). As much as palm oil provides energy and fatty acid needs (Oguntibeju et al., 2009), much of the palm oil that is consumed as food is to some degree oxidized rather than in the fresh state, and this oxidation appears to be responsible for the health risk associated with consuming palm oil (Edem, 2002).

Ekwenye and Ijeomah (2005) reported that traditionally, palm oil has been used in the South Eastern Nigeria for the treatment of various diseases and skin infections. This was confirmed by their experiment carried out at the Michael Okpara University of Agriculture, Umudike, Abia State, Nigeria using five microorganisms namely; *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Candida albicans*, and *Aspergillus niger* known to cause some infections treatable with palm oil and palm kernel oil. In this experiment,

*Correspondence: Bamidele Victor Owoyele

E-mail: deleyele@yahoo.com or owoyele@unilorin.edu.ng

Received January 11, 2014; Accepted May 20, 2014; Published August 31, 2014

doi: <http://dx.doi.org/10.5667/tang.2014.0004>

© 2014 by Association of Humanitas Medicine

This is an open access article under the CC BY-NC license.

(<http://creativecommons.org/licenses/by-nc/3.0/>)

0.04 ml of 100% concentration palm oil and palm kernel oil extracts was impregnated into the sterilized discs according to the method of Ekpa and Ebana (1996). The anti-microbial test was done using the disc diffusion method of Brauer et al. (1996) was adapted using nutrient agar, sabouraud dextrose agar, and the impregnated discs. Palm oil has also been reported to be anodyne, anti-dotal, aphrodisiac and a diuretic (Irvin, 1985; Sasidharan et al., 2012). It is folk remedy for headaches, pains, rheumatism, cardiovascular diseases, arterial thrombosis and an atherosclerosis (Ekwenye and Ijeomah, 2005; Honstra, 1986; Irvin, 1985; Sasidharan et al., 2012). The palm oil is known to be effective against many forms of intestinal disorders, especially diarrhoea and dysentery in infants (Ekpa and Ebana, 1996.). The fruit mesocarp oil and palm kernel oil are administered as poison anti-dote and used externally with several other herbs as lotion for skin diseases. Palm kernel oil is applied to convulsing children to regulate their body temperature. Folk remedies of oil palm also include treatment for cancer and liniment (Irvin, 1985; Sasidharan et al., 2012).

Anti-oxidant activity

Palm oil provides a rich source of beta-carotene and vitamin E, namely tocopherols and tocotrienols which are recognized nutritional anti-oxidants that act as scavengers of the oxygen atom or free radicals (Chong and Ng, 1991; Ekwenye and Ijeomah, 2005; Goh et al., 1985). The oxygen atom or free radicals can arise during the body's normal oxidative metabolism or from the action of toxic pollutants that contaminate our food and have been implicated in ageing, heart disease and cancer (Ekwenye and Ijeomah, 2005). *E. guineensis* is also used as a wound healing agent among the natives of Africans and as therapeutic agent in other parts of the World (Sasidharan et al., 2010; Sasidharan and Vijayarathna, 2012). The palm oil is also used for biofuels and some manufactured products (Naher et al., 2013). Carotenoids are potent anti-oxidants. Packer et al. (1992) showed that alpha-carotene is a more potent anti-oxidant than beta-carotene. Red palm oil is rich in natural phytonutrients that are important for health. These phytonutrients (i.e. tocotrienols and carotenes) are also powerful anti-oxidants that help maintain the stability of the oil during cooking process and may extend the shelf life of food prepared with red palm oil

Research has shown that consumption of red palm oil significantly enhanced vitamin A levels in humans, and it is beneficial in preventing vitamin A deficiency (Manorama and Rukmini, 1991; Roo, 2000; Solomons, 1998) and it is used for combating vitamin A deficiency in developing countries (Rukmini, 1994). Additionally, some workers have advised that nursing mothers should take red palm oil as supplement with their food in order to prevent vitamin A deficiency (Lietz et al., 2000). Vitamin A deficiency may lead to blindness, skin disease and weakened immune function. The vitamin A content of the red palm oil plays important roles in growth, development and in visual process (Edem, 2009). The human body is able to convert provitamin-A carotenoids (alpha- and beta-carotene) when there is a deficiency, hence it is safer to supplement with carotenes than consuming vitamin A (retinoids) directly. Excessive consumption of retinoids may lead to toxicity with symptoms ranging from mild, such as headache, nausea and dry, itchy skin to severe, such as liver damage (Solomons, 1998).

Tocotrienols are members of the vitamin E family. In the body, vitamin E acts as an anti-oxidant that protects lipid from peroxidation and help quench free radicals. However, there is a difference in anti-oxidant potency between tocotrienol and its sibling tocopherol. Tocotrienol has been shown to be 40 to 60 times more potent than tocopherol as an anti-oxidant. Palm oil

is the only vegetable oil available on the world market that naturally contains tocotrienols (Cottrell, 1991; Ebong et al., 1999; Elson, 1992; Van Rooyen et al., 2008) and is the richest natural source of beta - carotene (500 - 700 mg/l) which is responsible for the characteristic colour of the oil. Similar to the tocopherols, tocotrienols consist of 4 members: alpha, beta, gamma, and delta isomers (Serbinova, 1991). Alpha tocopherols and gamma tocotrienols have anti-oxidative effects on lipid peroxidation, in the presence of a xenobiotic metabolizing enzyme that induces lipid peroxidation (Zuzana et al., 2005). The vitamin E, particularly the tocotrienol present in palm oil can suppress the synthesis of cholesterol in the liver (Mcintosh et al., 1991; Qureshi et al., 1991; Qureshi et al., 1980). Some scientists in South Africa have been able to establish that oxidative stress plays a role in inflammatory and chronic disease such as HIV/AIDS and TB and contribute significantly to depletion of immune factors, micronutrients and progression of disease and that red palm oil could potentially retard the process because of rich anti-oxidants (Oguntibeju et al., 2009).

Anti-diabetic action

Studies have indicated that the potential mechanism of action for the improvement in glucose metabolism with *E. guineensis* involves inhibition of the enzyme dipeptidyl peptidase-4 (DPP-4; Abdullah, 2009) the effect of which is to prevent degradation of gastric inhibitory polypeptide, which itself stimulates insulin secretion, suppresses glucagon secretion and slows gastric emptying. *E. guineensis* is rich in catechins and polyphenols (Jaffri, 2011). Prior studies in streptozotocin-induced hyperglycemic rats showed that *E. guineensis* improved proteinuria and reduced oxidative stress levels (Rosalina Tan, 2011; Yamabe, 2006). This suggests a potential benefit for the pre-diabetic and diabetic states. Administration of tocopherol, Vitamin E or tocotrienol-rich fractions of palm oil, have been shown to recover glycemic status, inhibit oxidative damage; prevent DNA damage in animal studies and prevent glycosylation of end-products in serum and decrease in diabetic rats (Budin et al., 2006; Nazaimoon and Khalid, 2002; Obahiagbon, 2012)

Cardiovascular activities

Peer review journals have documented palm tocotrienol complex's promising hypocholesterolemic properties (Qureshi et al., 1991, 1995). Also, the daily consumption of tocotrienols-enriched fraction of palm oil (200 mg palmvitee capsule) can result in a significant reduction of serum cholesterol, Low Density Lipoprotein cholesterol, APOB, thromboxane, platelet factor 4 and glucose of hyper cholesterolemic subjects within four weeks of administration (Qureshi et al., 1991; Packer, 1992; Song and DeBose-Boyd, 2006). A number of human feeding studies reported that palm oil diets showed a reduction of blood cholesterol values ranging from 7 to 38% (Mattson and Grundy, 1985; Bonanome and Grundy, 1988). A comparative study in young Australian adults showed that the total blood cholesterol, triglycerides and High Density Lipoprotein (HDL) - cholesterol levels of those fed on palm oil (palm olein) and olive oil were lower than those fed on the usual Australian diet (Choudhury et al., 1995). A number of studies have also shown that palm oil increased HDL cholesterol and Apo A-1 levels (Sundram et al., 1992; Truswell et al., 1992). Other studies have also shown the beneficial effects of palm oil on the cholesterol level of the body (Marzuki et al., 1991; Ng et al., 1992, 1991; Zhang et al., 1997). The position of the saturated and unsaturated fatty acid chains in a triglyceride backbone of palm oil molecule determines whether the fat will elevate cholesterol level in the blood

(Kritchevsky, 1988, 1996). High blood pressure or hypertension is one of the major risk factors of cardiovascular diseases and strokes. In a human clinical trial, patients supplemented with palm tocotrienol complex for two months resulted in significant reduction in aortic systolic blood pressure (Rasool et al., 2006). In an earlier review by Obahiagbon (2012), he pointed out that Tocotrienol-rich Fraction (TRF) of palm oil exhibited cardio-protective ability in animal trials (Das et al., 2008). The cardio-protective effects produced by the isomers of tocotrienol were of the order of: $\gamma > \alpha > \delta$. The inhibition of normal cellular gene, C-Src activation and proteasome stabilization were found to be reasons behind the cardio-protective properties of TRF (Das et al., 2008). Feeding experiments using various animal models have highlighted that red palm oil is beneficial to health by reducing oxidative stress (Ebong et al., 1999). Many studies have demonstrated the protective effects of red palm oil in an ischemia/reperfusion model of oxidative stress (Bester et al., 2006; Engelbrecht et al., 2006; Esterhuysen et al., 2005).

Palm oil has been shown to possess anti-clotting effect and it prevents the formation of thrombus in the blood vessels (Oguntibeju et al., 2010). A human study (Kooyenga et al., 1997) showed that tocotrienol (from palm oil) supplementation can reduce stenosis of patients with carotid atherosclerosis. Vitamin E in palm oil has been linked with inhibition of platelets from sticking to each other. Other reports showed that palm oil diets increases the production of prostacyclin or thromboxane (Ng et al., 1992; Sundram et al., 1990). Thus scientific evidence indicates that the palm oil diet is anti-thrombotic. Studies in animals confirmed that palm oil do not promote the formation of plaques in the arteries. A study was conducted on rabbits to test the effect of palm oil on atherosclerosis. After feeding the rabbits for one and a half years, palm oil and sunflower oil diets caused the lowest degree of atherosclerosis in comparison with fish oil, linseed oil and olive oil. Similarly, the effects of palm oil was also compared with other types of plants derived oils and at the end of the 14-month feeding period, coconut oil fed rabbits had the most atherosclerosis lesions, while in palm oil-fed rabbits; the number of lesions was no different from that with the other oils (Oguntibeju et al., 2010).

Anti-bacterial

The anti-bacterial activity of this plant extract against different micro-organisms and anti-oxidant activity have already been reported (Manjunatha, 2005; Sasidharan et al., 2009). Moreover, Chong together with Sasidharan and some others (Chong et al., 2008) described the potential of *E. guineensis* leaf methanol extract as an infected wound healing agents. They observed that the bacterial count in the *E. guineensis* extract treated rats was significantly reduced to 102 CFU/g tissues on day 16. In a further study, Sasidharan tested wound healing activity without infection and the expression of matrix metalloproteinases (Sasidharan et al., 2012).

Toxicity

Several studies have been conducted to confirm the non-toxicity of different parts of *E. guineensis* at normal doses (Rajoo et al., 2010; Syahmi et al., 2010). Syahmi et al. (2010) evaluated the acute oral toxicity of methanol extract of the leaves using a dose of 5g and found no toxicity. They also use the brine shrimp bioassay and also reported no toxicity. Also, Anyanji et al. (2013) showed that the ethanol extract of the palm leaves do not cause any toxicity at 2 g/kg but it appears toxic at 5 g/kg based on a single administration after seven days. However, the indices of toxicity were partially reversed after 14 days of administration.

Conservation and preservation

Several literature have reported the use of palm oil in preservation purposes from various parasites including cowpea weevil, *Callosobruchus maculatus* (Law-Ogbomo and Egharevba, 2006), *Sitophilus zeamais* and *Callosobruchus maculatus* (Abulude et al., 2007).

Anti-cancer action

Studies have shown that tocotrienols fractions of palm oil were able to induce an inhibitory action on the human breast cancer cells, whereas the alpha-tocopherols were not able (Nesaretnam et al., 2004, 2008; McIntyre et al., 2000). Palm oil has been reported to be with wide range of protective properties against disease, aging as well as being modulators for cellular processes / functions where photo oxidative processes predominate by acting as scavengers of oxygen and peroxy radicals (Van Rooyen et al., 2008). It has been shown that fresh palm oil has no adverse effect on body weight and morphology of body tissues, lowers the level of serum lipids and inhibits tumour growth (Kritchevsky, 2000), enhances intestinal uptake of protein and the metabolism of sulphur-amino acids and promotes reproductive capacity (Ebong et al., 1999). Several researches have been conducted on cancer with the view to finding a lasting solution to the disease. Sundram and his colleagues (1989) were able to conduct an experiment to show that RPO significantly reduce tumor incidence in some experimental rats compared with the control groups. RPO, when compared with saturated fats and oils, may help fight cancer, especially breast cancer (Nesaretnam, et al., 1992). This may be due to tocotrienols (Nesaretnam, 1998; Elson, 1992) or other phytonutrients (Guthrie et al., 1997) present in palm oil. Indeed, Professors K. K. Carroll of the Centre for Human Nutrition at the University of Western Ontario and David Kritchevsky of the Wistar Institute recently concluded that evidence from animal and in vitro studies indicate that tocotrienols of palm oil are effective anti-cancer agents and provide adequate justification for clinical trials in human cancer patients (Nesaretnam, 1998). This oil has equally been shown to reduce the incidence of azoxymethane-induced aberrant crypt foci in rats and may therefore have a beneficial effect in reducing the incidence of colon cancer (Boateng et al., 2006).

Anti-inflammatory effects

The TRF of palm oil has been shown by Wu et al. (2008) to possess anti-inflammatory activities in a study involving the injection of lipopolysaccharide-induced inflammatory response. The mediators of cellular inflammation such as nitric oxide (NO), prostaglandin E2, and transcription of pro inflammatory cytokines were significantly reduced. The following were equally blocked; inducible NO, cyclooxygenase 2 expression and NF-kappa B expression.

Leaves of palm oil

A review of the health benefits of the leaf extracts of oil palm has been provided extensively by Mohammed (2014), however, the present review is to provide additional information.

Anti-cancer activities

Methanolic oil palm (*Elaeis guineensis*) leaf extract is rich in polyphenols (Runnie et al., 2003; Jaffri, 2011). The methanolic leaf extract of *E. guineensis* has also been shown to possess anti-cancer activities (Sasidharan and Vijayarathna, 2012). Studies have observed the presence of a large number of bioactive compounds in the methanolic extracts of this plant

including tannins, alkaloids, steroids, saponins, terpenoids, and flavonoids which exhibit various biological activities (Sasidharan et al., 2010; Gulecha and Sivakuma, 2011; Kumar et al., 2011). These compounds are present in a number of food items and hold great potential as drug candidates due to their safety, low toxicity and wide acceptance amongst the public.

Wound healing

Sasidharan et al. (2012) reported in their article that in traditional medicine, the leaf of *E. guineensis* is squeezed and the juice that is obtained is placed on wounds to promote healing (Irvin, 1985). They tried to establish this in an experimental research conducted in Malaysia which showed that *E. guineensis* leaf extract had potent wound healing capacity as evident from the better wound closure ($p < 0.05$), improved tissue regeneration at the wound site, and supporting histopathological parameters pertaining to wound healing (Sasidharan et al., 2012). The wounds were treated topically with 10% formulated crude extract (5 g of the extract in 50 g of yellow soft paraffin), while the control rats were treated only with yellow soft paraffin for 16 days. The decrease in wound diameters during the healing process was measured with an analytical perimeter. The wounded animals were kept for 25 days for further observations (Sasidharan et al., 2012). Although the leaf of the oil palm is a waste product, the alcohol extract of the leaf contains large amounts of phenolic compounds (Sasidharan et al., 2009; Soundararajan and Sreenivasan, 2012) that reportedly promote vascular relaxation and anti-oxidant activity in vitro. A pooled methanolic extract was dried under vacuum using a rotary evaporator and the resultant waxy residue collected, freeze dried, flushed with nitrogen and stored at -20°C . For vascular function studies, a stock solution (100 mg/ml) was prepared using a 1:1 v/v mixture of methanol: saline and serially diluted. Aliquots (25 – 50 μL) were added directly in a cumulative fashion to the bath (aortic rings) or injected intraluminally (mesenteric vascular bed (Abeywardena et al., 2002). In a recent study of streptozotocin (STZ)-induced hyperglycemic rats, *E. guineensis* leaf extract reduced glycemia and lipid oxidation in a dose-dependent manner, possibly by inhibiting dipeptidyl peptidase-4 (DPP-4) secretion (Tan et al., 2011).

Cardiovascular activities

The polyphenol-rich leaf extract of *E. guineensis* showed vasodilative properties on noradrenaline-precontracted rat aorta and mesenteric arterial bed, mainly via endothelium-dependent mechanisms (Abeywardena et al., 2002). It also effectively inhibited low-density lipoprotein oxidation better than other edible plant extracts (Salleh et al., 2002). Also, in a 12-week study conducted by Jaffri et al. (2011) on rats, the leaf extract showed good anti-hypertensive and anti-oxidant effects under NO deficiency, it was not hypotensive to normal rats and produced no chronic cardiovascular toxicity in any of the rats throughout the study.

Hepatoprotective effects

Sasidharan et al. (2009) demonstrated the hepatoprotective effects of *E. guineensis* against paracetamol induced liver damage by looking at the histopathology of mice liver. This was subsequently followed up by a serum analysis in which the same authors (Sasidharan et al., 2012), reported that the methanol extract of the leaves of the plant also offer hepatoprotection against paracetamol induced-liver damage in mice by reducing serum markers of liver injury such aspartate aminotransferase, alanine aminotransferase, and bilirubin.

Anti-diabetic effects

Kalman et al. (2013) reported that the ethanol-derived leaf extract of *E. guineensis* provided a clinically significant, positive effect on fasting plasma glucose levels in individuals with pre-diabetes who were treated with the leaf extract in a dose dependent manner in human subjects. A 500 mg low dose of *E. guineensis* was shown to have had a more consistent effect on reducing glycemia than the higher 1000 mg dose over an eight week period.

Anti-inflammatory activity

Oil palm ethanolic leaf extract (OPLE) at 150 mg/kg body weight showed significant pro-inflammatory activity with enhanced 46% late phase inflammation recovery effects. While at high dose, inflammation was significantly suppressed prior to the sixth hour compared to other groups, and did not require much inflammation suppression between the 18th and 48th hour. OPLE 150 mg/kg decreased lymphocyte counts, but was not as severely as dexamethasone treatment. This result suggests that OPLE extract possess strong in-vivo inflammatory-regulatory effects (Anyanji et al., 2013).

The main problem for the use of oil palm leaf extract as food in its natural form is its high content of insoluble fibre. The OPLE effectively reduced blood glucose and lipid oxidation in Type II diabetic humans and diabetes-induced rodents. The optimum dose in animal studies is equivalent to consuming 5 cups of 1% palm leaf extract for diabetic humans, to prevent liver and kidney damage (Mohammed, 2014).

Sap of oil palm

Multiple actions

The oil palm sap or wine can be described as the exudates that flow when the palm is tapped (Obahiagbon, 2012). The sap of this plant is also used as a laxative and the partially fermented palm wine is administered to nursing mothers to improve lactation (Sasidharan et al., 2012). The sap has also been recorded to be involved in malaria, jaundice and measles treatment. Just like the sap of other palms like *Raphia*, researches conducted by authors like Obahiagbon and Oviasogie (2007), and Obahiagbon et al. (2007) have shown that the sap contains numerous phytonutrients which plays significant roles in human health. Soap prepared with ash from fruit-husk is used for the preparation of a soap used for skin infections (Sasidharan et al., 2012).

Roots of oil palm

Multiple actions

A root decoction is used in Nigeria for headaches. The pulverized roots are added to drinks for gonorrhea, menorrhagia and as a cure for bronchitis (Sasidharan et al., 2012; Irvin, 1985). Chong et al. (2009) demonstrated the in vitro anti-microbial activity and fungitoxicity of syringic acid, caffeic acid and 4-hydroxybenzoic acid which are found in oil palm root. They also showed that of the three substances, syringic acid was the most fungitoxic against *G. boninense*.

CONCLUSION

Oil palm (*E. guineensis*) parts and products have been used in various parts of the world for different purposes. The traditional uses of the different parts of the plant have been well documented in this review. Virtually all the different parts of the plant have one or more therapeutic effects. The plant has

been used extensively in local treatment of various ailments while the scientific uses of the plants have also been documented. The therapeutic effects of this plant have been attributed to the abundant anti-oxidants present in various parts of the plant. However investigations on the actual mechanisms by which the anti-oxidants perform their functions are still ongoing. Also, this review shows that products obtained from oil palm are safe for consumption at moderate doses in humans and rodents.

ACKNOWLEDGEMENTS

The authors are grateful to their respective institutions for the conducive environment provided for this review.

CONFLICT OF INTEREST

The authors declare that there are no conflicts of interest.

REFERENCES

- Abdullah Z, Hussain K, Zhari I, Rasadah MA, Mazura P, Jamaludin F, Sahdan R. Evaluation of extracts of leaf of three *Ficus deltoidea* varieties for antioxidant activities and secondary metabolites. *Phcog Res.* 2009;1:216-223.
- Abeywardena M, Runny I, Nizar M, Mohamed S, Head R. Polyphenol-enriched extract of oil palm fronds (*Elaeis guineensis*) promotes vascular relaxation via endothelium-dependent mechanisms. *Asia Pac J Clin Nutr.* 2002;11:S467-S472.
- Abulude FO, Ogunkoya MO, Ogunleye RF, Akinola AO, Adeyemi AO. Effect of palm oil in protecting stored grains from *Sitophilus zeamais* and *Callosobruchus maculatus*. *J Entomol.* 2007;4:393-396.
- Anyanji VU, Mohamed S, Zokti JA, Ado MA. Anti-inflammatory properties of oil palm leaf (*Elaeis guineensis* Jacq.) extract in aged rats. *Int J Pharm Pharm Sci.* 2013;5:134-136.
- Bester DJ, van Rooyen J, Du Toit EF, Esterhuyse AJ. Red palm oil protects against the consequences of oxidative stress when supplemented with dislipidaemic diets. *Med Tech SA.* 2006;21:3-6.
- Boateng J, Verghese M, Chawan CB, Shackelford L, Walker LT, Khatiwada J, Williams DS. Red palm oil suppresses the formation of azoxymethane (AOM) induced aberrant crypt foci (ACF) in fisher 344 male rats. *Food Chem Toxicol.* 2006;44:1667-1673.
- Bonanome A, Grundy SM. Effect of dietary stearic acid and plasma cholesterol and lipoprotein levels. *New Eng J Med.* 1988;318:1244-1248.
- Bauer AW, Kirby WM, Sherris JC, Turck M. Antibiotic susceptibility testing by a standardized single disk method. *Am J Clin Pathol.* 1966;45:493-496.
- Budin SB, Rajab NF, Osman K, Top AGM, Mohammed WNW, Bakar MA, Mohamed J. Effects of palm vitamin E against oxidative damage in *Streptozotocin*-induced diabetic rats. *Malaysian J Biochem Mol Biol.* 2006;13:11-17.
- Chong KP, Rossall S, Atong M. In vitro antimicrobial activity and fungitoxicity of syringic acid, caffeic acid and 4-hydroxybenzoic acid against *Ganoderma boninense*. *J Agric Sci.* 2009;1:15-20.
- Chong KH, Zuraini Z, Sasidharan S, Devi PVK, Latha LY, Ramanathan S. Antimicrobial of *Elaeis guineensis* leaf. *Pharmacologyonline.* 2008;3:379-386.
- Chong YH, Ng TK. Effects of palm oil on cardiovascular risk. *Med J Malaysia.* 1991;46:41-50.
- Choudhury N, Tan L, Truswell AS. Comparison of palm olein and olive oil: Effects on plasma lipids and Vitamins E in young adults. *Am J Clin Nutr.* 1995; 61:1043-1051.
- Corley RHV. Oil Palm Research. In the genus *Elaeis*, Corley RHV, Hardon JJ, Wood BJ ed. (Amsterdam, Netherlands: Elsevier Scientific Publisher), p. 532, 1976.
- Cottrell RC. Introduction: nutritional aspects of palm oil. *Amer J Clin Nutr.* 1991;53:989S-1009S.
- Das S, Lekli I, Das M, Szabo G, Varadi J. Cardioprotection with palm oil tocotrienols: Comparison of different isomers. *Am J Physiol Heart Circ Physiol.* 2008;294:70-78.
- Duke AJ. Handbook of Energy Crops. pp. 1-5, 1983. Available at: https://www.hort.purdue.edu/newcrop/duke_energy/Elaeis_guineensis.html (accessed on 01st May 2014).
- Ebong PE, Owu DU, Isong EU. Influence of palm oil (*Elaeis guineensis*) on health. *Plant Food Hum Nutr.* 1999;53:209-222.
- Edem DO. Palm oil: Biochemical, physiological, nutritional, hematological and toxicological aspects: a review. *Plant Foods Hum Nutr.* 2002;57:319-341.
- Edem DO. Vitamin A: A review. *Asian J Clin Nutr.* 2009;1:65-82.
- Edem DO, Akpanabiatu MI. Effects of palm oil-containing diets on enzyme activities of rats. *Pak J Nut.* 2006;5:301-305.
- Ekpa OD, Ebana RUB. Comparative Studies of Mmanyanga, Palm and Coconut Oils; Anti-microbial Effects of the Oils and their Metallic Soaps on Some Bacteria and Fungi. *Global J Pure and Appl Sci.* 1996;2:155-163.
- Ekwenye UN, Ijeomah CA. Antimicrobial effects of palm kernel oil and palm oil. *KMITL Sci J.* 2005;5:502-505.
- Elson CE. Tropical Oils: Nutritional and Scientific Issues. *Crit Rev Food Sci Nutr.* 1992;31:79-102.
- Engelbrecht AM, Esterhuyse AJ, Du Toit EF, Lochner A, Van Rooyen J. P38-MAPK and PKB/Akt, possible role players in red palm oil-induced protection of the isolated perfused rat heart? *J Nutr Biochem.* 2006;17:265-271.
- Esterhuyse AJ, Du Toit EF, Van Rooyen J. Dietary red palm oil supplementation protects against the consequences of global ischemia in the isolated perfused rat heart. *Asia Pac J Clin Nutr.* 2005;14:340-347.

- Goh SH, Khor HT, Gee PT. Phospholipids of palm oil (*Elaeis guineensis*). J Am Oil Chem Soc. 1982;59:296-299.
- Goh SH, Choo YM, Ong SH. Minor constituents of palm oil. J Am Oil Chem Soc. 1985;62:237-240.
- Gulecha V, Sivakuma T. Anticancer activity of *Tephrosia purpurea* and *Ficus religiosa* using MCF 7 cell lines. Asian Pac J Trop Med. 2011;4:526-529.
- Guthrie N, Gapor A, Chambers AF, Carroll KK. Palm oil tocotrienols and plant flavonoids act synergistically with each other and with tamoxifen in inhibiting proliferation and growth of estrogen receptor-negative MDA-MB-435 and -positive MCF-7 human breast cancer cells in culture. Asia Pac J Clin Nutr. 1997;6:41-45.
- Hartley CWS. The Oil Palm. 1st ed. (London, England: Longmans), p. 618, 1977.
- Honstra G. Beneficial Effects of Palm oil on Arterial Thrombosis (Rat) and Atherosclerosis (Rabbit). (Kuala Lumpur, Malaysia: PORIM), pp. 35-45, 1986.
- Irvin TT. Wound healing. Arch Emerg Med. 1985;2:3-10.
- Jaffri JM, Mohamed S, Rohimi N, Ahmad IN, Noordin MM, Manap YA. Antihypertensive and cardiovascular effects of catechin-rich oil palm (*Elaeis guineensis*) leaf extract in nitric oxide-deficient rats. J Med Food. 2011;14:775-783.
- Kalman DS, Schwartz HI, Feldman S, Krieger DR. Efficacy and safety of *Elaeis guineensis* and *Ficus deltoidea* leaf extracts in adults with pre-diabetes. Nutr J. 2013;12:36.
- Kooyenga DK, Gerler M, Watkins TR, Gapor A, Diakoumakis E, Bierenbaum ML. Palm oil antioxidant effects in patients with hyper-lipidaemia and carotid stenosis-2 year experience. Asia Pac J Clin Nutr. 1997;6:72-75.
- Krichevsky D. Effects of triglyceride structure on lipid metabolism. J Nutr Rev. 1988;46:177-181.
- Kritchevsky D, Tepper SA, Chen SC, Meijer GW. Influence of triglyceride structure on experimental atherosclerosis in rabbits. FASEB J. 1996;10:A187.
- Kritchevsky D. Impact of red palm oil on human nutrition and health. Food Nutr Bull. 2000;21:182-188.
- Kumar RS, Rajkapoor B, Perumal P. In vitro and in vivo anticancer activity of *Indigofera cassioides* Rottl. Ex. DC. Asian Pac J Trop Med. 2011;4:379-385.
- Law-Ogbomo KE, Egharevba RKA. The use of vegetable oils in the control of *Callosobruchus maculatus* (F) (Coleoptera: Bruchidae) in three cowpea varieties. Asian J Plant Sci. 2006;5:547-552.
- Lietz G, Henry CJK, Mulokozi G, Mugyabuso J, Ballart A. Use of red palm oil for the promotion of maternal vitamin A status. Food Nutr Bull. 2000;21:215-218.
- Manorama R, Rukmini C. Nutritional evaluation of crude palm oil in rats. Am J Clin Nutr. 1991;53:1031-1033.
- Marzuki A, Arshad F, Razak TA, Jaarin K. Influence of dietary fat on plasma lipid profiles of Malaysian adolescents. Am J Clin Nutr. 1991;53:1010-1014.
- Mattson FH, Grundy SM. Comparison of effects of dietary saturated, monounsaturated and polyunsaturated fatty acids on plasma lipids and lipoproteins in man. J Lipid Res. 1985;26:194-202.
- McIntosh GH, Whyte J, McArthur R., Nestle, P. J. Barley and wheat foods; Influence on plasma cholesterol concentrations in hypercholesterolemic men. Am J Clin Nutr. 1991;53:1205-1209.
- McIntyre BS, Briski KP, Gapor A, Sylvester PW. Antiproliferative and apoptotic effects of tocopherols and tocotrienols on preneoplastic and neoplastic mouse mammary epithelial cells. Proc Soc Exp Biol Med. 2000;224:292-301.
- Mohamed S. Oil Palm Leaf: A New Functional Food Ingredient for Health and Disease Prevention. J Food Process Technol. 2014;5:300.
- Naher L, Yusuf UK, Ismail A, Tan SG, Mondal MMA. Ecological status of Ganoderma and basal stem rot disease of oil palms (*Elaeis guineensis* Jacq.) AJCS. 2013;7:1723-1727.
- Nazaimoon WMW, Khalid BA. Tocotrienols-rich diet decreases advanced glycosylation end products in non-diabetic rats and improves glycemic control in Streptozotocin-induced diabetic rats. Malays J Pathol. 2002;24:77-82.
- Nesaretnam K. Tocotrienols inhibit the growth of human breast cancer cells irrespective of estrogen receptor status. Lipids. 1998;33:461-469.
- Nesaretnam K, Khor HT, Ganeson J, Chong K, Gapor A. The effect of vitamin E tocotrienols from palm oil on chemically-induced mammary carcinogenesis in female rats. Nutr Res. 1992;12:879-892.
- Nesaretnam K, Ambra R, Selvaduray KR, Radhakrishnan A, Reimann K, Razak G, Virgili F. Tocotrienol-rich fraction from palm oil affects gene expression in tumors resulting from MCF-7 cell inoculation in athymic mice. Lipids. 2004;39:459-467.
- Ng TKW, Hayes KC, De Witt GF, Jegathesan M, Satgunasingham N, Ong ASH, Tan DTS. Dietary palmitic and oleic acids exert similar effects on serum cholesterol and lipoprotein profiles in normocholesterolemic men and women. J Am Coll Nutr. 1992;11:383-390.
- Ng TKW, Hassan K, Lim JB, Lye MS, Ishak R. Nonhypercholesterolemic effects of a palm oil diet in Malaysian volunteers. Am J Clin Nutr. 1991;53:1015S-1020S.
- Obahiagbon FI. A Review: Aspects of the African Oil Palm (*Elaeis guineensis* Jacq.) and the Implications of its Bioactives in Human Health. Amer J Biochem Mol Biol. 2012;2:106-119.
- Obahiagbon FI, Oviasogie PO, Ukhun ME, Oviawe PN. Comparative studies of the thiamine and riboflavin contents in the sap of *Raphia hookeri* palm by the application of fluorimetric method and arrhenius equation. Tren Appl Sci Res. 2007;2:260-263.

- Obahiagbon FI, Oviasogie PO. Changes in the physicochemical characteristics of processed and stored *Raphia hookeri* palm sap (shelf-life studies). *Am J Food Technol*. 2007;2:323-326.
- Oguntibeju OO, Esterhuysen AJ, Truter EJ. Possible role of red palm oil supplementation in reducing oxidative stress in HIV/AIDS and TB patients: A review. *J Med Pl Res*. 2010;4:188-196.
- Oluba OM, Onyeneke CE, Ojeh GC, Eidangbe GO, Orole RT. Effects of Palm Oil Supplementation on Lipid Peroxidation and Glutathione Peroxidase Activity in Cholesterol – Fed Rats. *The Internet J Cardio Res*. 2009;6:1.
- Packer L. Interactions among antioxidants in health and disease: Vitamin E and its redox cycle. *Exp Biol Med*. 1992;200:271-276.
- Packer L, Serbinova E, Choo M. Distribution and antioxidant activity of a palm oil carotene in rats. *Biochem Inter*. 1992;28:881-886.
- Qureshi AA, Qureshi N, Wright JJ, Shen Z, Kramer G. Lowering of serum cholesterol in hypercholesterolemic humans by tocotrienols (palmvite). *Am J Clin Nutr*. 1991;53:1021-1026.
- Qureshi A, Burger WC, Prentice N, Bird HR, Sunde ML. Regulation of Lipid metabolism in chicken liver by dietary cereals. *J Nutr*. 1980;110:388-393.
- Qureshi AA, Pearce BC, Parker RA, Deason ME. Response of hypercholesterolemic subjects to administration of tocotrienols. *Lipids*. 1995;30: 1171-1177.
- Rajoo A, Ramanathan S, Sasidharan S, Mansor SM. Standardization of *Elaeis guineensis* with respect to authenticity, assay and chemical constituent analysis. *Afr J Biotechnol*. 2010;9:7544-7549.
- Rasool AHG, Yuen KH, Yusoff K, Wong AR, Rahman AR. Dose dependent elevation of plasma tocotrienol levels and its effect on arterial compliance, plasma total antioxidant status, and lipid profile in healthy humans supplemented with tocotrienol rich vitamin E. *J Nutr Sci Vitaminol*. 2006;52:473-478.
- Rao BSN. Potential use of red palm oil in combating vitamin A deficiency in India. *Food Nutr Bull*. 2000;21:202-211.
- Rosalina Tan RT, Mohamed S, Samaneh GF, Noordin MM, Goh YM, Manap MYA. Polyphenol rich oil palm leaves extract reduce hyperglycaemia and lipid oxidation in STZ-rats. *Int Food Res J*. 2011;18:179-188.
- Rukmini C. Red palm oil to combat vitamin A deficiency in developing countries. *Food Nutr Bull* 1994;15:126-129.
- Runnie I, Mustapha NM, Radzali M, Azizah H, Hapizah N, Abeywardena MY, Mohamed S. Antioxidant and hypocholesterolemic effects of *Elaeis guineensis* frond extract on hypercholesterolemic rabbits. *Asean Food J*. 2003;12:137-147.
- Salleh MN, Runnie I, Roach PD, Mohamed S, Abeywardena MY. Inhibition of low-density lipoprotein oxidation and up-regulation of low-density lipoprotein receptor in HepG2 cells by tropical plant extracts. *J Agric Food Chem*. 2002;50:3693-3697.
- Sasidharan S, Vijayarathna S. Cytotoxicity of methanol extracts of *Elaeis guineensis* on MCF-7 and Vero cell lines. *Asian Pac J Trop Biomed*. 2012;2:826-829.
- Sasidharan S, Nilawaty R, Xavier R, Yoga Latha L, Amala R. Wound healing potential of *Elaeis guineensis* Jacq leaves in an infected albino rat model. *Molecules*. 2010;15:3186-3199.
- Sasidharan S, Selvarasoo L, Latha LY. Wound Healing Activity of *Elaeis guineensis* Leaf Extract Ointment. *Int J Mol Sci*. 2012;13:336-347.
- Sasidharan S, Vijayarathna S, Jothy SL, Ping KY, Latha LY. Hepatoprotective potential of *Elaeis guineensis* leaf against paracetamol induced damage in mice: A serum analysis. *IPCBE*. 2012;39:231-234.
- Sasidharan S, Sharmini R, Vijayarathna S, Yoga Latha L, Vijenthir R, Amala R, Amutha S. Antioxidant and hepatoprotective activity of methanolic extracts of *Elaeis Guineensis* Jacq leaf. *Pharmacologyonline*. 2009;3:84-90.
- Serbinova E, Kagan V, Han D, Packer L. Free Radical Recycling and Intramembrane Mobility in the Antioxidant properties of Alpha-tocopherol and Alpha-tocotrienol. *Free Radic Biol Med*. 1991;10:263-275.
- Solomons NW. Plant sources of vitamin A and human nutrition: red palm oil does the job. *Nutr Rev*. 1998;56:309-311.
- Song BL, DeBose-Boyd RA. Insig-dependent ubiquitination and degradation of 3-hydroxy-3-methylglutaryl coenzyme A reductase stimulated by δ - and γ -tocotrienols. *J Biol Chem*. 2006;281:25054-25061.
- Soundararajan V, Sreenivasan S. Antioxidant activity of *Elaeis guineensis* leaf extract: An alternative nutraceutical approach in impeding aging. *APCBEE Procedia*. 2012;2:153-159.
- Sundram K, Khor HT, Ong SHA, Pathmanathan R. Effect of dietary palm oils on mammary carcinogenesis in female rats induced by 7, 12-dimethylbenz (a) anthra-cene. *Cancer Res*. 1989;49:1447-1451.
- Sundram K, Hornstra G, Houwelingen AC. Replacement of dietary fat with palm oil: Effect on human serum lipid, lipoprotein and apoproteins. *Br J Nutr*. 1992;68:677-692.
- Sundram K, Khor HT, Ong ASH. Effect of dietary palm oil and its fractions on rat plasma and high density lipoprotein lipids. *Lipids*. 1990;25:187-193.
- Syahmi ARM, Vijayarathna S, Sasidharan S, Latha LY, Kwan YP, Lau YL, Shin LN, Chen Y. Acute oral toxicity and brine shrimp lethality of *Elaeis guineensis* jacq., (oil palm leaf) methanol extract. *Molecules*. 2010;15:8111-8121.
- Tan RT, Mohamed S, Samaneh GF, Noordin MM, Goh YM, Manap MYA. Polyphenol rich oil palm leaves extract reduce hyperglycaemia and lipid oxidation in STZ-rats. *Int Food Res J*. 2011;18:179-188.
- Truswell AS, Choudhury N, Roberts DCK. Double blind comparison of plasma lipids in healthy subjects eating potato

crisps fried in palmolein or canola oil. *Nutr Res.*1992;12:S34-S52.

Van Rooyen J, Esterhuyse AJ, Engelbrecht A, Toit EF. Health benefits of natural carotenoid rich oil: a proposed mechanism of protection against ischemia reperfusion injury. *Asia Pac J Clin Nutr.* 2008;17:316-319.

Wu SJ, Liu PL, Ng LT. Tocotrienol-rich fraction of palm oil exhibits anti-inflammatory property by suppressing the expression of inflammatory mediators in human monocytic cells. *Mol Nutr Food Res.* 2008;52:921-929.

Yamabe N, Yokozawa T, Oya T, Kim M. Therapeutic potential of (–)-epigallocatechin 3-*O*-gallate on renal damage in diabetic nephropathy model rats. *J Pharmacol Exp Ther.* 2006;319:228-236.

Zhang J, Wang CR, Dai J, Chen XS, Ge KY. Palm oil diet may benefit mildly hypercholesterolaemic Chinese adults. *Asia Pac J Clin Nutr.* 1997;6:22-25.

Zuzana M, Suhana M, Zalinah A, Gapor MT, Ngah WZW. Comparative effects of α -tocopherol and γ tocotrienol on lipid peroxidation status in Hep G2 cell line transfected with CYZE1 gene. *Eur J Sci Res.*2005;7:41-56.