

Review

The role of thymic stromal lymphopoietin on mast cell-mediated allergic inflammatory reactions

Tae-Yong Shin*

College of Pharmacy, Woosuk University, Wanju-gun, Jeonbuk, 55338, Republic of Korea

ABSTRACT

Thymic stromal lymphopoietin (TSLP) is a novel interleukin (IL)-7-like cytokine and was originally discovered in the supernatant of a murine thymic stromal cell line. TSLP signal initiates via complex of the TSLP receptor and the IL-7 receptor α chain. TSLP expression is closely connected with many diseases such as atopic dermatitis, allergic rhinitis, asthma, inflammatory arthritis, eosinophilic esophagitis, rheumatoid arthritis, inflammatory bowel diseases, and cancer. In this review, I discussed biological roles of TSLP on mast cell-mediated allergic responses. In addition, this review summarizes the effective drugs in allergic-inflammatory reactions induced by TSLP on mast cells.

Keywords thymic stromal lymphopoietin, atopic dermatitis, allergic rhinitis, mast cells

INTRODUCTION

1. Mast cells

Mast cells play a critical role in the initiation and propagation of allergic inflammatory processes such as, allergic asthma, life-threatening anaphylaxis, inflammatory response, atopic dermatitis (AD), and allergic rhinitis (AR) (El-Agamy, 2012; Galli and Tsai, 2012). The cross-linking of IgE triggers mast cell de-granulation and the synthesis of a variety of mediators including biogenic amines, serglycin proteoglycans, serine proteases, cytokines, and growth factors (Plaut et al., 1989). Histamine released by mast cells evoked vascular permeabilization and hence enhances the migration of eosinophils, macrophages, and neutrophils into inflamed tissues (Oskeritzian et al., 2015). Thus, mast cell degranulation and activations intensifies and extends TH2-dependent inflammatory response (Galli and Tsai, 2012). Furthermore, interleukin (IL)-3, IL-5, IL-9, IL-6, tumor necrosis factor (TNF)- α , macrophage inflammatory protein-2, intercellular adhesion molecule-1, and thymic stromal lymphopoietin (TSLP) induce the recruitment of inflammatory cells leading to the late allergic response expressed by nasal congestion, chronic wheeze, and nasobronchial hyper-reactivity (Mandhane, 2011; Moon and Kim, 2011; Perlman, 1999; Xu et al., 1995). Intracellular signaling pathways such as mitogen-activated kinases and caspase-1 regulate the production of these pro-inflammatory cytokines (Oh et al., 2012). The mRNA expressions of IL-1, IL-6, IL-8, TNF- α , and TSLP are induced by activation of a transcription factor, nuclear factor (NF)- κ B. NF- κ B bound to a specific consensus DNA element of promoter region increases the transcription of inflammatory cytokine genes (Jeong et al., 2002).

Mast cells are also multifunctional immune cells involved

in the pathogenesis of various chronic inflammatory disorders, autoimmune diseases, and cancers (Galli and Tsai, 2012). Increased number of mast cells has been found in AD, AR, asthma, rheumatoid arthritis, tumor, and mast cell leukemia (El-Agamy, 2012; Galli and Tsai, 2012; Jeong et al., 2013). Mastocytosis is disorders determined by abnormal proliferation and accumulation of mast cells in various tissues, including liver, skin, gastrointestinal tract, bone marrow, spleen, and lymph nodes (Horny et al., 2008). The number of mast cells is increased by migration, proliferation, and survival (Kneilling and Röcken, 2009). Proliferation of mast cells was mainly induced by stem cell factor and IL-3 (da Silva et al., 2014). Recently, we have been demonstrated that TSLP also increased the proliferation of mast cells (Han et al., 2014a). Proliferation of mast cells exaggerates the several inflammatory diseases. Thus, inhibition of mast cell proliferation is regarded as an attractive therapeutic strategy for various inflammatory diseases.

2. TSLP

TSLP is a novel interleukin (IL)-7-like cytokine and was originally discovered in the supernatant of a murine thymic stromal cell line (Friend et al., 1994; Sims et al., 2000). TSLP has been known to promote Th2 cell-associated inflammation. TSLP is mostly expressed by epithelial cells of barrier surfaces in response to inflammatory cytokines, microbial products, or physical injury (Allakhverdi et al., 2007). The TSLP receptor (TSLPR) complex consists of a heterodimer of the IL-7 receptor α chain (IL-7R α) and TSLPR (Pandey et al., 2000; Park et al., 2000). TSLP signaling is initiated by heterodimerization of the IL-7R α and TSLPR (Pandey et al., 2000; Park et al., 2000). TSLP affects immune responses of nonhematopoietic and hematopoietic cell lineages, including mast cells, dendritic cells, natural killer, CD4⁺, and CD8⁺ T cells, basophils, eosinophils, epithelial cells, and B cells (Ziegler et al., 2013). TSLP is involved in the initiation and progression of numerous disorders including skin disorders, respiratory diseases, intestine inflammation, cancer, and autoimmune diseases (Ziegler et al., 2013). Mast cell activation via the TSLP produced by epithelial cells has a central function in allergic responses and it exacerbates infection and

*Correspondence: Tae-Yong Shin

E-mail: tyshin@woosuk.ac.kr

Received July 14, 2016; **Accepted** August 16, 2016; **Published** August 31, 2016

doi: <http://dx.doi.org/10.5667/tang.2016.0021>

© 2016 by Association of Humanitas Medicine

This is an open access article under the CC BY-NC license.

(<http://creativecommons.org/licenses/by-nc/3.0/>)

Table 1. An inhibitory effect of drugs on TSLP production

Cell/Animal models	Drugs	Reference
HMC-1 cell	2-(4-{2-[(phenylthio)acetyl]carbonohydrzonoyl}phenoxy)acetamide	Moon et al., 2016
DNFB-induced atopic dermatitis animal model(BALB/c mice)	Resveratrol	Caglayan et al., 2016
HMC-1 cell	Alumium-doped zinc oxide nanoparticle	Kim et al., 2016
HMC-1 cell	Hydrogen sulfide	Han et al., 2016a
HMC-1 cell	Cordycepin	Yoo et al., 2016a
DNFB-induced atopic dermatitis animal model(NC/Nga mice)	<i>Acer tegmentosum</i> , <i>Salidroside</i>	Yang et al., 2016
DNFB-induced atopic dermatitis animal model(NC/Nga mice), HMC-1 cell	Niram (natural dye)	Han et al., 2014b
DNFB-induced atopic dermatitis animal model(NC/Nga mice), HMC-1 cell	Beta-sitosterol	Han et al., 2014c
HMC-1 cell	Hyperoside	Han et al., 2014d
DNFB-induced atopic dermatitis animal model(NC/Nga mice), HMC-1 cell, Splenocyte	Tryptanthrin	Han et al., 2014e
DNFB-induced atopic dermatitis animal model(NC/Nga mice), HMC-1 cell, Splenocyte, Allergic rhinitis animal model	Naju jjok	Han et al., 2014f Jeong et al., 2014
THP-1 (human monocyte)	Bamboo salt	Nam et al., 2014
Allergic rhinitis animal model, Eo1 (eosinophil cell line)	Kaempferol	Oh et al., 2013a
Allergic rhinitis animal model, HMC-1	Ginsenoside Rg1	Oh et al., 2013b
DNFB-induced atopic dermatitis animal model(NC/Nga mice)	<i>Catalpa ovata</i>	Yang et al., 2013
HMC-1 cell	Curcumin	Moon et al., 2013
THP-1 (human monocyte)	BaekJeol-Tang, Chondroitin sulfate	Jeong et al., 2012
HMC-1 cell	Schizandrin	Moon et al., 2012a
HMC-1 cell	BAPTA-AM (calcium chelator)	Han et al., 2012
HMC-1 cell	Selenium	Moon et al., 2012b
HMC-1 cell	Epigallocatechin-3-O-gallate (EGCG)	Moon et al., 2012c
HMC-1 cell	Berberine	Moon et al., 2011a
HMC-1 cell	Naringenin	Moon et al., 2011b

inflammation (Allakhverdi et al., 2007). TSLP is expressed and released by NF- κ B and caspase-1 in activated mast cells (Moon and Kim, 2011). In addition, TSLP stimulation of mast cells leads to signal transducers and activators of transcription (STAT6) phosphorylation, murine double minute 2 (MDM2) expression, and IL-13 production and then increased mast cell-mediated allergic inflammatory reaction (Han et al., 2014a).

TSLP plays a pivotal role as a chief regulator of allergic inflammation in mice and human and is a new target for anti-allergic inflammatory therapy. In this review, we will focus on TSLP and describe how TSLP plays a role in development of allergic inflammation, and discuss TSLP as an attractive therapeutic target for allergic diseases.

2.1. TSLP in AD

TSLP was highly overexpressed by keratinocytes in the skin lesions of AD patients and mice (Soumelis et al., 2002; Corrigan et al., 2009). It causes the initiation of allergic inflammation by stimulating activation and infiltration of Langerhans cells into skin-draining lymph nodes (Soumelis et al., 2002; Zhang and Zhou, 2012). TSLP was also important for local antigen-induced Th2 cytokine production by directly acting on skin-infiltrating antigen-specific CD4⁺ T cells (He et al., 2008; Zhang and Zhou, 2012). Consistent with a requirement for TSLP in AD, challenged TSLPR deficient mice showed significantly decreased allergic skin inflammation, decreased eosinophils infiltration, and decreased Th2 cytokines expression compared with wild type mice in the skin (He et al., 2008). TSLP-deficient mice failed to generate AD development

through the down-regulation of STAT6 and MDM2 (Han et al., 2014a).

2.2. TSLP in AR

TSLP is also critical factor for the pathophysiology of AR. In nasal epithelial cells, expression of TSLP was significantly higher in AR patients than in normal controls (Mou et al., 2008; Zhu et al., 2009; Kimura et al., 2011). TSLP level in nasal tissue biopsies was closely associated with IL-4 levels and AR severity (Mou et al., 2008). Highly TSLP levels were detected in nasal polyps of AR patients (Kimura et al., 2011; Liu et al., 2011).

3. Anti-allergic inflammatory effects of drugs via blockade of TSLP function

TSLP as a developmental factor of AD and AR was released and expressed by the intracellular calcium/caspase-1/ receptor-interacting protein 2/NF- κ B pathways on mast cells (Han et al., 2012; Moon and Kim, 2011). Overexpressed-TSLP deteriorates the pathogenesis of AD and AR via the beginning of allergic inflammatory reactions (Han et al., 2014a; Oh et al., 2013a). TSLP stimulation of mast cells leads to STAT6 phosphorylation, murine double minute 2 expression, and IL-13 production and then increased mast cell-mediated allergic inflammatory reaction (Han et al., 2014a). Therefore, TSLP may be a useful therapeutic target for mast cell-mediated allergic diseases. Based on the inhibitory effect of TSLP production reported, drugs were shown in Table 1. Anti-proliferative effects of drugs on TSLP-induced the mast cell proliferation were also shown in

Table 2. An inhibitory effect of drugs on TSLP-induced mast cell proliferation

Cell/Animal models	Drugs	Reference
HMC-1 cells	Tryptanthrin	Han et al., 2016b
HMC-1 cell (mast cell), HaCa T cell (keratinocyte), pollen-induced allergic conjunctivitis mouse model.	Rosmarinic acid	Yoou et al., 2016b
HMC-1 cells	Beta-sitosterol	Han et al., 2015
HMC-1 cells	Acteoside	Yoou et al., 2015
HMC-1 cells	Zinc oxide nanoparticle	Kim and Jeong 2016

Table 2.

CONCLUSION

The role of cytokines towards inflammation and modulation of various allergic diseases is a hot topic today. TSLP is an important factor in the pathogenesis of various allergic diseases and involved in the initiation and progression of allergic response. TSLP also induces the expression of a cascade of various potent inflammatory cytokines. Therefore, I suggested that TSLP is an attractive therapeutic target for allergic inflammatory diseases.

ACKNOWLEDGEMENTS

None

CONFLICT OF INTEREST

The authors state no conflict of interest.

REFERENCES

Allakhverdi Z, Comeau MR, Jessup HK, Yoon BR, Brewer A, Chartier S, Paquette N, Ziegler SF, Sarfati M, Delespesse G. Thymic stromal lymphopoietin is released by human epithelial cells in response to microbes, trauma, or inflammation and potently activates mast cells. *J Exp Med*. 2007;204:253-258.

Caglayan Sozmen S, Karaman M, Cilaker Micili S, Isik S, Arikan Ayyildiz Z, Bagriyanik A, Uzuner N, Karaman O. Resveratrol ameliorates 2,4-dinitrofluorobenzene-induced atopic dermatitis-like lesions through effects on the epithelium. *PeerJ*. 2016;4:e1889.

Corrigan CJ, Jayaratnam A, Wang Y, Liu Y, de Waal Malefyt R, Meng Q, Kay AB, Phipps S, Lee TH, Ying S. Early production of thymic stromal lymphopoietin precedes infiltration of dendritic cells expressing its receptor in allergen-induced late phase cutaneous responses in atopic subjects. *Allergy*. 2009;64:1014-1022.

da Silva EZ, Jamur MC, Oliver C. Mast cell function: a new vision of an old cell. *J Histochem Cytochem*. 2014;62:698-738.

El-Agamy DS. Anti-allergic effects of nilotinib on mast cell-mediated anaphylaxis like reactions. *Eur J Pharmacol*. 2012; 680:115-121.

Friend SL, Hosier S, Nelson A, Foxworthe D, Williams DE, Farr A. A thymic stromal cell line supports in vitro development of surface IgM+ B cells and produces a novel

growth factor affecting B and T lineage cells. *Exp Hematol*. 1994;22:321-328.

Galli SJ, Tsai M. IgE and mast cells in allergic disease. *Nat Med*. 2012;18: 693-704.

Han NR, Go JH, Kim HM, Jeong HJ. Hyperoside regulates the level of thymic stromal lymphopoietin through intracellular calcium signalling. *Phytother Res*. 2014d;28:1077-1081.

Han NR, Kang SW, Moon PD, Jang JB, Kim HM, Jeong HJ. Genuine traditional Korean medicine, Naju Jjok (*Chung-Dae, Polygonum tinctorium*) improves 2,4-dinitrofluorobenzene-induced atopic dermatitis-like lesional skin. *Phytomedicine*. 2014f;21:453-460.

Han NR, Kim HM, Jeong HJ. The potential anti-proliferative effect of β -sitosterol on human mast cell line-1 cells. *Can J Physiol Pharmacol*. 2015;93:979-983.

Han NR, Kim HM, Jeong HJ. The β -sitosterol attenuates atopic dermatitis-like skin lesions through down-regulation of TSLP. *Exp Biol Med* (Maywood). 2014c;239:454-64

Han NR, Kim HM, Jeong HJ. Thymic stromal lymphopoietin is regulated by the intracellular calcium. *Cytokine*. 2012;59:215-217.

Han NR, Kim HM, Jeong HJ. Tryptanthrin reduces mast cell proliferation promoted by TSLP through modulation of MDM2 and p53. *Biomed Pharmacother*. 2016b;79:71-77

Han NR, Moon PD, Jeong HJ, Kim HM. Hydrogen sulfide diminishes the levels of thymic stromal lymphopoietin in activated mast cells. *Arch Dermatol Res*. 2016a;308:103-113

Han NR, Moon PD, Kim HM, Jeong HJ. Tryptanthrin ameliorates atopic dermatitis through down-regulation of TSLP. *Arch Biochem Biophys*. 2014e;542:14-20

Han NR, Oh HA, Nam SY, Moon PD, Kim DW, Kim HM, Jeong HJ. TSLP induces mast cell development and aggravates allergic reactions through the activation of MDM2 and STAT6. *J Invest Dermatol*. 2014a;134:2521-2530

Han NR, Park JY, Jang JB, Jeong HJ, Kim HM. A natural dye, Niram improves atopic dermatitis through down-regulation of TSLP. *Environ Toxicol Pharmacol*. 2014b;38:982-990.

He R, Oyoshi MK, Garibyan L, Kumar L, Ziegler SF, Geha RS. TSLP acts on infiltrating effector T cells to drive allergic skin inflammation. *Proc Natl Acad Sci USA*. 2008;105:11875-11880.

Horny HP, Sotlar K, Valent P, Hartmann K. Mastocytosis: a

- disease of the hematopoietic stem cell. *Dtsch Arztebl Int*. 2008;105:686-692.
- Jeong HJ, Koo HN, Na HJ, Kim MS, Hong SH, Eom JW, Kim KS, Shin TY, Kim HM. Inhibition of TNF-alpha and IL-6 production by Aucubin through blockade of NF-kappaB activation RBL-2H3 mast cells. *Cytokine*. 2002;18:252-259.
- Jeong HJ, Nam SY, Oh HA, Han NR, Kim YS, Moon PD, Shin SY, Kim MH, Kim HM. Interleukin-32-induced thymic stromal lymphopoietin plays a critical role in macrophage differentiation through the activation of caspase-1 in vitro. *Arthritis Res Ther*. 2012;14:R259
- Jeong HJ, Oh HA, Lee BJ, Kim HM. Inhibition of IL-32 and TSLP production through the attenuation of caspase-1 activation in an animal model of allergic rhinitis by *Naju Jjok* (*Polygonum tinctorium*). *Int J Mol Med*. 2014;33:142-150.
- Jeong HJ, Oh HA, Nam SY, Han NR, Kim YS, Kim JH, Lee SJ, Kim MH, Moon PD, Kim HM, Oh HA. The critical role of mast cell-derived hypoxia-inducible factor-1 α in human and mice melanoma growth. *Int J Cancer*. 2013;132:2492-2501.
- Kim MH, Jeong HJ. Zinc oxide nanoparticles demoted MDM2 expression to suppress TSLP-induced mast cell proliferation. *J Nanosci Nanotechnol*. 2016;16:2492-2498.
- Kim MH, Seo JH, Kim HM, Jeong HJ. Aluminum-doped zinc oxide nanoparticles attenuate the TSLP levels via suppressing caspase-1 in activated mast cells. *J Biomater Appl*. 2016;30:1407-1416.
- Kimura S, Pawankar R, Mori S, Nonaka M, Masuno S, Yagi T, Okubo K. Increased expression and role of thymic stromal lymphopoietin in nasal polyposis. *Allergy Asthma Immunol Res*. 2011;3:186-193.
- Kneilling M, Röcken M. Mast cells: novel clinical perspectives from recent insights. *Exp Dermatol*. 2009;18:488-496.
- Liu T, Li TL, Zhao F, Xie C, Liu AM, Chen X, Song C, Cheng L, Yang PC. Role of thymic stromal lymphopoietin in the pathogenesis of nasal polyposis. *Am J Med Sci*. 2011;341:40-47.
- Mandhane SN, Shah JH, Thennati R. Allergic rhinitis: an update on disease, present treatments and future prospects. *Int Immunopharmacol*. 2011;11:1646-1662.
- Moon PD, Choi IH, Kim HM. Berberine inhibits the production of thymic stromal lymphopoietin by the blockade of caspase-1/NF- κ B pathway in mast cells. *Int Immunopharmacol*. 2011a;11:1954-1959.
- Moon PD, Choi IH, Kim HM. Epigallocatechin-3-O-gallate inhibits the production of thymic stromal lymphopoietin by the blockade of caspase-1/NF- κ B pathway in mast cells. *Amino Acids*. 2012c;42:2513-2519
- Moon PD, Choi IH, Kim HM. Naringenin suppresses the production of thymic stromal lymphopoietin through the blockade of RIP2 and caspase-1 signal cascade in mast cells. *Eur J Pharmacol*. 2011b;671:128-132.
- Moon PD, Jeong HJ, Kim HM. Down-regulation of thymic stromal lymphopoietin by curcumin. *Pharmacol Rep*. 2013;65:525-531.
- Moon PD, Jeong HJ, Kim HM. Effects of schizandrin on the expression of thymic stromal lymphopoietin in human mast cell line HMC-1. *Life Sci*. 2012a;91:384-388.
- Moon PD, Kim HM. The suppression of thymic stromal lymphopoietin expression by selenium. *Amino Acids*. 2012b;43:999-1004
- Moon PD, Kim HM. Thymic stromal lymphopoietin is expressed and produced by caspase-1/NF- κ B pathway in mast cells. *Cytokine*. 2011;54:239-243.
- Mou Z, Xia J, Tan Y, Wang X, Zhang Y, Zhou B, Li H, Han D. Overexpression of thymic stromal lymphopoietin in allergic rhinitis. *Acta Otolaryngol*. 2008;8:1-5.
- Nam SY, Oh HA, Choi Y, Park KY, Kim HM, Jeong HJ. Inhibition of IL-32 signaling by bamboo salt decreases pro-inflammatory responses in cellular models of allergic rhinitis. *J Med Food*. 2014;17:939-948
- Oh HA, Han NR, Kim MJ, Kim HM, Jeong HJ. Evaluation of the effect of kaempferol in a murine allergic rhinitis model. *Eur J Pharmacol*. 2013a;718:48-56
- Oh HA, Ryu JG, Cha WS, Kim HM, Jeong HJ. Therapeutic effects of traditional Korean medicine, Jeechool-Whan in allergic rhinitis model. *TANG*. 2012;2: e9.
- Oh HA, Seo JY, Jeong HJ, Kim HM. Ginsenoside Rg1 inhibits the TSLP production in allergic rhinitis mice. *Immunopharmacol Immunotoxicol*. 2013b;35:678-686.
- Oskeritzian CA, Hait NC, Wedman P, Chumanevich A, Kolawole EM, Price MM, et al. The sphingosine-1-phosphate/sphingosine-1-phosphate receptor 2 axis regulates early airway T-cell infiltration in murine mast cell-dependent acute allergic responses. *J Allergy Clin Immunol*. 2015;135:1008-1018.
- Pandey A, Ozaki K, Baumann H, Levin SD, Puel A, Farr AG, Ziegler SF, Leonard WJ, Lodish HF. Cloning of a receptor subunit required for signaling by thymic stromal lymphopoietin. *Nat Immunol*. 2000;1:59-64.
- Park LS, Martin U, Garka K, Gliniak B, Di Santo JP, Muller W, Largaespada DA, Copeland NG, Jenkins NA, Farr AG, Ziegler SF, Morrissey PJ, Paxton R, Sims JE. Cloning of the murine thymic stromal lymphopoietin (TSLP) receptor: Formation of a functional heteromeric complex requires interleukin 7 receptor. *J Exp Med*. 2000;192:659-670.
- Perlman DS. Pathophysiology of the inflammatory response. *J Allergy Clin Immunol*. 1999;104:S132-S136.
- Plaut M, Pierce JH, Watson CJ, Hanley-Hyde J, Nordan RP, Paul WE. Mast cell lines produce lymphokines in response to cross-linkage of Fc epsilon RI or to calcium ionophores. *Nature*. 1989;339:64-67.
- Sims JE, Williams DE, Morrissey PJ, Garka K, Foxworthe D, Price V, Friend SL, Farr A, Bedell MA, Jenkins NA, Copeland NG, Grabstein K, Paxton RJ. Molecular cloning and biological

characterization of a novel murine lymphoid lymphoid growth factor. *J Exp Med*. 2000;192:671-680.

Soumelis V, Reche PA, Kanzler H, Yuan W, Edward G, Homey B, Gilliet M, Ho S, Antonenko S, Lauerman A, Smith K, Gorman D, Zurawski S, Abrams J, Menon S, McClanahan T, de Waal-Malefyt Rd R, Bazan F, Kastelein RA, Liu YJ. Human epithelial cells trigger dendritic cell mediated allergic inflammation by producing TSLP. *Nat Immunol*. 2002;3:673-680.

Xu WB, Haddad EB, Tsukagoshi H, Adcock I, Barnes PJ, Chung KF. Induction of macrophage inflammatory protein 2 gene expression by interleukin 1 β in rat lung. *Throx*. 1995;50:1136-1140.

Yang G, An D, Lee MH, Lee K, Kim B, Suman CK, Ham I, Choi HY. Effect of *Acer tegmentosum* bark on atopic dermatitis-like skin lesions in NC/Nga mice. *J Ethnopharmacol*. 2016;177:53-60.

Yang G, Choi CH, Lee K, Lee M, Ham I, Choi HY. Effects of *Catalpa ovata* stem bark on atopic dermatitis-like skin lesions in NC/Nga mice. *J Ethnopharmacol*. 2013;145:416-423.

Yoo MS, Jin MH, Lee SY, Lee SH, Kim B, Roh SS, Choi IH, Lee MS, Kim HM, Jeong HJ. Cordycepin Suppresses Thymic Stromal Lymphopoietin Expression via Blocking Caspase-1 and Receptor-Interacting Protein 2 Signaling Pathways in Mast Cells. *Biol Pharm Bull*. 2016a;39:90-96

Yoo MS, Kim HM, Jeong HJ. Acteoside attenuates TSLP-induced mast cell proliferation via down-regulating MDM2. *Int Immunopharmacol*. 2015;26:23-29.

Yoo MS, Park CL, Kim MH, Kim HM, Jeong HJ. Inhibition of MDM2 expression by rosmarinic acid in TSLP-stimulated mast cell. *Eur J Pharmacol*. 2016b;771:191-198

Zhu DD, Zhu XW, Jiang XD, Dong Z. Thymic stromal lymphopoietin expression is increased in nasal epithelial cells of patients with mugwort pollen sensitive-seasonal allergic rhinitis. *Chin Med J (Engl)*. 2009;122:2303-2307.

Ziegler SF, Roan F, Bell BD, Stoklasek TA, Kitajima M, Han H. The biology of thymic stromal lymphopoietin (TSLP). *Adv Pharmacol*. 2013;66:129-155.