

## Original article

# EVALUATION OF OXYTOCIN LIKE EFFECTS OF *Uvariodendron kirkii* (Verdec.) EXTRACTS ON ISOLATED UTERINE STRIPS OF WISTAR RATS

Esther Wairimu Kinyua<sup>1\*</sup>, Charles Irungu Maina<sup>1</sup>, Catherine Kaluwa Kaingu<sup>2</sup>, David Kayaja Wafula<sup>3</sup>

<sup>1</sup>Department of Biological Sciences, Egerton University, Nairobi, Kenya, <sup>2</sup>Department of Veterinary Anatomy and Physiology, University of Nairobi, Nairobi, Kenya. <sup>3</sup>Department of Medical Physiology, University of Nairobi, P.O Box 30197-00100, Nairobi, Kenya.

## ABSTRACT

Uterotonics have the ability to contract uterus. Such plants might be useful in augmenting or inducing labour, expelling retained afterbirth and for abortifacient purposes. Limitations associated with conventional treatments have made herbal medicines a feasible alternative for the management of these conditions. The aim of this study was to evaluate the contractile effects of *Uvariodendron kirkii* extracts on isolated uterine strips of female Wistar rats. Isolated strips of Wistar rats' uteri were treated with 20, 40, 80 and 160 mg/ml concentrations of *Uvariodendron kirkii* aqueous extract. The plant extract was also tested against prostaglandin and oxytocin induced uterine contractions. *Uvariodendron kirkii* extract concentrations (20, 40, 80 and 160 mg/ml) increased the frequency of uterine contraction (16.53, 25.12, 33.48 and 56.39 percentages respectively) compared to the control. The graded extract concentrations caused a significant increase in amplitude (force) of uterine contractions by 2.87, 9.22, 16.37 and 24.32 percentages respectively. The concentrations significantly increased the frequency of oxytocin induced uterine contractions by 6.92; 28.31; 47.06, 58.78 percentages respectively. The graded extract concentrations also significantly increased the amplitude of oxytocin induced uterine contractions by 6.07; 9.40; 15.19 and 23.56 percentages respectively. *Uvariodendron kirkii* extract concentrations significantly increased the frequency and amplitude of prostaglandin induced contractions. The percentage increase in frequency was 11.44, 8.92, 20.65 and 35.71 at 20, 40, 80 and 160 mg/ml respectively. The mean amplitude of prostaglandin induced uterine contractions also increased (4.75, 3.89, 8.29 and 15.91% at 20, 40, 80 and 160 mg/ml respectively). The extract caused a dose dependent increase in uterine frequency and amplitude of contraction. The findings of this study are useful in generating a novel uterotonic agent that will be useful in augmenting labour or in expelling retained after birth in cattle. More studies at molecular level will further elucidate the plant mechanism of action.

**Keywords:** *Uvariodendron kirkii*, Oxytocin induced, Prostaglandin induced, Uterine contractions

## INTRODUCTION

Post-partum haemorrhage is the single leading cause of maternal morbidity and mortality in developing countries (Fukami *et al.*, 2019). Maternal mortality occurs within 24 hours of childbirth, mostly from excessive bleeding (Ngwenya and Bulawayo, 2016). Medicinal plants that can be used to induce labour, treat post-partum haemorrhage and retained afterbirth are of great importance especially in rural parts of developing countries where health facilities are far and have inadequate supplies of emergency medicines (Kaingu *et al.*, 2012). *Uvariodendron kirkii* (CK008) is a native plant in Tana River county, Kenya that is traditionally used to treat menorrhagia, dysmenorrhea, retained placenta and post-partum haemorrhage (Kaingu *et al.*, 2013). It has a fertility regulating effects such as anti-fertility and

anti-implantation properties (Kaingu, 2016). The plant causes uterine contractions thereby explaining the anti-implantation effect. This study therefore, evaluated the contractile effects of aqueous root bark extracts of *Uvariodendron kirkii* on isolated rat uterine tissue in the presence and absence of oxytocin and prostaglandin F<sub>2α</sub>.

## MATERIALS AND METHODS

### Collection and preparation of the plant material

The plant was harvested from Tana River County, Kenya. The plant material was collected specifically in the Itsowe, Garsen and Ngao subdivisions because of widespread use of herbal medicine and inaccessibility to health facilities (Kaingu *et al.*, 2013). *Uvariodendron kirkii* roots (CK008) were collected, dried, ground and the powder stored as per the method described in Kaingu *et al.*, 2017.

### Extract preparation

As per method described by (Kaingu *et al.*, 2017). Extract was then weighed to determine yield.

\*Correspondence: Esther Wairimu Kinyua

E-mail: estherkinyua54@gmail.com

Received Nov 12, 2019; Accepted Dec 23, 2019; Published Feb 28, 2020

doi: <http://dx.doi.org/10.5667/tang.2020.0002>

©2020 by CellMed Orthocellular Medicine Pharmaceutical Association

This is an open access article under the CC BY-NC license.

(<http://creativecommons.org/licenses/by-nc/3.0/>)

# EVALUATION OF OXYTOCIN LIKE EFFECTS OF *Uvariiodendron kirkii* (Verdec.) EXTRACTS ON ISOLATED UTERINE STRIPS OF WISTAR RATS

## Aqueous stock solution

160 mg/ml of *Uvariiodendron kirkii* extract was prepared by weighing 1600 mg of freeze-dried extract and reconstituting with 10 millilitres of distilled water. Similarly, stock solution of 80, 40 and 20 mg / ml dose levels were prepared through double dilutions of the stock solution (Kaingu *et al.*, 2012).

## Experimental Animals

Female Wistar rats aged 10 - 12 weeks old and weighing between 180-220 grams were used. They were kept in cages (5 per cage), in an air-conditioned room at  $22 \pm 2^\circ\text{C}$  and 60-70% relative humidity. Cages contained untreated wood shavings as beddings which were changed every other day. The animals were exposed to 12 hours light and 12 hours dark cycle, fed with commercial rat pellets (Unga Feeds Limited, Kenya) and had free access to clean water *ad libitum* (Bozcheloei *et al.*, 2017). All the rats were handled humanely in accordance with the institution's Faculty of Veterinary Medicine Biosafety, Animal Use and Ethics Committee (FVM BAUEC/2019/190) guidelines and allowed to acclimatize for one week before study onset.

## Experimental design

A completely randomized design was adopted. Female laboratory inbred Wistar rats, from the Department of Biochemistry, University of Nairobi, were used. Five rats were used for positive control (oxytocin). Five rats for each extract concentration (20, 40, 80, and 160 mg/ml), 5 rats for prostaglandin induced concentration and 5 rats for oxytocin induced contractions.

## Preparation of the uterine tissues

Non-pregnant female rats were used and their oestrus cyclicity was closely monitored daily between 9 and 10 a.m. for about 15 days. Vaginal smears were collected, and slides observed under a light microscope in order to ascertain the oestrus stage (Marcondes *et al.*, 2002). Rats in proestrus and oestrus stages of the oestrus cycle were selected as this is when the uterus is more receptive to contractility. One of the mechanisms that successfully deliver sperm to the fallopian tubes. In order to increase uterine sensitivity to contractile agents, the rats were injected with 2 mg/kg diethylstilbestrol subcutaneously 24 hours prior to study onset (Misonge *et al.*, 2014). The rats were then humanely sacrificed using diethyl ether (Goodies *et al.*, 2015). The uterine horns were immediately harvested and placed into a Petri dish containing previously warmed and aerated de Jalon solution. The uterine tissues were excised of all connective tissue. The uterine horns were cut into 2 cm strips (Watcho *et al.*, 2010) and each strip was mounted in an organ bath containing 20 ml de Jalon's solution (Pakoussi *et al.*, 2015; Kaingu *et al.*, 2012).

De Jalon solution was maintained at a temperature of  $37 \pm 0.50^\circ\text{C}$  and continuously bubbled with a mixture of 95% oxygen and 5% carbon dioxide (Misonge *et al.*, 2014). Each uterine strip was mounted vertically within an organ bath. The upper end of the segment was hooked to an isometric force transducer (ML500/A, AD Instrument) coupled with Power Lab data acquisition system (Power Lab 8/30). The frequency of contraction (number of peaks recorded) and amplitude of contraction (in microvolts) were recorded and analysed using Chart5 software for windows.

## Positive control contractions.

Non-pregnant uterine strips were harvested and prepared as described above. They were mounted and allowed (one at a time) to equilibrate for 30 minutes in de Jalon solution alone, after

which its contractions were recorded for 10 minutes. These initial contractions were taken as the negative control recordings. After 10 minutes of negative control contractions the tissue was exposed to 1.0 ml Oxytocin which contained 10 IU synthetic oxytocin (Bafor *et al.*, 2017). The isometric contractions were recorded for 10 minutes (taken as positive control), Frequency (rate) and amplitude (force) of uterine contractions was analysed from the chart recordings. Frequency was taken as the number of contractions recorded over the 10-minute period, that is, the number of peaks recorded. The amplitude or force was taken as the mean height in microvolts ( $\mu\text{V}$ ) of the peaks produced over the 10-minute period. By comparing the frequency and amplitude of contraction for the control and test groups one was able to determine whether the extract increased, reduced or did not cause any effect in terms of frequency and amplitude of uterine contraction.

## Effect of *Uvariiodendron kirkii* aqueous extracts on non-pregnant isolated rats' uteri

Fresh uterine horns were prepared as previously described. Non-pregnant uterine tissue mounting was done. Soon after the first 10 minutes of negative control contractions, the tissue was exposed to 1 ml of 20 mg/ml *Uvariiodendron kirkii* aqueous extract. The extract concentration was added to the organ bath inner chamber (containing de Jalon solution). Isometric contractions were recorded for 10 minutes, after which fresh de Jalon solution was used to wash the organ chamber before mounting a fresh uterine strip, which was exposed to 40 mg/ml extract concentration (Kaingu *et al.*, 2012). The process was repeated using 80 and 160 mg/ml. Each extract concentration challenge was carried out using five rats. Frequency and amplitude of uterine contraction was calculated and recorded.

## The effect of *Uvariiodendron kirkii* aqueous extracts on oxytocin induced uterine contractions

Fresh uterine tissues were prepared. The tissue was mounted, one at a time, and after 10 minutes of negative control contractions was exposed to 1.0 ml oxytocin (containing 10 IU synthetic oxytocin). The isometric contractions were recorded for 10 minutes (taken as positive control), after which the strip was not rinsed and organ bath contents not altered. The uterine strip was thereafter exposed to 1 ml of 20 mg/ml of *Uvariiodendron kirkii* aqueous extract and isometric contractions recorded for 10 minutes. The extract concentration was added into the organ bath inner chamber. The process was repeated 5 more times using fresh strips (Bafor *et al.*, 2017; 2018). Similarly, fresh uterine strips were exposed to 40, 80 and 160 mg/ml extract concentration challenge. Frequency and amplitude of uterine contraction was calculated and recorded as previously described.

## The effect of *Uvariiodendron kirkii* aqueous extracts on prostaglandin induced uterine contractions

The above procedure was repeated using 1.0 ml prostaglandin F2 $\alpha$  which contained 250  $\mu\text{g}$  active ingredient, cloprostenol.

## Data analysis

All values were expressed as percentage increase or decrease in mean  $\pm$  standard error of mean (SEM) relative to the controls, using the following formula.

$$\% \text{ contractions} = \frac{\text{frequency or amplitude after treatment} - \text{control contractions}}{\text{control contractions}} \times 100$$

\* 100

GraphPad prism 8.0.1(244) was used to analyse the data, that is


# EVALUATION OF OXYTOCIN LIKE EFFECTS OF *Uvariadendron kirkii* (Verdec.) EXTRACTS ON ISOLATED UTERINE STRIPS OF WISTAR RATS

frequency and amplitude of uterine contractions using one-way ANOVA. P-values ( $P < 0.05$  \*,  $P < 0.01$  \*\* and  $P < 0.001$  \*\*\*) was considered significant. A post hoc Tukey's multiple comparison test was conducted to analyse statistical differences among groups.


## Positive control

The data is expressed as mean  $\pm$  standard error of mean (SEM) as a percentage increase or decrease in contraction frequency and amplitude relative to the control. Upon addition of 1ml of oxytocin the frequency and amplitude of contraction increased by  $11.29 \pm 0.774$  % and  $18.48 \pm 4.363$  % respectively, compared to the negative control with DE Jalon solution only.


## RESULTS




**Fig.1A** Effect of *Uvariadendron kirkii* extract on isolated uterine frequency of contraction.




**Fig.1B** Effect of *Uvariadendron kirkii* extract on isolated uterine amplitude of contraction.




**Fig.2A** The effect of *Uvariadendron kirkii* extract on frequency of oxytocin induced uterine contractions.



**Fig.2B** Effect of *Uvariadendron kirkii* extract concentrations on amplitude of oxytocin induced uterine contractions.



**Fig.3A** The effect of *Uvariadendron kirkii* extract on frequency of prostaglandin induced uterine contractions.



**Fig.3B** The effect of *Uvariadendron kirkii* extract concentrations on amplitude of prostaglandin induced uterine contractions.

#### Effect of *Uvariadendron kirkii* aqueous extract on frequency and amplitude of uterine contraction

Figure 1A and 1B represent the effect of *Uvariadendron kirkii* extracts on uterine frequency and amplitude of contraction. Upon addition of 20 mg/ml of the extract, the uterine contraction frequency increased by  $16.53 \pm 2.4$  %,  $25.12 \pm 4.88$  % at 40 mg/ml;  $33.48 \pm 7.37$  % at 80 mg/ml and  $56.39 \pm 9.82$  % at 160 mg/ml. The mean uterine contraction frequencies were significant ( $p < 0.05$ ) at 40 mg/ml, ( $p < 0.01$ ) at 80 mg/ml and ( $p < 0.0001$ ) at 160 mg/ml. Upon addition of 20, 40 and 80 mg/ml *Uvariadendron kirkii* extract, the amplitude increased by  $2.87 \pm 0.333$  %,  $9.22 \pm 3.779$  % and  $16.37 \pm 5.802$  % respectively. 20, 40 and 80 mg/ml *Uvariadendron kirkii* extract concentrations caused a non-significant difference in amplitude of uterine contractions at ( $p < 0.05$ ). However, at 160 mg/ml, there was a significant ( $p < 0.01$ ) difference in amplitude of contraction, as the amplitude increased by  $24.32 \pm 5.997$  % compared to the control. There was no significant difference in frequency of uterine contractions at 20, 40 and 80 mg/ml, and oxytocin (positive control) at ( $p < 0.05$ ). However, there was a significant difference in frequency of uterine contractions between 160 mg/ml *Uvariadendron kirkii* extract and oxytocin at  $p < 0.0001$ . There was a non-significant difference in amplitude of uterine contractions in all *Uvariadendron kirkii* extract concentrations and oxytocin at  $p < 0.05$  level of significance.

#### Effect of *Uvariadendron kirkii* aqueous extract on frequency and amplitude of oxytocin- induced uterine contraction

Figure 2A and 2B represent the effect of *Uvariadendron kirkii* extracts on oxytocin induced uterine frequency and amplitude of contraction. Oxytocin frequency and amplitude of uterine contraction was taken as control. The uterine contraction frequency increased by  $6.92 \pm 5.582$  % at 20 mg/ml,  $28.31 \pm 7.193$  % at 40 mg/ml,  $47.06 \pm 9.684$  % at 80 mg/ml and  $58.78 \pm 11.75$  % at 160 mg/ml. The significance difference was at ( $p < 0.01$ ) at 80 mg/ml and ( $p < 0.001$ ) at 160 mg/ml. At 20, 40, 80 and 160 mg/ml *Uvariadendron kirkii* respectively the mean amplitude of uterine contraction increased by  $6.07 \pm 1.617$  %;  $9.40 \pm 2.087$  %;  $15.19 \pm 1.250$  % and  $23.56 \pm 3.096$  % respectively. The increase in amplitude was significant ( $p < 0.05$ ) at 40 mg/ml, ( $p < 0.001$ ) at 80 mg/ml and ( $p < 0.0001$ ) at 160 mg/ml.

#### Effect of *Uvariadendron kirkii* aqueous extract on prostaglandin-induced uterine contraction

Figure 3A and 3B show representatives on frequency and amplitude of prostaglandin induced uterine contractions. Prostaglandin frequency and amplitude of uterine contraction was taken as control. The percentage increase in frequency of uterine contraction was  $11.44 \pm 2.626$  % at 20 mg/ml,  $8.916 \pm 1.257$  % at 40 mg/ml,  $20.65 \pm 1.462$  % at 80 mg/ml and  $35.71 \pm 8.821$  % at 160 mg/ml. There was a significant difference ( $p < 0.05$ ) at 80 mg/ml and ( $p < 0.0001$ ) at 160 mg/ml. The mean amplitude uterine contractions increased by  $4.75 \pm 0.694$  % at 20 mg/ml,  $3.89 \pm 0.639$  % at 40 mg/ml,  $8.29 \pm 0.766$  % at 80 mg/ml and  $15.91 \pm 3.46$  % at 160 mg/ml. There was significant difference ( $p < 0.05$ ) at 80 mg/ml and ( $p < 0.0001$ ) at 160 mg/ml.

## DISCUSSION

#### Effect of *Uvariadendron kirkii* extract on uterine smooth muscle

The most significant finding of this study was the increase in

frequency (rate) and amplitude (force) of uterine contractions caused by graded concentrations of *Uvariadendron kirkii* aqueous extract, as shown in Figures 1A and 1B. The plant has been used in Tana River county to expel the placenta, as a contraceptive and also as an abortifacient. The root bark is boiled in water and the decoction taken orally daily for the fertility regulation and the leaves are inserted in the womb after delivery to expel the placenta.

Other plants have been reported to have similar effects like *Uvariadendron kirkii* on the uterus. For instance, Ahangarpour *et al.*, (2010) reported that aqueous extract of *Cassia italica* leaves increased uterine contractions in rats, with the highest doses of the extract having the most contractile effect on the peak and frequency of uterus contraction. Another study by Kaingu *et al.*, (2012), reported that aqueous extracts of *Euclea divinorum* and *Ricinus communis* enhanced uterine contractility directly. Pakoussi *et al.*, (2018), also reported *Spondias mombin* leaves extract induced a spontaneous increase in uterine contraction amplitude by inducing prostaglandins release. Bafor *et al.*, (2009) also found out that the aqueous extract of the leaves of *Ficus exasperata* increased the frequency of rhythmic uterine contractions in Sprague-Dawley Rats, thereby justifying its use in easing of childbirth.

Nikolajsen *et al.*, (2011), have reported several Tanzanian plants to have induced strong and frequent uterine contractions in Sprague-Dawley Rats. The plants that increased both the force and frequency of uterine contractions included *Bidens pilosa*, *Commelina africana*, *Desmodium barbatum*, *Manihot esculenta*, *Ocimum suave*, *Oldenlandia corymbosa*, *Sphaerogyne latifolia*, *Obetia radula*, *Rubia cordifolia*, and *Triumfetta microphylla*. Ibrahim *et al.*, (2018), reported that the intensity of the contractions in presence of *Ficus deltoidea* var. *Angustifolia* aqueous extract increased moderately following the administration of the extract in a dose-dependent manner which is similar to this study. These results support their traditional claim of use of the plants in assisting labour, improving menstrual circulation, removal of retained placenta and treating post-partum hemorrhage.

The effect of *Uvariadendron kirkii* on uterine contractions can be explored in terms of treating or managing retained placenta as one of the causes is failed uterine contractions by the retroplacental myometrium (Urner *et al.*, 2014). Retained placenta is defined as failure of the foetal placenta (tufts) to separate from the maternal placenta (crypts). It is a serious problem in dairy cattle as is the main cause of cattle infertility and decreased milk yield (Zubair and Ahmad, 2014). Several studies have reported the use of medicinal plants in expelling retained placenta in cattle. The leaves of *Vernonia amygdalica* has been used to treat retained placenta by mixing with table salt, then administered to cow orally, and after some time the animal is relieved from retained placenta (Birhanu and Abera, 2015; Tekle, 2014).

The action of chemical extract from *Vernonia amygdalica* act as increasing uterine contraction by decreasing progesterone concentration, this increment of uterine contraction enhance removal of retained placenta (Yeap *et al.*, 2010). The chemicals such as tannins and flavonoids contained in the extracts of uterotonic plants act as antioxidants, antimicrobial and initiates the contractility of the uterus, thus placenta is easily detached and removed from the uterus (Abdisa, 2018). Tannin for instance produced by the plant has an astringent property that act by shrinking the small blood vessels which in turn lessens the capillary pressure and in turn lead to separation of the foetal membranes from the uterus (Amiera *et al.*, 2014).

Flavonoids on the other hand has antioxidant properties and also assist in oestrogen activation. Oestrogen production leads to activation of contraction proteins, connexins, which in turn aid in placenta removal by increasing uterine contractions (Abdisa, 2018). A study by Kenana *et al.*, (2019) on phytochemical analysis showed *Uvariiodendron kirkii* to contain flavonoids in high concentrations and tannins which may cause the uterine contractility and therefore may aid in the removal of the placenta.

A study on *Ricinus communis* has suggested its use to treat retained foetal membranes (Birhanu and Abera, 2015). Other plants that have been suggested by the traditional medical practitioners for treating retained foetal membrane include: the leaves of *Ensete ventricosum* which are given to cattle during parturition (Mesfin *et al.*, 2016). *Grewia ferruginea* extracts are useful for ease of expulsion of the placenta (Yadav *et al.*, 2014). Bamboo leaves *Bambusa vulgaris* used together with black pepper *Qunda barbaree* assists in placenta expulsion (Abdisa, 2018). Raspberry leaves when consumed during the last 45 days of gestation period have low chances of development of periparturient diseases that is, prolonged labour and retained placenta (Simpson *et al.*, 2001). Whole 100g of *Argemone mexicana* plant when used in feeds together with any available local grass once a day can be used for removal of retained placenta in cows (Yadav *et al.*, 2014). Leaves of both *Opuntia ficus indica* and *Urera hypselodendron* are used for expulsion of retained placenta. The leaves are chopped, then mixed with water and administered to the animal orally (Bobaso *et al.*, 2019). *Achyranthes aspera* L. has been reported to be used by people of Kalahindi district, Orissa, India, where the plant is applied on the genital part and also through inhalation to aid removal of the placenta (Sadangi and Sahu, 2004).

However, (Kaaria *et al.*, 2019) reported that *Asparagus racemosus* caused a relaxation instead of uterine contraction. Similarly, (Pio *et al.*, 2019) showed that *Rhaphiodon echinus* (Re) Crude Ethanol Extract, Lyophilized-Re and Hexane-Re relaxed utero fragment rats, pre-contracted in presence of potassium ions with concentration-dependent response. *Echinophora platyloba* and peppermint oil are reported to reduce uterine contractions. Peppermint oil antispasmodic effect on uterine muscle is by blockage of the calcium channels (Bahmani *et al.*, 2015). Kausar *et al.*, (2016), also reviewed on *Saraca asoka* bark and reported that it is used as a uterine tonic drug where it treats disorders associated with dysmenorrhea, abnormal bleeding, menorrhagia and threatened abortion.

#### Effect of extract on oxytocin and prostaglandin F2 $\alpha$ induced uterine contractions

Physiologically, during labour and the immediate post-partum period, levels of oxytocin and prostaglandins are high within the uterus. During parturition, the levels of Oxytocin and prostaglandin gradually increase to facilitate parturition (Kaingu *et al.*, 2012). The findings of the present study also showed that *Uvariiodendron kirkii* increased the rate and force of Oxytocin induced uterine contractions as shown in Figures 2A and 2B. *Uvariiodendron kirkii* graded contractions also increased the rate and force of prostaglandin induced uterine contractions as shown in Figures 3A and 3B. These findings corroborate those of Katuura *et al.*, (2018), who reported that leaf extracts of *Vernonia amygdalina*, *Maesa lanceolata* and *Rhus natalensis* caused high frequency and amplitude of contractility in isolated rabbit uterus strips similar to, and in most cases higher, than that observed in oxytocin alone. Similarly, Eze *et al.*, (2016) reported that addition of increasing concentrations of oxytocin in the presence of 80 mg ml<sup>-1</sup> aqueous extract of *Sida acuta* produced

significant highest uterine contraction response compared to oxytocin and extract alone which is similar to the response stimulated by *Globimetula braunii* extract as studied by Ie and Zam, (2008). Both the plants potentiated the effect of oxytocin, which is similar to this study.

Oxytocin enhance uterine contractions by two main mode of action in the uterus. One, oxytocin binds to the myometrium oxytocin receptor to produce contractions and two, it binds to endometrium oxytocin receptor to increase prostaglandin production (Amiera *et al.*, 2014). Once oxytocin is stimulated, there is a markedly increase in intracellular concentrations of calcium ions and inositol triphosphate. The release of calcium ions stimulates calcium dependent calmodulin which activates the myosin light chain kinase which in turn catalyzes the contraction response (Arrowsmith and Wray, 2014). PGF2 $\alpha$  on the other hand is produced by the uterine decidual cells. It enhances parturition by induction of luteolysis via the F prostanooids receptors (FP) in the luteal cells. Luteolysis results in cessation of progesterone production and luteal cell apoptosis. A decrease in levels of progesterone leads to the expression of the myometrium genes required for successful parturition (Sugimoto *et al.*, 2015).

This study is also corroborated by Kaingu *et al.*, (2012) where aqueous *Euclea divinorum* and *Ricinus communis* aqueous extracts enhanced prostaglandin and oxytocin induced uterine contractility. *Vitex doniana* bark aqueous extract induced graded uterine contractions and potentiated the effects of prostaglandins, ergometrine and oxytocin (Ladeji *et al.*, 2005). *Trichosanthes kirilowii* (Maxim.) root extract also similarly strengthened spontaneous contractions and enhanced response of uterus muscle to prostaglandin F2 $\alpha$  and that of oxytocin (Gruber and Brien, 2011). Similarly, in this study *Uvariiodendron kirkii* potentiated prostaglandin and oxytocin induced contractions as shown in figures 2A, 2B, 3A and 3B.

However, PGF2 $\alpha$  induced uterine contractions were significantly reduced by ginger oil (*Zingiber officinale*) as shown by (Buddhakala *et al.*, 2008). Other species of plants that have shown smooth muscle relaxation effect with anti-prostaglandin effect under laboratory conditions include *Citrus orantiifolia*, *Rosmarinus officinalis* and *Psidium guajava* (Anel *et al.*, 2014). *Echinophora platyloba* is reported to reduce uterine contractions (Bahmani *et al.*, 2015). *Foeniculum vulgare* inhibited prostaglandins induced uterine contraction (Mirabi *et al.*, 2014).

Uterotonic plants can also be used to prevent or treat postpartum haemorrhage (PPH) (Gruber and Brien, 2011). As earlier stated PPH is caused mainly by uterine atony, and therefore conservative management techniques such as use of uterotonic medications are used as the first line of therapy. This is done to counteract uterine atony and therefore assist in retained placenta expulsion and removal of blood clots. One of the commonly used uterotonic is oxytocin (Likis *et al.*, 2015). This study suggests *Uvariiodendron kirkii* to be a uterotonic agent similar to oxytocin and therefore suggests its use in treatment of PPH.

Attah *et al.*, (2012), suggests use of *Commelina africana*, *Vernonia amygdalina*, *Sida corymbosa*, *Ocimum gratissimum*, *Hyptis suaveolens*, *Duranta repens*, *Calotropis procera*, *Sclerocarya birrea*, and *Saba comorensis* in treatment of PPH. *Commelina africana*, *Vernonia amygdalina* and *Sida corymbosa* however yielded the biggest increase in uterine contractility in a pharmacological uterine model. The pharmacological results are in agreement with our results as *Uvariiodendron kirkii* aqueous extracts increased uterine contractility. It is therefore of vital

# EVALUATION OF OXYTOCIN LIKE EFFECTS OF *Uvariadendron kirkii* (Verdec.) EXTRACTS ON ISOLATED UTERINE STRIPS OF WISTAR RATS

importance to pursue alternative source of remedies for labour management.

## CONCLUSION

In conclusion *Uvariadendron kirkii* root bark possess contractile effects towards isolated rats' uteri. The contractile effect of the plant explains its use in treatment of retained placenta and postpartum haemorrhage by women in Tana River county. The determination of uterotonic activity of crude extract of *Uvariadendron kirkii* provide a starting point towards its potential use in human as a natural source of uterotonic agent.

## RECOMMENDATIONS

Further studies should be carried out to elucidate *Uvariadendron kirkii* mechanism of action in the uterus. It would also be important to isolate phytochemical compounds responsible for its uterotonic effect. Use of collagen gel uterine contractility assays to study the long-lasting uterotonic effect of *Uvariadendron kirkii* extract is also recommended.

## ACKNOWLEDGEMENTS

The authors acknowledge the lab space, materials and equipments' provision by Egerton University, biological sciences department, University of Nairobi, Department of Veterinary Anatomy and Physiology and department of Medical physiology.

## CONFLICT OF INTEREST

The authors declare no conflict of interest

## REFERENCES

Abdisa T. Mechanism of retained placenta and its treatment by plant medicine in ruminant animals in Oromia, Ethiopia. *Journal of Veterinary Medicine and Animal Health*. 2018; 10(6): 135–147.

Ahangarpour A, Oroojan AA and SB. The effects of *Cassia italica* leaves aqueous extract on non-pregnant uterus contraction in rats. *Iranian Journal of Reproductive Medicine*. 2010; 8(4): 179–184.

Amiera ZUR, Nihayah MI, Wahida F and Rajab NF. Phytochemical Characteristic and Uterotonic Effect of Aqueous Extract of *Ficus deltoidea* Leaves in Rats Uterus. *Pakistan Journal of Biological Sciences*. 2014;17(9): 1046–1051.

Andel TV, De Boer HJ, Barnes J and Vandebroek I. Medicinal plants used for menstrual disorders in Latin America, the Caribbean, sub-Saharan Africa, South and Southeast Asia and their uterine properties: A review. *Journal of Ethnopharmacology*. 2014;155(2014), 992–1000.

Arrowsmith S and Wray S. Oxytocin: its mechanism of action and receptor signalling in the myometrium. *Journal of Neuroendocrinology*. 2014;26(6): 1–37.

Attah AF, Brien MO, Koehbach J, Sonibare MA, Moody JO, Smith TJ and Gruber CW. Uterine contractility of plants used to facilitate childbirth in Nigerian ethnomedicine. *Journal of Ethnopharmacology*. 2012;143(1): 377–382

Bafor EE, Omogbai EK, and Ozolua RI. Evaluation of the Uterotonic Activity of the Aqueous Leaf Extract of *Ficus exasperata* Vahl (Moraceae). *Research Journal of Medicinal Plant*. 2009; 3(2), 34–40.

Bafor EE, Nwogu JK, Elvis-Offiah UB, Amaechina F, Ofeimun J, Omoruyi BAOO, Viegelmann C and RuAngeliev E. Modulation of ex-vivo uterine contraction by the methanol leaf extract of *Alchornea laxiflora* Benth. (Euphorbiaceae) and preliminary spectrometric identification of associated secondary metabolites. *Journal of Medicinal Plants for Economic Development*. 2018;2(1): 1–13.

Bafor EE, Omokaro WO, Uwumarongie OH, Elvis-offiah UB, Omoruyi O, Viegelmann CV and Edrada-ebel R. *Dryopteris filix-mas* (Dryopteridaceae) leaves inhibit mouse uterine activity. *Journal of Medicinal Plants for Economic Development*. 2017;1(1): a25.

Bahmani M, Eftekhari Z, Jelodari M, Saki K, Abdollahi R, Majlesi M, Rafieian-Kopaei M and Rasouli S. Effect of Iranian herbal medicines in dysmenorrhea phytotherapy. *Journal of Chemical and Pharmaceutical Research*. 2015;7(2): 519–526.

Birhanu T and Abera D. Survey of ethno-veterinary medicinal plants at selected Horro Guduru Districts, Western Ethiopia. *African Journal of Plant Science*. 2015;9(3): 185–192.

Bobaso H, Abera Z and Asefa K. Survey of Veterinary Herbal Folk Medicine and its Threats in West Arsi Zone, Survey of Veterinary Herbal Folk Medicine and its Threats in. *Global Veterinaria*. 2019;21(3): 127–136.

Bozcheloei MI, Sani MG, Sedaghat K, Jarrahi M, Nejat H and Khorasani MZ. The Effect of Hydro-Alcoholic Extract of *Rosa Damascena* on Uterine Contraction of Virgin rats. *Middle East Journal of Rehabilitation and Health Studies*. 2017;4(3): e13089.

Buddhakala N, Talubmook C, Sriyotha P, Wray S and Kupittayanant S. Inhibitory Effects of Ginger Oil on Spontaneous and PGF 2  $\alpha$  -Induced Contraction of Rat Myometrium. *Planta Med*. 2008;74: 385–391.

Eze OBL, Nwodo OFC, Ogugua VN and Joshua PE. Uterine Contractile Effect of Ethanol Extracts of *Sida acuta* Burm F. Leaves. *Journal of Investigational Biochemistry*. 2016;5(2), 37–40.

Fukami T, Koga H, Goto M, Ando M, Matsuoka S, Tohyama A, Yamamoto H, Nakamura S, Koyanagi T, To Y, Kondo H, Eguchi F and Tsujioka H. Incidence and risk factors for postpartum hemorrhage among transvaginal deliveries at a tertiary perinatal medical facility in Japan. *PLoS ONE*. 2019;14(1), 1–8.

# EVALUATION OF OXYTOCIN LIKE EFFECTS OF *Uvariadendron kirkii* (Verdec.) EXTRACTS ON ISOLATED UTERINE STRIPS OF WISTAR RATS

- Goodies ME, Oghenesuvwe EE and Ejiroghene A. Effects of inhaled anaesthetic agents (Chloroform and Diethyl ether) on fasting blood glucose and haematological parameters in Wistar rats. *Sky Journal of Biochemistry Research*. 2015;4(2): 013–015.
- Gruber CW and Brien MO. Uterotonic Plants and their Bioactive Constituents. *Planta Med*. 2011;77(3): 207–220.
- Ibrahim FW, Derased NI, Romaizat U, Zainudin A, Mohammad N, and Rajab NF. Mechanism Identification of Ficus Deltoidea Aqueous Extract in Rat Uterine Contractions. *Journal Sains Kesihatan Malaysia Isu Khas*. 2018;1: 75–81.
- Ie O and Zam N. Oxytocin properties of the aqueous extract of Globimetula braunii (Loranthaceae). *Parkistan Journal of pharmacological Science*. 2008;21: 356–360
- Kaaria LM, Oduma AJ, Kaingu CK, Mutai PC and Wafula DK. Effect of Asparagus racemosus on selected female reproductive parameters using Wistar rat model. *Discovery Phytomedicine*. 2019;6(4): 199–204.
- Kaingu, C, K. Evaluation of Anti-fertility Potential of Selected Medicinal Plants of Tana River County, Kenya. 2016. University of Nairobi. PHD Thesis
- Kaingu CK, Oduma JA and Kanui T. Preliminary investigation of contractile activity of Ricinus communis and Euclea divinorum extracts on isolated rabbit uterine strips. *Journal of Ethnopharmacology*. 2012;142(2): 496–502.
- Kaingu CK, Oduma JA, Mbaria JM, and Kiama SG. Medicinal plants traditionally used for the management of female reproductive health dysfunction in Tana River County, Kenya. *Tang Humanitas Medicine*. 2013;3(2): 1–10.
- Kaingu CK, Oduma JA, Mbaria JM and Kiama SG. Effects of Croton menyharthii and Uvariadendron kirkii extracts on ovarian corpora lutea and reproductive hormones. *Discovery Phytomedicine*. 2017;4(1): 21–25.
- Katuura, E., Kalabika, E. and Lubega, A. Uterotonic Potential of Selected Plants Used by Ugandan Local Communities in the Treatment of Malaria. *European Journal of Medicinal Plants*. 2018;24(2): 1–12.
- Kauser AS, Hasan A, Parrey SA and Ahmad W. Ethnobotanical, phytochemical and pharmacological properties of Saraca asoca bark: A Review. *European Journal of Pharmaceutical and Medical Research*. 2016;3(6): 274–279.
- Kenana JK, Mbaria JM, Kaingu CK and Okumu PO. Toxicological and Phytochemical Evaluation of Uvariadendron kirkii. *Journal of Ethnobiology and Traditional Medicine*. 2019;130(2019): 1487–1504.
- Ladeji O, Udoh FV and Okoye ZSC. Activity of Aqueous Extract of the Bark of Vitex doniana on Uterine Muscle Response to Drugs. *Phytotherapy Research*; 2005;19: 804–806.
- Likis FE, Sathe NA, Morgans AK, Katherine EH, Young JL, Carlson-Bremer D, Schorn M and Surawicz T. Management of Postpartum Hemorrhage. 2015;151; ES1-ES5
- Marcondes FK, Bianchi FJ and Tanno AP. Determination of the estrous cycle phases of rats: some helpful considerations. *Brazilian Journal of Biology*. 2002;62(4A): 609–614
- Mesfin MM, Fitsum T, Melese Y, Tewodros G and Mebratu A. Documentation of Ethno Veterinary Practices in Selected Sites of Wolaita and Dawuro Zones, Ethiopia. 2016. Available at: <https://globaljournals.org/item/6165-documentation-of-ethno-veterinary-practices-in-selected-sites-of-wolaita-and-dawuro-zones-ethiopia> (Accessed 20th September 2019).
- Mirabi P, Alamolhoda SH, Esmaeilzadeh S and Mojab F. Effect of medicinal herbs on primary dysmenorrhoea - A systematic review. *Iranian Journal of Pharmaceutical Research*. 2014;13(3): 757–768.
- Misonge JO, Ogeto JO, Sengera GO, Mwalukumbi JM, Mwaura AM and Juma SD. Evaluation of phytochemistry and uterotonic activity of root aqueous extract of Uvariadendron anisatum verdec. used in childbirth in Eastern / Central Kenya. *Journal of Pharmacy*. 2014;4(12): 48–53.
- Ngwenya S, and Bulawayo M. Postpartum haemorrhage: incidence, risk factors, and outcomes in a low-resource setting. *International Journal of Women's Health*. 2016;8: 647–650.
- Nikolajsen T, Nielsen F, Rasch V, Sørensen PH, Ismail F, Kristiansen U, and Jäger AK. Uterine contraction induced by Tanzanian plants used to induce abortion. *Journal of Ethnopharmacology*. 2011;137(1), 921–925.
- Pakoussi T, Metowogo K, Ap M, KEG, Ak A. and Gbeassor M. Effect of Spondias Mombin L. (Anacardiaceae) on Rat Uterine Smooth Muscle Contraction. *International Journal of Basic and Applied Science*. 2015;3(3): 18–28.
- Pakoussi T, Mouzou AP, Metowogo K, Aklikokou KA, and Gbeassor M. How do Spondias mombin L.(Anacardiaceae) leaves extract increase uterine smooth muscle contractions to facilitate child birth in parturient women. *African Health Science*. 2018;18(2), 235–243.
- Pio IDSL, Lavor AL, Damasceno CMD, Menezes PMN, Silva FS and Maia GLA. Traditional knowledge and uses of medicinal plants by the inhabitants of the islands of the São Francisco river, Brazil and preliminary analysis of Rhipidodon echinus (Lamiaceae). *Brazilian Journal of Biology*. 2019;79(1): 87–99.
- Sadangi N and Sahu RK. Traditional veterinary herbal practice of Kalahandi district, Orissa, India. *Journal of Natural Remedies*. 2004;4(2): 131–136.
- Simpson M, Parsons M, Greenwood J and Wade K. Raspberry leaf in pregnancy: its safety and efficacy in labor. *Journal of Midwifery & Women's Health*. 2001;46(2), 51–59.
- Sugimoto Y, Inazumi T and Tsuchiya S. Roles of prostaglandin receptors in female reproduction. *Journal of Biochemistry*. 2015;157: (2), 73–80.
- Tekle Y. An ethno-veterinary botanical survey of medicinal plants in Kochore district of Gedeo Zone, Southern Nations Nationalities and Peoples Regional State (SNNPRs), Ethiopia.

EVALUATION OF OXYTOCIN LIKE EFFECTS OF *Uvariadendron kirkii* (Verdec.) EXTRACTS ON ISOLATED  
UTERINE STRIPS OF WISTAR RATS

*Journal of Scientific and Innovative Research*. 2014;3(4): 433–445.

Urner F, Zimmermann R, and Krafft A. Manual Removal of the Placenta after Vaginal Delivery: An Unsolved Problem in Obstetrics. *Journal of Pregnancy*. 2014;2014: 1–6.

Watcho P, Ngadjui E, Alango NP, Benot NT and Kamanyi A. Evaluation of in vitro uterotonic activities of fruit extracts of *ficus asperifolia* in rats. *Evidence-Based Complementary and Alternative Medicine*. 2010;2010, 1–7.

Yadav SS, Bhukal RK, Bhandoria MS, Ganie SA, Gulia SK and Raghav TBS. Ethnoveterinary Medicinal plants of Tosham block of district Bhiwani (Haryana) India. *Journal of Applied Pharmaceutical Science*. 2014;4(06): 40–48.

Yeap SK, Ho WY, Ky H and Alitheen NB. *Vernonia amygdalina*, an ethnoveterinary and ethnomedical used green vegetable with multiple bio- activities. *Journal of Medicinal Plant Research*. 2010;4(25): 2787–2812.

Zubair M and Ahmad M. Review Article An insight into the Recent Advances on the Physiology and Treatment of Retention of Fetal Membranes in Cattle. *Advances in Animal and Veterinary Sciences*. 2014;2(2): 73–77.