

## 세포교정영양요법(OCNT)을 이용한 사마귀 개선 사례

백민호 약사

경기도 파주시 조리읍 고봉로 1039 파주백화점약국

### A Case Study on the Improvement of Warts Using Ortho-Cellular Nutrition Therapy (OCNT)

Pharmacist, Minh Baek

Paju Baekwajeom Pharmacy, 1039, Gobong-ro, Jori-eup, Paju-si, Gyeonggi-do, Republic of Korea

#### ABSTRACT

**Objective:** To report a case of wart improvement through Ortho-Cellular Nutrition Therapy (OCNT).

**Methods:** CNT was administered to a Korean woman in her 50s who complained of aesthetic and lifestyle discomfort due to warts on her neck.

**Results:** After OCNT administration, the patient's wart symptoms significantly improved, and aesthetic enhancement was confirmed.

**Conclusion:** Appropriate administration of OCNT to wart patients can aid in symptom improvement.

**Keywords** Ortho-Cellular Nutrition Therapy (OCNT), warts, human papillomavirus

#### Introduction

Warts are a common condition with a global prevalence of 7-12%. This disease is a proliferative disorder that occurs when human papillomaviruses (HPV), belonging to the *Papillomaviridae* family, infect keratinocytes.<sup>1</sup> This virus has been linked to genital cancers such as cervical cancer, some head and neck cancers, oropharyngeal cancers, and warts, with more than 170 different subtypes identified to date. These subtypes known to cause warts include types 2, 3, 10, and 27.<sup>2</sup>

Warts are classified and named according to their appearance and location as common warts, plantar or palmar warts, flat warts, and genital warts (condyloma). While these are known to be caused by different HPV subtypes, they generally share common clinical features. It includes papillary shapes protruding from the skin, punctate blood vessels referring to dot-like vascular patterns visible on the skin, and bleeding symptoms that occur when the lesion is scratched.<sup>3</sup>

Although warts are not life-threatening, they can cause stress to patients for aesthetic reasons when they occur in exposed areas such as the face or neck.<sup>4</sup> Moreover, as warts are an infectious disease caused by HPV infection, there is a possibility of spreading to other parts of the body or even to other people.<sup>5</sup> Therefore, appropriate treatment should be

administered to prevent the occurrence and spread of warts.

Treatment options for warts include cryotherapy, thermotherapy, laser therapy, or local injections. However, depending on the patient, these treatments may cause pain and itching at the treatment site and may be accompanied by side effects such as inflammation and pigmentation after the procedure. Additionally, some treatments may be difficult to apply depending on the type of wart and specific patient groups, such as pregnant women or infants.<sup>4</sup> Therefore, applying a treatment method tailored to the patient's condition and situation is essential.

In this case study, the patient complained of discomfort in daily life and appearance due to warts on the neck and showed significant improvement after undergoing Ortho-Cellular Nutrition Therapy (OCNT).

#### Case Report

##### 1. Subject

One case of a wart patient was studied.

- 1) Name: Yoon OO (55 years old/F)
- 2) Diagnosis: Warts
- 3) Onset: 2022
- 4) Treatment period: July 2024 - present
- 5) Chief complaint: Redness and lump formation on the neck area
- 6) Past history: Laser treatment for wart removal in the early 40s
- 7) Social history: None
- 8) Family history: Warts in the paternal family
- 9) Present illness and current medications: None

\*Correspondence: Minh Baek

E-mail: homin8111@naver.com

Received Aug 30, 2024; Accepted Aug 30, 2024; Published Aug 30, 2024  
doi: <http://dx.doi.org/10.5667/CellMed.spc.093>

©2024 by CellMed Orthocellular Medicine Pharmaceutical Association  
This is an open access article under the CC BY-NC license.

(<http://creativecommons.org/licenses/by-nc/3.0/>)

† This report has been translated and edited by the CellMed editor-in-chief, Prof. Beom-Jin Lee.

## 2. Method

Initially, a single wart cream containing alpha-hydroxy acid (AHA), EGCG, and tocopherol as main ingredients was prescribed and instructed to be applied frequently only to the wart lesion area. After one week, the patient was instructed to use it twice daily, in the morning and evening. At this point, Cyaplex balm was additionally prescribed and instructed to be applied in conjunction with the wart cream. This prescription was to be applied widely to the wart lesion area and surrounding areas.

## Results

Initially, warts were observed on the back of the neck, and redness was present in the area. Therefore, OCNT was initiated at this point. About two weeks after application, the redness in the area was confirmed to have faded. (Fig. 1.)

Wart symptoms with skin discoloration and multiple small lumps were also observed on the front of the neck, and OCNT was applied for symptom improvement. After OCNT application, the redness in the patient's lesion area gradually and significantly decreased. The lumps were also observed to have disappeared or reduced in size. (Fig. 2.)

## Discussion

The case patient was a woman in her 50s who visited a pharmacy due to sudden lesions in her neck area. Upon observation, the lesions were suspected to be warts, and through questioning, it was confirmed that she had previously undergone laser treatment for warts long ago.

Currently, there are various methods to treat warts. These include treatments that dissolve keratin using salicylic acid or trichloroacetic acid (TCA),<sup>6</sup> ointments containing sinecatechin or imiquimod that kill viruses or induce immunotherapy,<sup>7</sup> and treatments that directly remove the affected area such as cryotherapy, laser therapy, or injection therapy.<sup>8-10</sup> However, keratin-dissolving treatments do not eliminate HPV, the cause of warts, and ointments that induce virus killing or immunotherapy have the disadvantage of not being applicable for long periods.<sup>7</sup>

Moreover, these treatments have limitations when


performed alone. Therefore, in most cases, multiple treatments are combined.<sup>11</sup> However, patients may feel aversion or discomfort towards specific treatments, and treatment effectiveness may vary among individuals, necessitating the appropriate implementation of personalized treatments.

The wart cream prescribed to the patient was selected to contain various ingredients for improving wart symptoms. Alpha-hydroxy acid (AHA), one of the main ingredients in the cream, helps increase skin moisture content by inducing plasticization and exfoliation of keratinocytes and promoting normal differentiation of epidermal cells.<sup>12</sup> It also contains coix seed extract, which can aid skin whitening by inhibiting the activity of MITF, TYR, TRP-1, and TRP-2, which induce melanin production.<sup>13</sup>


The prescribed cream contains a large amount of EGCG, a natural substance extracted from green tea, which shows excellent antioxidant, anti-inflammatory, and antibacterial effects. Due to these characteristics, it is known to be beneficial for various skin conditions such as psoriasis, dermatitis, and atopic dermatitis.<sup>14</sup> It also contains *Artemisia annua* extract, known as Qing Hao in traditional Chinese medicine, which has been shown to have excellent antibacterial and antifungal effects through numerous studies.<sup>15</sup>

A critical aspect of improving wart symptoms is strengthening the immune function of the skin barrier to prevent recurrence due to latent viruses. Tocopherol, contained in the prescribed wart cream, induces the production of Interleukin-2 (IL-2) and helps proliferate T cells, thereby strengthening immune activity.<sup>16</sup> It also contains carotenoids, which help enhance immunity by promoting the proliferation of natural killer and T cells. Therefore, it is thought that these ingredients positively improved the patient's wart symptoms.

*Centella asiatica* extracts contained in Cyaplex balm are known to positively affect cell recovery by suppressing inflammation through active ingredients such as asiaticoside and madecassoside, calming the lesion area and increasing collagen synthesis.<sup>17</sup> Additionally, anthocyanins help cell recovery by reducing oxidative stress caused by reactive oxygen species and increasing cell mobility.<sup>18</sup> Lanolin has been shown through research to strengthen the skin barrier function and help other active ingredients absorb smoothly.<sup>19</sup> It is believed that these ingredients strengthened the damaged skin tissue and created an excellent synergy effect, allowing the



**Fig. 1. Changes in the patient's wart condition on the back of the neck during OCNT administration.** (A) Photo taken on August 5, 2024, and (B) August 20, 2024. After OCNT was administered, the redness in the lesion area can be seen to have faded.



**Fig. 2. Changes in the patient's wart condition on the front of the neck during OCNT administration.** (A) Photo taken on August 10, 2024, (B) August 12, 2024, and (C) August 20, 2024. As OCNT progressed, the redness in the lesion area can be seen to have faded, and the lumps have either reduced in size or disappeared.

prescribed wart cream to work effectively.

This case study was conducted on a single patient and has limitations in being universally applicable to all wart patients. However, it is considered significant that through simple OCNT, the patient's wart symptoms were significantly improved, the appearance of the lesion area was enhanced, and the quality of life could be improved. Therefore, with the patient's consent, this case is being reported.

## References

- Shahid SK. Recent patents in anti-wart treatment. *Pharm Pat Anal.* May 2020;9(2):53-62.
- Accardi R, Gheit T. Cutaneous HPV and skin cancer. *Presse Med.* Dec 2014;43(12 Pt 2):e435-43.
- Zalaudek I, Giacomel J, Cabo H, et al. Entodermoscopy: a new tool for diagnosing skin infections and infestations. *Dermatology.* 2008;216(1):14-23.
- Zhu P, Qi RQ, Yang Y, et al. Clinical guideline for the diagnosis and treatment of cutaneous warts (2022). *J Evid Based Med.* Sep 2022;15(3):284-301.
- Sabry AO, Patel BC. Papilloma. *StatPearls.* StatPearls Publishing. Copyright © 2024, StatPearls Publishing LLC.; 2024.
- Truong K, Joseph J, Manago B, Wain T. Destructive therapies for cutaneous warts: A review of the evidence. *Aust J Gen Pract.* Oct 2022;51(10):799-803.
- Schöfer H, Tatti S, Lynde CW, et al. Sinecatechins and imiquimod as proactive sequential therapy of external genital and perianal warts in adults. *Int J STD AIDS.* Dec 2017;28(14):1433-1443.
- Andrews MD. Cryosurgery for common skin conditions. *Am Fam Physician.* May 15 2004;69(10):2365-72.
- Maranda EL, Lim VM, Nguyen AH, Nouri K. Laser and light therapy for facial warts: a systematic review. *J Eur Acad Dermatol Venereol.* Oct 2016;30(10):1700-1707.
- Muse ME, Stiff KM, Glines KR, Cline A, Feldman SR. A review of intralesional wart therapy. *Dermatol Online J.* Mar 15 2020;26(3)
- Huang K, Li M, Xiao Y, et al. The application of medical scale in treating plantar warts: analysis and prospect. *J Dermatolog Treat.* Mar 2022;33(2):637-642.
- Babilas P, Knie U, Abels C. Cosmetic and dermatologic use of alpha hydroxy acids. *J Dtsch Dermatol Ges.* Jul 2012;10(7):488-91.
- Goelzer Neto CF, do Nascimento P, da Silveira VC, de Mattos ABN, Bertol CD. Natural sources of melanogenic inhibitors: A systematic review. *Int J Cosmet Sci.* Apr 2022;44(2):143-153.
- Messire G, Serreau R, Berteina-Raboin S. Antioxidant Effects of Catechins (EGCG), Andrographolide, and Curcuminoids Compounds for Skin Protection, Cosmetics, and Dermatological Uses: An Update. *Antioxidants (Basel).* Jun 21 2023;12(7)
- Ekiert H, Klimek-Szczykutowicz M, Rzepliela A, Klin P, Szopa A. Artemisia Species with High Biological Values as a Potential Source of Medicinal and Cosmetic Raw Materials. *Molecules.* Sep 29 2022;27(19)
- Reiter E, Jiang Q, Christen S. Anti-inflammatory properties of alpha- and gamma-tocopherol. *Mol Aspects Med.* Oct-Dec 2007;28(5-6):668-91.
- Bylka W, Znajdek-Awiżeń P, Studzińska-Sroka E, Dańczak-Pazdrowska A, Brzezińska M. Centella asiatica in dermatology: an overview. *Phytother Res.* Aug 2014;28(8):1117-24.
- Van de Velde F, Esposito D, Grace MH, Pirovani ME, Lila MA. Anti-inflammatory and wound healing properties of polyphenolic extracts from strawberry and blackberry fruits. *Food Res Int.* Jul

2019;121:453-462.

19. Carrer V, Guzmán B, Martí M, Alonso C, Coderch L. Lanolin-Based Synthetic Membranes as Percutaneous Absorption Models for Transdermal Drug Delivery. *Pharmaceutics*. Jun 21 2018;10(3)