

The Efficacy of Psychological Interventions for Complex Trauma: A Systematic Review and Meta-Analysis

Hyunjung Choi^{1†} Wooyeol Lee¹ Simyang Heo² Jahye Kim²

¹Department of Psychology, Chungbuk National University, Cheongju; ²Traumahealing Center Human Hearts, Seoul, Korea

This study meta-analyzed interventions for symptom constructs of complex posttraumatic stress disorder related to complex trauma, such as child abuse, multiple interpersonal trauma, and organized violence. From 42 randomized controlled studies, 164 effect sizes were calculated comprising various treatment comparisons and outcomes. Cognitive processing therapy/cognitive therapy (CPT/CT), narrative exposure therapy (NET), phase-based trauma-focused cognitive behavior therapy (phased TF-CBT), single-phased TF-CBT (single TF-CBT), prolonged exposure (PE), eye movement desensitization and reprocessing (EMDR), and present-centered therapy (PCT) each showed moderate to large effect sizes on PTSD. For depression, CPT/CT, phased TF-CBT, single TF-CBT, NET, and PE showed moderate to large effect sizes. Meta regression revealed that efficacy on PTSD and depression was greater with trauma-focused than with present-centered interventions. Limited evidence made it impossible to compare phased treatments and single-phase treatments. Outcomes of disturbance in self organization (DSO) were relatively scarce; however, among those, CPT/CT, PE, and PCT showed large to moderate effect sizes on negative self-concept. For emotion regulation and interpersonal problems, phased TF-CBT seemed to be promising. Treatment completion rates were similar between trauma-focused and present-centered. Population and study characteristics did not affect efficacy; however, we discussed related clinical issues.

Keywords: complex trauma; evidence-based interventions; posttraumatic stress disorder; complex posttraumatic stress disorder; meta-analysis

Recently, the International Classification of Diseases 11th Revision (ICD-11) included a diagnosis of complex posttraumatic stress disorder (CPTSD), which consists of three core posttraumatic stress symptoms (reexperience, avoidance, and sense of threat) and additional symptoms of disturbance in self organization (DSO) represented by affect dysregulation, negative self-concept, and disturbances in relationships (World Health Organization, 2018). This inclusion reflects previous understandings about the complicated impacts of complex trauma (Briere, 1992; Her-

man, 1992; van der Kolk, 1996), and has foundations from recent empirical studies that revealed profiles of CPTSD or DSO that are independent from PTSD (Cloitre, Garvert, Brewin, Bryant, & Maercker, 2013; Knefel, Garvert, Cloitre, & Lueger-Schuster, 2015), or borderline personality disorder (BPD) (Cloitre, Garvert, Weiss, Carlson, & Bryant, 2014). A measurement for CPTSD has also been developed and validated emphasizing clinical utility as well as universal applicability (Cloitre et al., 2018).

Since the official recognition of CPTSD has shown recent progress, searching for empirically supported psychological treatment of CPTSD is required. For PTSD, clinical guidelines of psychological treatments are clearly established (American Psychological Association, 2017); however, some controversial issues remain in the treatment of CPTSD. There are clinical guidelines for CPTSD; for example, the International Society of Traumatic Stress Studies

[†]Correspondence to Hyunjung Choi, Department of Psychology, Chungbuk National University, 1 Chung-daero, Suwon-gu, Cheongju, Korea; E-mail: hchoi@chungbuk.ac.kr

Received Jan 03, 2020; Revised Feb 09, 2020; Accepted Feb 27, 2020

This work was supported by the Ministry of Education of the Republic of Korea and the National Research Foundation of Korea (NRF-2017S1A5B5A07062178).

(ISTSS) expert consensus, suggests that phase-oriented treatment may be beneficial to individuals with complex symptoms (Cloitre et al., 2012). Critiques pointed out that trauma-focused CBT (TF-CBT) developed for PTSD also may be efficacious for people with complex trauma, and that phase orientation lacks evidence in utility (e.g., De Jongh et al., 2016). However, this discussion may be refuted, that it focused on PTSD and depressive symptoms and exclude broad symptoms of DSO and did not consider the possible adverse effects, such as increased distress during the exposure process. A systematic review among women survivors of childhood abuse reported that exposure-based CBTs had shown above moderate effect size for PTSD; however, with CPTSD, affect management interventions brought better improvement and lower drop out (Dorrepal et al., 2014). They concluded that phase-based treatment would be more suitable for CPTSD which prepares clients with distress tolerance for subsequent trauma-focused and exposure-based treatments.

This study aimed to meta-analyze the efficacy of interventions for CPTSD constructs and to identify optimal treatment for complex trauma, including PTSD and related pervasive symptoms. Unlike review studies that rely on a subjective synthesis, a meta-analysis calculates the effect size (Jang & Shin, 2011) and by comparing randomized controlled trials, meta-analysis can give a comprehensive understanding in which treatment for whom may be efficacious (Seidler & Wagner, 2006), providing critical evidence to clinical guidelines and policy making (Cooper, 2010).

Karatzias et al. (2019), which is the first meta-analysis according to the ICD-11 CPTSD, recently reported the efficacy of 51 randomized controlled studies that included outcomes of PTSD and at least one DSO symptom. Studies using compact measurement tools for CPTSD should be first considered, as Brewin et al. (2017) mentioned that not the type of trauma but the symptomatology post trauma would decide the diagnosis of CPTSD. However, because to date, efficacy studies using a dedicated measure for CPTSD were rare (Karatzias et al., 2019), focusing on the type of the trauma for a meta-analysis may provide additional guidelines in real-world settings. To expand the findings, our study focused on populations that had experienced what represents previous definition of complex trauma.

Earlier, Herman (1992, 1997) had identified complex trauma re-

sulting from multiple, repeated, prolonged, and systematic traumatization, especially of an interpersonal nature, such as child abuse, domestic violence, sexual exploitation, combat, or torture may complicate posttraumatic adaptation. Also, Hobfoll and colleagues discovered that it is the degree of resource loss during sustained trauma exposure that leads to severe and chronic symptom trajectories (Hobfoll, Mancini, Hall, Canetti, & Bonanno, 2011). Similarly, as reported by Courtois and Ford (2013), complex trauma involves “traumatic attachment (p. 25)”, which is repetitive and cumulative, causing severe alterations in one’s ordinary developmental process, life, and identity. These conditions were considered to be associated with symptom complications, such as emotion dysregulation, identity problems, disruptions in relationships, somatization, alterations in consciousness, and altered systems of meaning (Briere, 1992; Herman, 1992; Pelcovitz et al., 1997), in addition to core symptoms of PTSD.

Therefore, we focused on studies including the above previous descriptions of complex trauma, and those that dealt with symptoms, such as PTSD, depression, anxiety, dissociation, negative self-concept, emotion dysregulation, interpersonal problems, borderline personality disorder, and self-harm behaviors. Systematic review and meta-analyses included randomized controlled studies of major psychological treatments, including phase-based treatments, trauma-focused treatments, and other psychosocial approaches. For trauma-focused treatments, effect size of each treatment was calculated, because in the applied setting each treatment is trained and conducted distinctively, and fidelity of each treatment is an important issue. Predictors of treatment efficacy were examined using meta-regression analyses. We hypothesized that trauma-focused treatments would be efficacious for PTSD compared to non-trauma focused treatments. For other complex symptoms, phased-based trauma-focused treatments would show higher efficacy compared to a single-phase trauma-focused treatment.

Methods

Study selection

The search period covered from 1990 to September 2017, since it was in the 1990s that discussion of complex trauma first began (e.g., Terr, 1990), up to the date when the search was initiated. The

PsychINFO, PubMed, PILOTS, and Medline databases were searched using the EndNote \times 5. Selections were organized by the PICOT guideline (Institute of Medicine, 2011). For population (P), intervention (I), and outcomes (O), search terms were entered as follows; (complex trauma OR multiple trauma OR child abuse OR child neglect OR child adversity OR maltreatment OR torture OR sexual trauma OR domestic violence OR intimate partner violence OR prostitution OR bereavement) AND (psychotherapy OR psychological treatment OR exposure OR EMDR OR trauma-focused cognitive behavioral therapy OR cognitive processing OR cognitive restructuring OR dialectical behavior therapy OR present-centered OR stress management OR gestalt OR psychoanalytic OR mentalization OR somatic experiencing OR sensorimotor psychotherapy OR advocacy OR group therapy OR psychoeducation) AND (PTSD OR suicid* OR self-harm OR self-injury OR emotion regulation OR self-regulation OR self-organization OR substance OR eating OR borderline personality disorder OR depress* OR identity OR interperson* OR relation*).

For comparison (C), treatment as usual, waiting list, and active treatment comparison studies were selected; and for study type (T), conducting a randomized controlled trial, using validated instruments, and peer reviewed studies were eligible. Exclusion criteria included non-peer reviewed literature, review article, participants under the age 18, offender study, non-face-to face therapy, and non-individual therapy (couple or family therapy, or group therapy as major interventions).

Intervention listed recommended treatments for PTSD (APA, 2017), which are trauma-focused treatments such as prolonged exposure (PE; Foa & Rothbaum, 1998), cognitive processing (Resick & Schnicke, 1992) or cognitive therapy (Ehlers, Clark, Hackmann, McManus, & Fennell, 2005) for PTSD, cognitive behavior therapy (CBT) composed of interventions based on cognitive behavioral principles, eye movement desensitization and reprocessing (EMDR; Shapiro, 1995), and dialectical behavior therapy (Linehan, 1993). Relatively newly recognized somatic based interventions such as somatic experiencing (SE; Levine, 2010) and sensorimotor psychotherapy (SP; Ogden & Fisher, 2015) were also included. Additionally, non-trauma focused, present-centered therapies (PCT) were included, as were treatments from traditionally established approaches in psychotherapy related to our main tar-

get symptoms. Last, "advocacy" was included to seek psychosocial interventions for complicated areas of human rights violations. Comparisons for treatment completers and intent-to-treat analysis were planned. Two researchers independently worked on the search and reviewed titles and abstracts consequently. A third researcher cross-checked and discussed disagreements to find resolutions. Finally, all three researchers reviewed full texts for final selection and resolved discrepancies.

Data extraction

Two researchers each with expertise in trauma studies and statistics, extracted data for standardized differences between treatment groups, seeking agreements. Intent-to-treat scores and follow-up data (versus post data) were firstly coded. When these were not available, completer scores or post data were coded and recorded separately as study characteristics. When both self-report measurements and clinical interviews were reported, clinical interview data were used. Means and SDs were mainly extracted, and SDs were calculated if standard errors were given instead. Because no studies reported pre-post data correlations, a moderate value of $r = .5$ was coded according to Becker (1990). Studies reporting raw differences with confidence intervals, and ds were coded as such. When the number of participants from pre-intervention to follow-up differed in the reported intent-to-treat analysis, we decided using the number of pre-intervention participants, and this calculation were confirmed by being most closely matched with an alternative calculation according to Morris (2008).

Study characteristics were independently coded by three researchers, and the first author cross-checked each data. Each study characteristic was organized into a table, including intervention arms, intervention type (trauma-focused or not, and phased-oriented or not), average number of sessions, number of participants, number of treatment completers (according to the definition of completion in each study), completer rate, follow-up duration, targeted outcomes, measurements of outcomes, analysis type (intent-to-treat or completer), and study location. Each participant's characteristics was also organized, such as female gender (%), low SES (percentage of people with lowest income or insecure social status level, based on the report of each study), and European/Caucasian ethnicity (%). The type of trauma experience (organized violence

such as torture, terrorism, living in a conflict region, detention, or persecution; child abuse; multiple interpersonal trauma; and military trauma), and intervention setting (outpatient, residential, or shelter) were also coded.

Analysis

Quality assessment

Following the methodology checklist for randomized controlled trials of NICE (2012), two researchers completed the quality assessment, and final decisions were made by discussing uncertainties. Low, unclear/unknown, and high risks of bias were assessed in four domains, such as selection, performance, attrition, and detection. Overall risk of bias was decided as low, moderate, and high depending on the level of risk in each domain. At least three low risks of bias on each domain were considered overall as low risk. Two low risks without any high risks were considered to be an overall moderate risk. Overall risk was rated as high when one or more high risks were detected in any domain.

Effect size and meta regression analysis

Comprehensive Meta-Analysis 3 was used for the meta-analyses. As by “shifting the unit of analysis” (Cooper, 1998), effect size was used as the unit of analysis. Hedges’ g was chosen as the effect size to overcome the shortcomings of Cohen’s d which often overestimates the effect size. The effect size and 95% confidence interval were calculated for each intervention. Effect size was interpreted as large ($>.80$), moderate (.50 to .80), small (.20 to .50), or none ($<.20$) according to Cohen (1988). Heterogeneity was assessed using the I^2 and τ^2 statistics. Because we assumed heterogeneity, a random effect model was chosen, with maximum likelihood estimation. We assessed publication bias using the funnel plot, and Trim and Fill (Duval & Tweedie, 2000) and conducted sensitivity analyses. Among comparisons of psychological treatment and TAU/WL, univariate meta-regression was conducted by entering each study characteristic such as type of trauma (childhood abuse, organized violence, multiple interpersonal trauma), compositions of low SES (%), female gender (%), and European/Caucasian ethnicity (%), session length, intent-to-treat (vs. completer) study, risk of bias levels (high vs. low, and high vs. moderate), and trauma-focused treatment (vs. non-trauma focused).

Results

Study selection

A flowchart of data selections is presented in Figure 1. A total of 5,773 articles were searched, then by screening via titles and abstracts, and adding 11 primary articles searched by ancestry approach from secondary analysis articles, a total of 430 articles remained for full text screening. Studies with bereaved participants were all excluded, because the researchers agreed that information to conclude whether it would be a multiple or complex experience were insufficient. The final screening excluded 372 articles, leaving 42 randomized controlled studies eligible for the full inclusion criteria.

Study characteristics

Study and participant characteristics of the 42 randomized controlled studies are presented in Table 1 and in Appendices I and II. Most of the studies were conducted in Western countries. For target symptoms, majority assessed efficacy on PTSD and depression. Studies assessing DSO related symptoms were relatively scarce. Studies were mainly conducted in outpatient settings. Seven studies (16.7%) did not report follow-up data, and the other 35 studies (83.3%) reported follow-up data in which duration ranged from one to 24 months. There were four studies (9.5%) that had a phased-based design. Twenty-four studies (57.1%) reported data of intent-to-treat analysis clearly enough to be included in the meta-analysis.

The total number of participants from all studies was 4,455, ranging from 18 to 347 individuals in each study. Based on type of trauma, child abuse, multiple interpersonal trauma, organized violence, and military-trauma studies were identified. For gender, most studies had higher composition of females, except eight studies (19.0%) with less than 50% of female participants were all related to organized violence. There was no male-only study.

For the ethnicity of the participants, 40.5% included European/Caucasian people at over 50%, and the rest included diverse ethnicities. For SES, twelve studies (28.6%) had less than 50% of people with low SES. There were 12 studies (28.6%) focusing mainly on a low-resource population, including refugees and asylum seekers, a culturally diverse population with low income, and people




Figure 1. Flow chart of search result.

ple in shelter setting.

The average number of individual sessions ranged from one (psychoeducation) to 38.6 (DBT-PE) sessions, and average for all studies was 11.20 sessions. The treatment completer rate ranged

from 48.2 to 100%; one study did not clearly report the number of completers. The average completer rate was calculated (all completers / all participants) to be 75.9%.

Table 1. *Number of Studies According to Study and Participant Characteristics*

Characteristics	n (%)	Characteristics	n (%)
Study characteristics		Participant characteristics	
Location		Type of trauma	
USA	22 (52.4)	Child abuse	13 (31)
Germany	9 (21.4)	Multiple interpersonal trauma	14 (33.3)
Australia	2 (4.8)	Organized violence	11 (26.2)
Canada	2 (4.8)	Military trauma	4 (9.5)
Chile	1 (2.4)	Gender	
China	1 (2.4)	Female only	24 (57.1)
Iraq	1 (2.4)	Female over 50%	8 (19)
Ireland	1 (2.4)	Female less than 50%	8 (19)
Kurdistan	1 (2.4)	Male only	0 (0)
Romania	1 (2.4)	Not clear	2 (4.8)
Thailand	1 (2.4)	Ethnicity	
Setting		European/Caucasian over 50%	17 (40.5)
Outpatient	39 (92.9)	Only non-European/Caucasian	10 (23.8)
Residential	2 (4.8)	Diverse (European/Caucasian less than 50%)	12 (28.6)
Shelter	1 (2.4)	Not clear	3 (7.1)
Target symptom		Social economic status (SES)	
PTSD	37 (88.1)	Low SES people over 50%	19 (45.2)
Depression	42 (100)	Low SES people less than 50%	12 (28.6)
Anxiety	15 (35.7)	Not clear	11 (26.2)
Dissociation	6 (14.3)		
Negative self-concept	7 (16.7)		
Emotion dysregulation	2 (4.8)		
Interpersonal problem	4 (9.5)		
Borderline personality disorder	2 (4.8)		
Substance use	1 (2.4)		
Non-suicidal self-injury and suicidality	1 (2.4)		
Analysis			
Reported follow-up data (1-24 Months)	35 (83.3)		
Conducted intent-to-treat analysis	24 (57.1)		

Quality assessments

Quality assessments of the 42 randomized controlled studies are shown in Appendix II. Assessment of selection bias showed that 20 studies had a low risk of bias in randomized selection with adequate concealment, whereas the other 22 studies were rated as unclear because of limited reporting. Assessing performance bias, 26 studies had low risk, 14 were unclear, and two had a high risk which was related to the uneven number of sessions between comparisons. Whether participants were blind to allocation was mostly not clearly reported or eventually not applicable, and this was also true for the blindness of the therapists. Results of attrition bias assessment showed that 34 had low risk, one was unclear, and seven had high risk. In attrition bias, high risk was mainly related to

attrition rates between comparison groups. Assessing risk for detection, 17 studies had low risk; unclear studies had limited reports on investigator's blindness or used self-report instruments instead of standardized interviews for outcome assessment. Overall, 17 studies (40.4%) had low risk, eight (19.0%) moderate risk, and 17 (40.4%) high risks.

Effect size outcomes

We calculated 164 effect sizes of various comparisons and outcomes from the 42 studies. Further analyses were based on each treatment. The Hedges' g with its confidence interval and the I^2 and τ^2 statistics for each comparison are provided in a supplementary Tables 1–4.

Single-phased trauma-focused psychotherapy

Effect size of single phased trauma treatments such as CPT/CT, NET, PE, single TF-CBT, and EMDR studies were as follows.

Cognitive processing therapy and cognitive therapy (CPT/CT)

There were ten studies (ten comparisons) examining the efficacy of CPT/CT. Outcomes of PTSD, anxiety, depression, dissociation, and negative self-concept symptoms were assessed. In nine comparisons with TAU/WL, CPT/CT showed large effect size on PTSD ($n = 9$, $g = 1.155$, 95% CI .684 to 1.625), depression ($n = 9$, $g = 1.172$, 95% CI .657 to 1.688), and negative self-concept ($n = 3$, $g = 1.110$, 95% CI .027 to 2.193). Small effect size was shown on anxiety ($n = 2$, $g = .390$, 95% CI .140 to .641). Only one study assessed dissociation, which had shown large effect size ($g = 1.352$). Treatment completion rate for CPT/CT ranged from 48.2 to 94.7%, and the average rate was 69.8% (384/550).

Narrative exposure therapy (NET)

Seven studies (eight comparisons) examined NET. Outcomes of PTSD, depression, dissociation, and BPD symptoms were assessed. Among eight comparisons with TAU/WL, NET had large effect size on PTSD ($n = 8$, $g = .969$, 95% CI .464 to 1.474) and moderate effect size on depression ($n = 8$, $g = .580$, 95% CI .320 to .841). One study comparing NET with treatment by experts assessed dissociation ($g = .181$) and BPD symptoms ($g = .221$) showing small effect sizes. Treatment completion rate ranged from 64.7 to 100%, and the average rate was 82.4% (131/159).

Prolonged exposure (PE)

Six studies (eight comparisons) examined PE (including PE with cognitive restructuring). Outcomes of PTSD, depression, anxiety, negative self-concept, and dissociative symptoms were assessed. PE compared with TAU/WL had moderate effect size on PTSD ($n = 7$, $g = .643$, 95% CI .371 to .915), depression ($n = 7$, $g = .514$, 95% CI .319 to .708), and negative self-concept ($n = 2$, $g = .643$, 95% CI .081 to 1.205). Small effect size was shown on anxiety ($n = 4$, $g = .245$, 95% CI .055 to .436). Treatment completion rate ranged from 53.2 to 65.8%, and the average rate was 61.5% (276/449).

Trauma-focused CBT (single TF-CBT)

We grouped five studies using a combination of trauma-focused CBT techniques, such as exposure and skills training, as single TF-CBT. Outcomes of PTSD, depression, anxiety, and substance use symptoms were assessed. In comparisons with TAU/WL, TF-CBT showed large effect size on PTSD ($n = 4$, $g = 1.268$, 95% CI .592 to 1.944), depression ($n = 3$, $g = 1.035$, 95% CI .194 to 1.876), and anxiety ($n = 2$, $g = 1.156$, 95% CI .331 to 1.980). One study reported the efficacy of exposure included CBT on substance use compared to the usual CBT, showing no superiority ($g = .154$). One study did not report the number of completers; other than this, treatment completion rate ranged from 67.1 to 98%, and the average rate was 82.1% (312/380).

Eye movement desensitization and reprocessing (EMDR)

Three studies (four comparisons) examined EMDR. Outcomes of PTSD, depression, anxiety, and negative self-concept symptoms were assessed. Among four comparisons with TAU/WL, EMDR showed large effect size on anxiety ($n = 4$, $g = .830$, 95% CI .351 to 1.309), moderate effect size on PTSD ($n = 4$, $g = .767$, 95% CI .361 to 1.173), and small effect size on depression ($n = 4$, $g = .374$, 95% CI .057 to .691). One study reported a result in negative self-concept that showed a small effect size ($g = .457$). Treatment completion rate ranged from 50 to 100%, and the average rate was 85.9% (55/64).

Phase-based trauma-focused cognitive behavior therapy (phased TF-CBT)

Four studies (four comparisons) examined phase-oriented treatment consisting of DBT based skills training and exposure therapy, including DBT-PTSD, STAIRS-PE (skills training in affect and interpersonal regulation-PE), and DBT-PE arms. Outcomes of PTSD, depression, anxiety, dissociation, BPD, emotion regulation, interpersonal problems, non-suicidal self-injury (NSSI), and suicidality symptoms were assessed. Among comparisons with TAU/WL, large effect size were shown on PTSD ($n = 2$, $g = 1.378$, 95% CI .984 to 1.772) and depression ($n = 2$, $g = .956$, 95% CI .195 to 1.717), and moderate effect size was found in dissociation ($n = 2$, $g = .718$, 95% CI .195 to 1.241). One study reported BPD symptoms, showing small effect size ($g = .440$). Another study reported large effect

size each on emotion regulation ($g = 1.305$) and interpersonal problems ($g = 1.270$). Treatment completion rate ranged from 58.8 to 84.8%, and the average rate was 76.1% (89/117).

Non-trauma focused psychotherapy

Studies of CBT without treating trauma materials, and PCTs were examined.

Non-trauma focused CBT (non-TF-CBT)

We included two studies of stabilization-oriented skills training and behavior activation in non-trauma focused CBTs (non-TF-CBT). These two comparisons with TAU/WL showed small effect size on PTSD symptoms ($n = 2$, $g = .216$, 95% CI $-.038$ to $.470$) and depression ($n = 2$, $g = .441$, 95% CI $.120$ to $.762$). A study of behavior activation showed moderate effect size on anxiety ($g = .513$). Treatment completion ranged from 54.3 to 71.9%, and the average rate was 67.8% (101/149).

Present-centered therapy (PCT)

There were seven PCT (seven comparisons) including three interpersonal psychotherapy (IPT) studies, one holographic reprocessing, one emotion-focused therapy, one present-centered therapy, and one brief psychodynamic psychotherapy. Outcomes of PTSD, depression, anxiety, dissociation, negative self-concept, and interpersonal problem symptoms were assessed. Compared with TAU/WL, present-centered therapies showed moderate effect size on PTSD ($n = 6$, $g = .562$, 95% CI $.203$ to $.921$), anxiety ($n = 3$, $g = .576$, 95% CI $.175$ to $.978$), and negative self-concept ($n = 3$, $g = .588$, 95% CI $.154$ to 1.021). They had small effect size on depression ($n = 6$, $g = .446$, 95% CI $.225$ to $.667$) and interpersonal problems ($n = 2$, $g = .332$, 95% CI $-.108$ to $.771$). One study reported dissociation showing large effect size ($g = 1.081$). Treatment completion ranged from 75 to 94.1%, and the average rate was 79.8% (198/248).

Other psychosocial interventions

One study reported the efficacy of one-session individual psychoeducation vs. WL. Small effect size was shown in PTSD symptoms ($g = .264$); however, none in depression ($g = .065$) or anxiety ($g = .070$). Completion rate was 97.4%. Two studies reported efficacy of advocacy vs. usual care on depression with interpersonal vio-

lence. Results showed no effects ($n = 2$, $g = .048$, 95% CI $-.133$ to $.229$). Finally, one study reported the efficacy of additional mentoring with usual care on depression compared to only usual care. It showed a moderate effect size on depression ($g = .546$). Completion rate was 79.6%.

Active treatment comparisons

There were six studies comparing major trauma-focused interventions. Comparing PE to PE with cognitive restructuring, there was no differences in efficacy on PTSD symptoms ($n = 2$, $g = .117$, 95% CI $-.117$ to $.352$) and depression ($n = 2$, $g = .166$, 95% CI $-.070$ to $.403$).

Two studies compared phased TF-CBT to only skills training, and phased TF-CBT had small effect size on PTSD symptoms ($n = 2$, $g = .403$, 95% CI $-.088$ to $.895$), depression ($n = 2$, $g = .391$, 95% CI $-.099$ to $.881$), and anxiety ($n = 2$, $g = .473$, 95% CI $-.021$ to $.967$).

One study reported efficacy for phased TF-CBT (DBT-PE) on dissociation ($g = .381$), and suicidality ($g = .286$), showing small effect size compared to DBT-only intervention. Phased TF-CBT (DBT-PE) compared to DBT-only showed no better efficacy on NSSI ($g = .190$). Another comparison reported that phased TF-CBT (STAIR/exposure) showed moderate effect size on emotion regulation ($g = .576$) and interpersonal problems ($g = .626$) compared to skills training only. Also, efficacy of phased TF-CBT (STAIR/exposure) compared to exposure with supportive counseling, showing large effect size on anxiety ($g = .922$), moderate effect size on interpersonal problems ($g = .586$), and small effect size on symptoms of PTSD ($g = .350$), depression ($g = .231$), and emotion regulation ($g = .390$).

Publication bias and sensitivity analysis

Presented in the Figure 2, publication bias was examined by the funnel plot of psychological treatments vs. TAU/WL, among outcomes of PTSD, depression, and anxiety symptoms, which had more than 10 comparisons required to draw the funnel plot. Funnel plot and Trim and Fill (Duval & Tweedie, 2000) revealed no publication bias.

Forest plots of psychological treatments vs. TAU/WL on each symptom outcome are presented in Supplementary Figures 1–5.




Figure 2. Funnel plot of comparisons on outcomes of PTSD, depression, and anxiety.

We conducted sensitivity analysis by comparing the random effect size by removing each single comparison. First, we analyzed based on outcomes that had more than five comparisons. Comparing psychological treatments to TAU/WL, for PTSD symptoms ($n = 32$, $g = .936$, 95% CI .712 to 1.161, $I^2 = 84.546$, $\tau^2 = .320$), effect size was maintained when removing any comparison ($g = .871$ –.967). Using Trim and Fill, effect size and 95% confidence interval values were unchanged, leaving out publication bias. Also, for depression ($n = 31$, $g = .688$, 95% CI .511 to .864, $I^2 = 73.293$, $\tau^2 = .171$), removing any comparison had little impact on the effect size ($g = .637$ –.710). By Trim and Fill, effect size and 95% confidence interval values remained unchanged. This was the same for anxiety symptoms ($n = 12$, $g = .699$, 95% CI .435 to .963, $I^2 = 79.894$, $\tau^2 = .161$), effect size range after each removal ($g = .593$ –.754), and effect size before and after Trim and Fill were maintained.

For negative self-concept ($n = 6$, $g = .762$, 95% CI .306 to 1.218, $I^2 = 68.884$, $\tau^2 = .219$), effect size range after each removal ($g = .522$ –.847) was maintained. For dissociation ($n = 6$, $g = .857$, 95% CI .539 to 1.179, $I^2 = 43.627$, $\tau^2 = .069$), there were some changes in effect size range after some removal ($g = .754$ –.968); removal of studies with BPD participants increased the effect size to .936 and .968, respectively.

Next, we conducted sensitivity analyses based on each psychological treatment. Among comparisons of CPT/CT and TAU/WL, removing two studies that provided less than four sessions, effect size for PTSD symptoms ($n = 7$, $g = 1.348$, 95% CI .805 to 1.891), and depression ($n = 7$, $g = 1.383$, 95% CI .767 to 2.000) increased. Among comparisons of NET and TAU/WL, removing a study that provided less than four sessions, effect size for PTSD symptoms ($n = 7$, $g = 1.103$, 95% CI .570 to 1.636) and depression ($n = 7$,

$g = .702$, 95% CI .399 to 1.004) increased. Additionally, removing a study with BPD participants, effect size for PTSD symptoms ($n = 6$, $g = 1.252$, 95% CI .710 to 1.794) and depression ($n = 6$, $g = .716$, 95% CI -.391 to 1.042) increased. Among comparisons of EMDR and TAU/WL, removing a study that provided two sessions decreased the effect size of PTSD symptoms ($n = 3$, $g = .655$, 95% CI .102 to 1.207), and depression ($n = 3$, $g = .277$, 95% CI -.129 to .683). We could not remove a BPD participant study in phased TF-CBT vs. TAU/WL comparison because there were too few comparisons.

Univariate meta-regression analyses

We selected comparisons of psychological treatments to TAU/WL and did meta-regression analyses on outcome symptoms of PTSD and depression. Other outcome symptoms did not have enough comparisons for meta-regression and thus could not be examined. We did univariate analyses of study characteristics. The type of trauma (childhood abuse vs. organized violence, childhood abuse vs. multiple interpersonal trauma), compositions of low/insecure SES (%), female gender (%), European/Caucasian ethnicity (%), studies mainly with low-resource populations, session length, intent-to-treat (vs. completer) study, and risk of bias levels (high vs. low, high vs. moderate) each did not predict efficacy on PTSD or depression. Trauma-focused treatment (vs. non-trauma focused) predicted efficacy for both PTSD ($n = 31$, $R^2 = .18$, coefficient = .639, 95% CI = .042–1.240, $Z = 2.10$, $p < .05$) and depression ($n = 31$, $R^2 = .19$, coefficient = .426, 95% CI = .015–.837, $Z = 2.04$, $p < .05$).

Discussion

This meta-analysis found 42 RCT studies indicating that psychological treatments promise efficacy for people with complex trauma in various degrees across constructs of CPTSD. Most studies were conducted in American or European regions. Compared to PTSD or depression studies, studies targeting DSO symptoms, dissociation, comorbid BPD, or self-harm behaviors were scarce. No studies included low intelligence participants. These symptoms were more likely to be included in the exclusion criteria, as previous discussions point out (Bradley, Greene, Russ, Dutra, & Westen, 2005; Leeman et al., 2017). There were more female par-

ticipant studies than male, and participant ethnicity was diverse. Some studies reported SES information of participants, however, more precise reports may be helpful for future analysis.

Our meta analyses showed that on PTSD symptoms of complex trauma, psychological treatments such as CPT/CT, NET, phased TF-CBT, and single TF-CBT had large effect size, and PE, EMDR, and PCT showed moderate effect size. This result supports the idea that CPTSD treatment may benefit building from current PTSD treatments (ISTSS Guidelines Committee, 2018; Karatzias et al., 2019). For depression, CPT/CT, phased TF-CBT, and single TF-CBT showed large effect size, NET and PE moderate effect size, and EMDR, non-TF-CBT, and PCT small effect size. CBT based trauma-focused treatments seemed to work favorably on depressive symptoms of complex trauma.

That PCT showed similar effect size to PE and EMDR on PTSD, showing that it may reduce complex trauma PTSD, indicates that PCT may be an option when trauma-focused therapies cannot be adjusted because of client preferences or accessibility of intervention. Nonetheless, our meta-regression revealed that trauma-focused interventions compared to PCT or non-trauma focused interventions had higher efficacy on PTSD and depression of complex trauma. This supports previous studies reporting the superiority of trauma-focused interventions in chronic PTSD (Bisson, Roberts, Andrew, Cooper, & Lewis, 2013), organized trauma (ter Heide, Mooren, & Kleber, 2016), military PTSD (Haagen, Smid, Knipscheer, & Kleber, 2015), and child abuse related PTSD (Ehring et al., 2014).

Considering CPT/CT and NET, the number of sessions (over four) seemed to increase the efficacy for symptoms of PTSD and depression; however, this was not true for EMDR. It is hard to conclude, since there were few comparisons; however, BPD symptoms seemed to negatively influence the efficacy of interventions. Previous quantitative review of treatments for child abuse CPTSD (defined as PTSD plus disorders of extreme stress or PTSD with personality disorders) concluded that exposure promised short-term improvements; however, within intent-to-treat analysis, affect modulation rather than exposure brought more favorable outcomes, including less dropout (Dorrepal et al., 2014). Therefore, complicated presentations, such as comorbidity of personality disorders related with complex trauma may require additional inter-

ventions, such as skills training prior to exposure. Skills training may aid the exposure process and maintain the efficacy of exposure by changing efficacy moderators, such as dissociation (e.g., Cloitre, Petkova, Wang, & Lu, 2012; Kleindienst et al., 2016) or emotion dysregulation (e.g., Hien, Lopez-Castro, Papini, Gorman, & Ruglass, 2017) frequent in complex trauma.

Among DSO symptoms, for negative self-concept, large effect size was found in CPT/CT, and moderate effect size in PE and PCT. For emotion regulation and interpersonal problems, comparisons were insufficient to reach any conclusions. Since phased TF-CBT interventions seemed to promise large efficacy, it may also be true for DSO symptoms that dealing with both trauma material and training for regulation strategies are beneficial. Not only for processing trauma memories but also for mastering how to regulate oneself and deal with outside world would lead to adjustments in current self and interpersonal life. However, we cannot confirm without more evidence using well agreed instruments for DSO symptoms. Further studies targeting DSO symptoms may find out whether this efficacy comes from sequential interventions or combinations of skills training and exposure.

Among 42 studies, 57.1% reported numbers to code intent-to-treat data. Meta-regression showed that intent-to-treat data did not affect the effect size. Because maintaining treatment completion is an important task for CPTSD treatment, future interventions reporting intent-to-treat data would provide rich evidence.

Average treatment completer rate from all studies was 75.9%, and average treatment completion rates for each intervention were as followed; CPT/CT was 69.8%, NET was 82.4%, PE was 61.5%, phased TF-CBT was 76.1%, EMDR was 85.9%, single TF-CBT was 82.1%, non-TF-CBT was 67.8%, and PCT was 79.8%. In a previous meta-analysis of dropout rates for PTSD treatments, average dropout rate was reported to be 18.3%, including dropout rate of trauma-specific treatments as 36%, whereas for present-centered treatment the dropout rate was 22% (Imel, Laska, Jakupcak, & Simpson, 2013), showing rates similar to those in our report. However, in our study, fluctuations in comparisons exist. It should be mentioned that in our study, unlike the PTSD study of Imel et al. (2013), non-trauma focused, present-centered treatments did not show a superior completion rate compared to trauma-focused treatments. The differences in completion rate across studies require further evidence

seeking participant or context-related moderators that influence attrition, which will support evidence-based decision-making in practice.

Six studies focused on active treatment comparisons; for instance, adding cognitive restructuring to exposure did not change efficacy on PTSD. However, adding skills training to exposure seemed to decrease DSO, also, on anxiety and interpersonal problems large to moderate efficacy was shown, and on PTSD, depression, and emotion regulation small efficacy was shown. Adding exposure to skills training showed moderate efficacy for emotion regulation and interpersonal problems, and small efficacy for PTSD, depression, anxiety, dissociation, and suicidality, but showed similar results in NSSI. From this evidence we may suggest that for complex traumas, multiple interventions, such as skills training along with trauma-focused exposure treatments, may have more benefits on DSO than does single intervention. However, as mentioned above, evidence is scarce to conclude whether this means that phase-based treatments are superior to non-phased treatments in CPTSD.

Since lack of social resources may influence trajectories of trauma recovery (e.g., Hall, Murray, Galea, Canetti, & Hobfoll, 2015; Miller & Rasmussen, 2010), it was important in our study to seek social conditions of participants as moderators of efficacy. Most studies were conducted in North America and Europe. Further evidence from diverse cultural regions is in need. We found that studies were evenly conducted among various ethnicities, and ethnicity had no influence on PTSD and depression efficacy outcomes in the meta-regression. Also, SES of participants did not affect treatment efficacy on PTSD and depression, which is in line with previous discussions that prolonged economic loss may have little effect on the imminent course of PTSD (Hall et al., 2015).

However, given that SES of participants was mostly not clearly reported, and that we did not use robust indicators for SES, it is difficult to draw conclusions. In addition, social conditions influencing efficacy on DSO symptoms could not be analyzed, bringing attention to the concern that PTSD focused treatment would overlook various difficulties and the effect of daily psychological distress of social conditions (Miller & Rasmussen, 2010). Future studies may help by seeking social conditions as moderators on outcomes of DSO. Advocacy and mentoring were adapted in interventions for interpersonal abuse; however, outcome measure-

ments were limited and showed no significant efficacy. We may mention that some studies of non-trauma focused brief interventions were conducted with participants of diverse ethnicity with unstable social economic resources, implying that people with the least resources may also have the fewest treatment opportunities. Compared to these studies, studies adjusting trauma-focused intervention to low-resource populations showed strong efficacy. Therefore, gathering resources to provide both trauma-focused interventions and consistent psychosocial resource support for marginalized people should be considered.

Most studies dealt with interpersonal trauma including child abuse; however, the type of trauma did not affect treatment efficacy. Previous meta regression reported that childhood onset of trauma is related to a smaller effect (Karatzias et al., 2019); so, more evidence is needed to conclude whether it is the onset age that matters, or cumulative aspect or loss of resources is what is associated.

Overall assessment of risk of bias concluded that 59.5% of the studies had low to moderate risk; however, high risk studies were up to 40%, reflecting the limitations of conducting a rigor methodology trial with complex trauma experienced people. A previous analysis reported that rigorous methodology was related to higher efficacy in CPTSD (Karatzias et al., 2019); however, in our study, having a high risk did not affect efficacy on PTSD and depression. In our study, PCT studies tend to have high risks; so, according to previous findings, lowering the risk of bias in PCT studies may change its efficacy.

We have limitations to consider. We included studies dealing with complex trauma and did not limit to CPTSD diagnosis. Therefore, we cannot conclude that our results indicate efficacy for ICD-11 CPTSD; however, we sought important evidence dealing with PTSD and the pervasive symptoms, including DSO, of complex trauma. Limitations are that meta-regression focused on only PTSD and depression, given that DSO symptoms or dissociation were measured in less than 16.6% of the studies. Further studies targeting CPTSD symptoms with validated instruments will improve future discussions in CPTSD treatment. Also, we excluded group therapies. Previous meta-analyses found that for PTSD, individual therapies were more effective than were group approaches (Ehrings et al., 2014); however, for CPTSD, no such evidence was found (Karatzias et al., 2019). Future research may ana-

lyze how group therapy would be effective in DSO symptoms. In study selection and coding procedures, four researchers were involved to minimize possible errors or omissions; however, the limitation is that the intercoder reliability was not calculated. Also, excluding non-peer reviewed articles hindered encompassing grey literature, giving a possibility of publication bias, so effect size may be less accurate (Conn, Valentin, Cooper, & Rantz, 2003). However, including grey literature in psychiatry research also have risks (Martin, Pérez, Sacristán, & Álvarez, 2005), so future research may consider including grey literature, only with sensitivity analysis supporting the evidence. Lastly, adverse effects of treatments were not examined, and further study should address evidence of possible distress following therapeutic gain.

Despite its limitations, this study synthesized the evidence in the treatment of complex trauma. Implications suggest that current PTSD interventions are equally beneficial for people with complex trauma, and a combination of skills training and exposure in a phase-oriented way seems promising. More evidence is in need to identify which intervention for whom, when, in which order, and for how long would be effective for specific profiles of CPTSD, and we may expect future advances and moderations in current treatments to improve efficacy on CPTSD.

Supplemental materials

Supplemental materials are available at <https://doi.org/10.15842/kjcp.2020.39.2.007>.

Author contributions statement

HJC, assistant professor in the Department of Psychology at Chungbuk National University, as the principal investigator of the research grant, designed the research, participated in selection and coding, analyzed data, and prepared the manuscript. WYL, assistant professor in the Department of Psychology at Chungbuk National University, supervised data coding and statistical analyses, and prepared the manuscript. SYH, a certified mental health professional, and JHK, a certified mental health professional who is now a doctoral student at Chungbuk National University, both selected, assessed, and coded research data. All authors provided critical feedback, participated in preparing the manuscript, and approved the final submission.

References

- American Psychological Association. (2017). *Clinical practice guideline for the treatment of posttraumatic stress disorder (PTSD) in adults*. Washington, DC: Author.
- Becker, B. J. (1990). Coaching for the scholastic aptitude test: Further synthesis and appraisal. *Review of Educational Research*, 60, 373-417.
- Bisson, J. I., Roberts, N. P., Andrew, M., Cooper, R., & Lewis, C. (2013). Psychological therapies for chronic post-traumatic stress disorder (PTSD) in adults. *Cochrane Database of Systematic Reviews*, 2013, CD003388.
- Bradley, R., Greene, J., Russ, E., Dutra, L., & Westen, D. (2005). A multidimensional meta-analysis of psychotherapy for PTSD. *American Journal of Psychiatry*, 162, 214-227.
- Brewin, C. R., Cloitre, M., Hyland, P., Shevlin, M., Maercker, A., Bryant, R. A., . . . Reed, G. M. (2017). A review of current evidence regarding the ICD-11 proposals for diagnosing PTSD and complex PTSD. *Clinical Psychology Review*, 58, 1-15.
- Briere, J. N. (1992). *Child abuse trauma: Theory and treatment of the lasting effects*. Thousand Oaks, CA: Sage Publications.
- Cloitre, M., Courtois, C., Ford, J., Green, B., Alexander, P., Briere, J., . . . van der Hart, O. (2012). *The ISTSS Expert Consensus Treatment Guidelines for Complex PTSD in Adults*. Retrieved from http://www.istss.org/AM/Template.cfm?Section=ISTSS_Complex_PTSD_Treatment_Guidelines&Template=/CM/ContentDisplay.cfm&ContentID=5185
- Cloitre, M., Garvert, D. W., Brewin, C. R., Bryant, R. A., & Maercker, A. (2013). Evidence for proposed ICD-11 PTSD and complex PTSD: A latent profile analysis. *European Journal of Psychotraumatology*, 4, 20706.
- Cloitre, M., Garvert, D. W., Weiss, B., Carlson, E. B., & Bryant, R. A. (2014). Distinguishing PTSD, complex PTSD, and borderline personality disorder: A latent class analysis. *European Journal of Psychotraumatology*, 5, 25097.
- Cloitre, M., Petkova, E., Wang, J., & Lu Lassell, F. (2012). An examination of the influence of a sequential treatment on the course and impact of dissociation among women with PTSD related to childhood abuse. *Depression and Anxiety*, 29, 709-717.
- Cloitre, M., Shevlin, M., Brewin, C., Bisson, J. I., Roberts, N. P., Maercker, A., . . . Hyland, P. (2018). The International Trauma Questionnaire: Development of a self-report measure of ICD-11 PTSD and complex PTSD. *Acta Psychiatrica Scandinavica*, 138, 536-546.
- Cohen, J. (1988). *Statistical power analysis for the behavioral science* (2nd ed.). Hillsdale, NJ: Erlbaum.
- Conn, V. S., Valentine, J. C., Cooper, H. M., & Rantz, M. J. (2003). Grey literature in meta-analysis. *Nursing Research*, 52, 256-261.
- Cooper, H. M. (1998). *Synthesizing research: A guide for literature reviews* (3rd ed.). Thousand Oaks, CA: Sage.
- Cooper, H. M. (2010). *Research synthesis and meta-analysis: A step by step approach*. New York, NY: Sage.
- Courtois, C. A., & Ford, J. D. (2013). *Treatment of complex trauma: A sequenced, relationship-based approach*. New York, NY: Guilford Press.
- De Jongh, A., Resick, P. A., Zoellner, L. A., van Minnen, A., Lee, C. W., Monson, C. M., . . . Bicanic, I. A. E. (2016). Critical analysis of the current treatment guidelines for complex PTSD in adults. *Depression and Anxiety*, 33, 359-369.
- Dorrepal, E., Thomaes, K., Hoogendoorn, A. W., Veltman, D. J., Draijer, N., & van Balkom, A. J. L. M. (2014). Evidence-based treatment for adult women with child abuse-related complex PTSD: A quantitative review. *European Journal of Psychotraumatology*, 5, 23613.
- Duval, S., & Tweedie, R. (2000). Trim and fill: A simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics*, 56, 455-463.
- Ehlers, A., Clark, D. M., Hackmann, A., McManus, F., & Fennell, M. (2005). Cognitive therapy for post-traumatic stress disorder: Development and evaluation. *Behaviour Research and Therapy*, 43, 413-431.
- Ehring, T., Welboren, R., Morina, N., Wicherts, J. M., Freitag, J., & Emmelkamp, P. M. (2014). Meta-analysis of psychological treatments for posttraumatic stress disorder in adult survivors of childhood abuse. *Clinical Psychology Review*, 34, 645-657.
- Foa, E. B., & Rothbaum, B. O. (1998). *Treating the trauma of rape: Cognitive-behavioral therapy for PTSD*. New York, NY: Guilford Press.
- Haagen, J. F., Smid, G. E., Knipscheer, J. W., & Kleber, R. J. (2015). The efficacy of recommended treatments for veterans with PTSD: A metaregression analysis. *Clinical Psychology Review*, 40, 184-194.
- Hall, B. J., Murray, S. M., Galea, S., Canetti, D., & Hobfoll, S. E. (2015). Loss of social resources predicts incident posttraumatic stress disorder during ongoing political violence within the Palestinian Authority. *Social Psychiatry and Psychiatric Epidemiology*, 50, 561-568.
- Herman, J. L. (1992). Complex PTSD: A syndrome in survivors of prolonged and repeated trauma. In M. J. Horowitz (Ed.), *Essential papers on post traumatic stress disorder*. New York, NY: New York University Press.
- Herman, J. L. (1997). *Trauma and recovery: The aftermath of violence-from domestic abuse to political terror*. New York, NY: Basic Books.
- Hien, D. A., Lopez-Castro, T., Papini, S., Gorman, B., & Ruglass, L. M. (2017). Emotion dysregulation moderates the effect of cognitive behavior therapy with prolonged exposure for co-occurring PTSD and substance use disorders. *Journal of Anxiety Disorders*,

- 52, 53-61.
- Hobfoll, S. E., Mancini, A. D., Hall, B. J., Canetti, D., & Bonanno, G. A. (2011). The limits of resilience: Distress following chronic political violence among Palestinians. *Social Science & Medicine*, 72, 1400-1408.
- Imel, Z. E., Laska, K., Jakupcak, M., & Simpson, T. L. (2013). Meta-analysis of dropout in treatments for posttraumatic stress disorder. *Journal of Consulting and Clinical Psychology*, 81, 394-404.
- Institute of Medicine. (2011). *Finding What Works in Health Care: Standards for Systematic Reviews*. Washington, DC: The National Academies Press.
- ISTSS Guidelines Committee. (2018). *ISTSS Guidelines Position Paper on Complex PTSD in Adults*. Oakbrook Terrace, IL: Author.
- Jang, D., & Shin, I. (2011). Historical development of meta-analysis as an educational research methodology. *Journal of Curriculum and Evaluation*, 14, 309-332.
- Karatzias, T., Murphy, P., Cloitre, M., Bisson, J., Roberts, N., Shevlin, M., . . . Hutton, P. (2019). Psychological interventions for ICD-11 complex PTSD symptoms: Systematic review and meta-analysis. *Psychological Medicine*, 49, 1-15.
- Kleindienst, N., Priebe, K., Görg, N., Dyer, A., Steil, R., Lyssenko, L., . . . Bohus, M. (2016). State dissociation moderates response to dialectical behavior therapy for posttraumatic stress disorder in women with and without borderline personality disorder. *European Journal of Psychotraumatology*, 7, 30375.
- Knefel, M., Garvert, D. W., Cloitre, M., & Lueger-Schuster, B. (2015). Update to an evaluation of ICD-11 PTSD and complex PTSD criteria in a sample of adult survivors of childhood institutional abuse by Knefel & Lueger-Schuster (2013): A latent profile analysis. *European Journal of Psychotraumatology*, 6, 25290.
- Leeman, R. F., Hefner, K., Frohe, T., Murray, A., Rosenheck, R. A., Watts, B. V., & Sofuoglu, M. (2017). Exclusion of participants based on substance use status: Findings from randomized controlled trials of treatments for PTSD. *Behaviour Research and Therapy*, 89, 33-40.
- Levine, P. A. (2010). *In an unspoken voice: How the body releases trauma and restores goodness*. Berkeley, CA: North Atlantic Books.
- Linehan, M. (1993). *Cognitive-Behavioral Treatment of Borderline Personality Disorder*. New York, NY: Guilford Press.
- Martin, J. L. R., Pérez, V., Sacristán, M., & Álvarez, E. (2005). Is grey literature essential for a better control of publication bias in psychiatry? An example from three meta-analyses of schizophrenia. *European Psychiatry*, 20, 550-553.
- Miller, K. E., & Rasmussen, A. (2010). War exposure, daily stressors, and mental health in conflict and post-conflict settings: Bridging the divide between trauma-focused and psychosocial frameworks. *Social Science & Medicine*, 70, 7-16.
- Morris, S. B. (2008). Estimating effect sizes from pretest-posttest-control group designs. *Organizational Research Methods*, 11, 364-386.
- National Institute for Health and Care Excellence (NICE). (2012). *The guidelines manual*. London: National Institute for Health and Care Excellence.
- Ogden, P., & Fisher, J. (2015). *Sensorimotor psychotherapy: Interventions for trauma and attachment (Norton series on interpersonal neurobiology)*. New York, NY: WW Norton & Company.
- Pelcovitz, D., van der Kolk, B. A., Roth, S., Mandel, E. S., Kaplan, S., & Resick, P. (1997). Development of a criteria set and a structured interview for disorders of extreme stress (SIDES). *Journal of Traumatic Stress*, 10, 3-16.
- Resick, P. A., & Schnicke, M. K. (1992). Cognitive processing therapy for sexual assault victims. *Journal of Consulting and Clinical Psychology*, 60, 748.
- Seidler, G. H., & Wagner, F. E. (2006). Comparing the efficacy of EMDR and trauma-focused cognitive-behavioral therapy in the treatment of PTSD: A meta-analytic study. *Psychological Medicine*, 36, 1515-1522.
- Shapiro, F. (1995). *Eye movement desensitization and reprocessing: Basic principles, protocols, and procedures*. New York, NY: Guilford Press.
- ter Heide, F. J. J., Mooren, T. M., & Kleber, R. J. (2016). Complex PTSD and phased treatment in refugees: A debate piece. *European Journal of Psychotraumatology*, 7, 28687.
- Terr, L. (1990). *Too scared to cry*. New York, NY: Basic Books.
- van der Kolk, B. A. (1996). The complexity of adaptation to trauma: Self-regulation, stimulus discrimination, and characterological development. In B. A. van der Kolk, A. C. McFarlane, & L. Weisæth (Eds.), *Traumatic stress: The effects of overwhelming experience on mind, body, and society* (pp.182-213). New York, NY: Guilford Press.
- World Health Organization. (2018). *International Classification of Disorders, 11th edition*. Geneva: World Health Organization.

Appendix I. Study and Population Characteristics of Selected Studies

Study	Main intervention	Trauma-focused	Treatment arms	Average structured individual session (min. per session)	N	Female (%)	Low social economic status (%)	European/Caucasian (%)	Pre n	Completer n	Treatment completor rate	F/U months
Adenauer 2011	NET	Y	NET	12 (108)	34	48.8	87.5	0	16	15	93.8	0
Bichescu 2007	NET	Y	WL			44.4	100	0	18	16	88.9	
			NET	5 (120)	18	0	-	100	9	9	100	6
			Psychoeducation	1		11.1	-	100	9	9	100	
Bohus 2013	DBT-PTSD	Y	DBT-PTSD	23 (45)	74	100	-	-	36	29	80.6	6
			TAU/WL			100	-	-	38	29	76.3	
Bolton 2014a	BA	N	BA	12	281	57	50	0	114	82	71.9	6
	CPT	Y	CPT	12		58.4	48	0	101	67	66.3	
			WL			59.1	56	0	66	53	80.3	
Bolton 2014b	Common elements	Y	CETA	9.7 (60)	347	61	59.9	0	182	148	81.3	0
	Treatment approach (CETA)		WL			64.2	58.8	0	165	126	76.4	
Chard 2005	CPT	Y	CPT	10 (60)	71	100	41	81.4	36	28	77.8	12
			WL			100	41	81.4	35	27	77.1	
Cloitre 2002	STAIR-PE	Y	STAIR-PE	16	58	100	31	46	31	22	71	0
			WL			100	31	46	27	24	88.9	
Cloitre 2010	STAIR-PE	Y	STAIR-exposure	16	104	100	32.3	37	33	28	84.8	6
		N	STAIR-support	16		100	28.1	37	38	28	73.7	
		Y	Support-exposure	16		100	46.4	33	33	20	60.6	
Duffy 2007	CBT	Y	CBT	12	58	34.5	-	100	29	20	69	4
			WL			44.8	-	100	29	29	100	
Edmond 1999	EMDR	Y	EMDR	6 (90)	59	100	38	85	20	20	100	3
		N	TAU	6		100	38	85	20	19	95	
			WL			100	38	85	19	13	68.4	
Foa 2005	PE	Y	PE	9 (90-120)	179	100	47.4	49.2	79	52	65.8	12
	PE/cognitive restructuring	Y	PE/cognitive restructuring	9 (90-120)		100	47.4	49.2	74	44	59.5	
			WL			100	47.4	49.2	26	25	96.2	
Galovski 2012	CPT	Y	CPT	7.5	100	-	67	42	53	33	62.3	3
			WL			-	67	42	47	37	78.7	
Ghafoori 2016	Psychoeducation	N	Psychoeducation	1 (90)	86	52.6	89.2	29.7	38	37	97.4	1
			WL			20.8	86.7	23.3	48	30	62.5	
Ghafoori 2017	PE	Y	PE	6.79 (60-90)	71	83	61.7	36.2	47	25	53.2	0
			Present-centered	8.67 (60-90)		83.3	66.7	12.5	24	18	75	
Harkness 2012	CBT	Y	CBT	16.56	203	63.5	-	-	70	47	67.1	12
	IPT	N	IPT	17.15		63.5	-	-	64	50	78.1	
	Antidepressant					63.5			69	43	62.3	

(Continued to the next page)

Appendix I. Continued

Study	Main intervention	Trauma-focused	Treatment arms	Average structured individual session (min. per session)	N	Female (%)	Low social economic status (%)	European/Caucasian (%)	Pre n	Completer n	Treatment completer rate	F/U months
Harned 2014	DBT-PE	Y	DBT-PE	38.6 (60)	26	100	75	80.8	17	10	58.8	3
Hensel-Dittmann 2011	DBT	N	DBT	28.8 (120–150)		100	75	80.8	9	5	55.6	3
Hijazi 2014	NET	Y	NET	10 (90)	28	-	89.3	2	15	12	80	12
Johnson 2011	NET	N	Stress inoculation	10 (90)		-	89.3	2	13	11	84.6	12
	NET	Y	NET	3 (60–90)	63	63.4	100	0	41	39	95.1	4
	WL	N	WL			40.9	100	0	22	21	95.5	
	Stabilization	N	Stabilization (HOPE)+standard shelter service	6.8 (60–90)	70	100	62.9	48.6	35	19	54.3	6
Jung 2013	CRIM (CT)	N	Standard shelter service			100	82.9	37.1	35	33	94.3	
		Y	Cognitive restructuring and imagery modification	2 (50–90)	34	100	-	89.3	17	14	82.4	1
Katz 2014	PE	Y	WL			100	-	89.3	17	14	82.4	
	Holographic reprocessing	N	PE	10	51	100	71	41	17	10	58.8	0
		N	Holographic reprocessing	10		100	47	41	17	16	94.1	
		N	Person centered	10		100	41	47	17	11	64.7	
Korte 2017	TF-CBT	Y	Exposure (COPE)	12 (90)	81	7.4	d/k	68.5	54		-	0
		N	Relapse prevention (CBT)	12 (90)		14.8	d/k	44.4	27		-	
Kubany 2003	CT	Y	Cognitive therapy	8.5 (90)	37	100	d/k	48.6	19	18	94.7	3
		N	WL			100	d/k	48.6	18			
McDonagh 2005	PE	Y	PE/cognitive restructure	14 (90–120)	74	100	42	90	29	17	58.6	6
	present-centered	N	Present-centered	14 (90–120)		100	50	95	22	20	90.9	
		N	WL			100	30	96	23	20	87	
Neuner 2010	NET	Y	NET	8.79 (120)	32	31.3	100	0	16	14	87.5	6
		N	TAU	-		31.3	100	0	16	16	100	
Nixon 2016	CPT	Y	CPT	3.5 (90)	47	92	26	83	25	15	60	12
		N	TAU	3.5		100	18	91	22	17	77.3	
Pabst 2014	NET	Y	NET	17.2 (90)	22	100	-	-	11	9	81.8	12
		N	TBE	14.4 (90)		100	-	-	11	8	72.7	
Paivio 2010	Emotion-focused (Imaginal Confrontation, IC)	N	Emotion-focused (IC)	16.9 (60)	56	50	20	95	27	20	74.1	6
		N	EF w/o imaginal confrontation	16.8 (60)		56	8	84	28	25	89.3	
Pigeon 2009	IPT	N	IPT	12	70	100	65	62	37	28	77	0
		N	TAU	6.3		100	73	54	33	19	58	
Resick 2002	CPT	Y	CPT	12 (60–90)	171	100	-	71	62	41	66.1	9
	PE	Y	PE	12 (60–90)		100	-	71	62	40	64.5	
		N	WL			100	-	71	47	40	85.1	

(Continued to the next page)

Appendix I. Continued

Study	Main intervention	Trauma-focused	Treatment arms	Average structured individual session (min. per session)	N	Female (%)	Low social economic status (%)	European/Caucasian (%)	Pre n	Completer n	Treatment completor rate	F/U months
Resick 2008	CPT	Y	CPT	12 (60)	162	100	79	62	56	27	48.2	6
	CPT without WA	N	CPT-C	12 (60)		100	46	62	51	29	56.9	
	Written accounts	Y	WA	12 (60-120)		100	42	62	55	30	54.5	
Scheck 1998	EMDR	Y	EMDR	2 (90)	67	100	-	62	34	30	88.2	3
		N	TAU	2 (90)		100	-	62	33	30	90.9	
Schnurr 2007	PE	Y	PE	10 (90)	284	100	37.6	56	141	88	62.4	6
		N	Present-centered	10 (90)		100	39.2	53	143	113	79	
Stenmark 2013	NET	Y	NET	10 (90)	81	33.3	100	0	51	33	64.7	6
		N	TAU	10 (90)		26.7	100	0	30	21	70	
Sullivan 1999	advocacy	N	Advocacy	20 (120)	278	100	59	42	143	135	94.4	24
			Control			100	59	42	135	130	96.3	
Suris 2013	CPT	Y	CPT	9.7	86	82.7	35	44	52	34	65.4	6
		N	Present-centered	10.5		88.2	41	44	34	28	82.4	
Taft 2011	Mentoring	N	Mentor+clinician_care		174	100	75.2	67.3	113	90	79.6	12
			Usual clinician care			100	70.5	60.7	61	43	70.5	
Talbot 2011	IPT	N	IPT	12.9	70	100	59	62	37	31	83.8	9
		N	TAU	6.3		100	58	55	33	22	66.7	
ter Heide 2011	EMDR	Y	EMDR	11 (90)	20	50	70	0	10	5	50	3
			Stabilization	11 (60)		30	70	0	10	5	50	
Tiwari 2010	Advocacy	N	Advocacy+usual service	12 (30)	200	100	30	0	100	100	100	9
			Usual service			100	32	0	100	100	100	
Vitriol 2009	Psychodynamic	N	Psychodynamic	12	87	100	-	0	44	33	75	6
		N	Standard treatment			100	-	0	43	23	53.5	
Weiss 2015	Common elements treatment approach (CETA)	Y	CETA	9.94 (50-60)	149	32.3	61.7	0	99	97	98	4
			WL			28	50	0	50	50	100	
	CPT	Y	CPT	12	193	32.6	57.3	0	129	107	82.9	5
			WL			37.5	54.7	0	64	64	100	

Note. BA = behavioral activation treatment; CBT = cognitive behavioral therapy; CETA = common elements treatment approach; COPE = Concurrent Treatment of PTSD and Substance Use Disorders Using Prolonged Exposure; CPT = cognitive processing therapy; CPT-C = Cognitive processing therapy for posttraumatic stress disorder; CRIM = Cognitive restructuring and imagery modification; CT = cognitive therapy; DBT = dialectical behavior therapy; DBT-PTSD = dialectical behavior therapy for posttraumatic stress disorder; EMDR = eye movement desensitization and reprocessing; EF w/o imaginal confrontation = emotion focused without imaginal confrontation; Emotion-focused (IC) = emotion-focused (Imaginal Confrontation); HOPE = Helping to Overcome PTSD through Empowerment; IPT = interpersonal psychotherapy; NET = narrative exposure therapy; PE = prolonged exposure; STAIR-PE = skills training in affect and interpersonal regulation-prolonged exposure; STAIR = skills training in affect and interpersonal regulation; TAU = treatment as usual; TBE = Treatment by Experts for Borderline Personality Disorder; TF-CBT = Trauma-Focused Cognitive Behavioral Therapy; WA = written accounts; WL = waiting list.

Appendix II. Study Characteristics and Risk of Bias Assessment of Selected Studies

Study	Main intervention	Outcome	Instrument	Trauma type	Study location	Setting	Phase-oriented	Intent-to-treat	Risk of bias
Adenauer 2011	NET	Depression	HAM-D	Organized violence (conflict/torture/detention/persecution)	Germany	Outpatient	N	N	Low
Bichescu 2007	NET	Depression	BDI	Organized violence (conflict/torture/detention/persecution)	Romania	Outpatient	N	Y	Moderate
Bohus 2013	DBT-PTSD	BPD Depression Dissociation PTSD	BSL BDI-II DES CAPS	Child abuse (under the age 18)	Germany	Residential	Y	Y	Low
Bolton 2014a	BA CPT	Anxiety Depression PTSD	HSCL-25 HSCL-25 HTQ	Organized violence (conflict/torture/detention/persecution)	Kurdistan	Outpatient	N	Y	Moderate
Bolton 2014b	Common elements treatment approach (CETA)	Anxiety Depression PTSD	HSCL-25 HSCL-25 HTQ	Organized violence (conflict/torture/detention/persecution)	Thailand	Outpatient	N	Y	Low
Chard 2005	CPT	Depression Dissociation PTSD	BDI-II DES-II CAPS-SX	Child abuse (average age of onset = 6.4 years)	USA	Outpatient	N	N	High
Cloitre 2002	STAIR-PE	Anxiety Depression Dissociation Emotion regulation Interpersonal = problem PTSD	STAI-S BDI DISS NMR IIP CAPS	Child abuse (under the age 18)	USA	Outpatient	Y	N	Moderate
Cloitre 2010	STAIR-PE	Anxiety Depression Emotion regulation Interpersonal = problem PTSD	STAI-S BDI NMR IIP CAPS	Child abuse (under the age 18)	USA	Outpatient	Y	Y	Low
Duffy 2007	CBT	Depression PTSD	BDI PDS	Organized violence (conflict/torture/detention/persecution)	Ireland	Outpatient	N	Y	High
Edmond 1999	EMDR	Anxiety Depression PTSD	STAI BDI IES	Child abuse (average age of onset = 6.5 years)	USA	Outpatient	N	N	High
Foa 2005	PE PE/cognitive restructuring	Depression PTSD	BDI PSS-I	Multiple interpersonal trauma	USA	Outpatient	N	Y	Low
Galovski 2012	CPT	Depression PTSD	BDI-II CAPS	Multiple interpersonal trauma	USA	Outpatient	N	Y	Low
Ghafoori 2016	psychoeducation	Anxiety Depression PTSD	BSI-18 BSI-18 PCL-C	Multiple interpersonal trauma	USA	Outpatient	N	N	High
Ghafoori 2017	PE	Anxiety Depression PTSD	BSI-18 BSI-18 PCL-5	Multiple interpersonal trauma	USA	Outpatient	N	Y	High
Harkness 2012	CBT IPT	Depression	HAM-D	Child abuse (under the age 17)	Canada	Outpatient	N	Y	Low

(Continued to the next page)

Appendix II. Continued

Study	Main intervention	Outcome	Instrument	Trauma type	Study location	Setting	Phase-oriented	Intent-to-treat	Risk of bias
Harned 2014	DBT-PE DBT	Anxiety Depression Dissociation NSSI PTSD Suicidality	HRSA HRSD DES-T SASII PSS-I SASII	Multiple interpersonal trauma	USA	Outpatient	Y	Y	Low
Hensel-Dittmann 2011	NET	Depression PTSD	HAM-D CAPS	Organized violence (conflict/torture/detention/persecution)	Germany	Outpatient	N	N	Moderate
Hijazi 2014	NET	Depression PTSD	BDI-II HTQ	Organized violence (conflict/torture/detention/persecution)	Germany	Outpatient	N	Y	Moderate
Johnson 2011	Stabilization	Depression PTSD	BDI CAPS	Multiple interpersonal trauma	USA	Residential	N	Y	High
Jung 2013	CRIM (CT)	Depression PTSD Self	BDI-II CAPS Rosenberg SES	Child abuse (average age of onset = 7.7 years)	Germany	Outpatient	N	N	High
Katz 2014	PE Holographic reprocessing	Anxiety Depression PTSD Self	BSI-18 BSI-18 PCL PTCI	Military trauma	USA	Outpatient	N	N	High
Korte 2017	TF-CBT	Depression PTSD Substance	BDI-II PCL-M MINI/ TLFB	Military trauma	USA	Outpatient	N	Y	High
Kubany 2003	CT	Depression PTSD Self	BDI CAPS Rosenberg SES	Multiple interpersonal trauma	USA	Outpatient	N	N	Moderate
McDonagh 2005	PE Present-centered	Anxiety Depression Dissociation PTSD Self	STAI BDI DES CAPS TSI	Child abuse (under the age 16)	USA	Outpatient	N	Y	High
Neuner 2010	NET	Depression PTSD	HSCL-25 PDS	Organized violence (conflict/torture/detention/persecution)	Germany	Outpatient	N	N	Low
Nixon 2016	CPT	Depression PTSD Self	BDI-II CAPS PTCI	Multiple interpersonal trauma	Australia	Outpatient	N	Y	Low
Pabst 2014	NET	BPD Depression Dissociation PTSD	BSL-23 HAM-D FDS PDS	Multiple interpersonal trauma	Germany	Outpatient	N	Y	High
Paivio 2010	Emotion-focused (Imaginal Confrontation)	Anxiety Depression Interpersonal = problem PTSD Self	STAI BDI-II IIP IES Rosenberg SES	Child abuse (under the age 18)	Canada	Outpatient	N	N	High
Pigeon 2009	IPT	Depression	HAM-D	Child abuse (onset age not clear)	USA	Outpatient	N	Y	High

(Continued to the next page)

Appendix II. Continued

Study	Main intervention	Outcome	Instrument	Trauma type	Study location	Setting	Phase-oriented	Intent-to-treat	Risk of bias
Resick 2002	CPT PE	Depression PTSD	BDI CAPS	Multiple interpersonal trauma	USA	Outpatient	N	Y	Moderate
Resick 2008	CPT CPT without WA written accounts	Depression PTSD Depression	BDI PDS BDI	Multiple interpersonal trauma	USA	Outpatient	N	Y	Low
Scheck 1998	EMDR	Anxiety Depression PTSD Self	STAI BDI IES TSCS	Child abuse (onset age not clear)	USA	Outpatient	N	N	High
Schnurr 2007	PE	Anxiety Depression PTSD	STAI BDI CAPS	Military trauma	USA	Outpatient	N	Y	Low
Stenmark 2013	NET	PTSD Depression	CAPS HAM-D	Organized violence (conflict/torture/detention/persecution)	Germany	Outpatient	N	N	Low
Sullivan 1999	Advocacy	Depression	CES-D	Multiple interpersonal trauma	USA	Shelter	N	N	Moderate
Suris 2013	CPT	Depression PTSD	QIDS CAPS	Military trauma	USA	Outpatient	N	N	Low
Taft 2011	Mentoring	Depression	EPDS	Multiple interpersonal trauma	Australia	Outpatient	N	N	High
Talbot 2011	IPT	Depression PTSD	HAM-D PSS	Child abuse (under the age 18)	USA	Outpatient	N	N	High
ter Heide 2011	EMDR	Anxiety Depression PTSD	HSCL-25 HSCL-25 HTQ	Organized violence (conflict/torture/detention/persecution)	Germany	Outpatient	N	N	Low
Tiwari 2010	Advocacy	Depression	BDI-II	Multiple interpersonal trauma	China	Outpatient	N	Y	Low
Vitriol 2009	Psychodynamic	Depression PTSD Interpersonal = problem	HAM-D PTO8 Lambert's-OQ-IR	Child abuse (under the age of 15)	Chile	Outpatient	N	Y	High
Weiss 2015	Common elements treatment approach (CETA) CPT	Anxiety Depression PTSD	HSCL-25 HSCL-25 HTQ	Organized violence (conflict/torture/detention/persecution)	Iraq	Outpatient	N	Y	Low

Note. BA = behavioral activation treatment; BDI = Beck Depression Inventory; BDI-II = Beck Depression Inventory-II; BSI-18 = Brief Symptom Inventory 18; BSL-23 = The short version of the Borderline Symptom List; CAPS = Clinician-Administered PTSD Scale; CAPS-SX = the 17-item Clinician-Administered PTSD Scale; CBT = cognitive behavioral therapy; CPT = cognitive processing therapy; CES-D = Center for Epidemiologic Studies-Depression Scale; CIDI = Composite International Diagnostic Interview; DBT-PTSD = dialectical behavior therapy for posttraumatic stress disorder; DES = dissociative experiences scale; DES-II = dissociative experiences scale - II; DES-T = dissociative experiences scale scale taxon; DISS = the 14-item Dissociation Scale; EMDR = eye movement desensitization and reprocessing; EPDS = Edinburgh Postnatal Depression Scale; FDS = Fragebogen für dissoziative symptom; HAM-D = Hamilton Depression Rating Scale; HRSA = Hamilton Rating Scale for Anxiety; HRSD = Hamilton Rating Scale for Depression; HSCL-25 = Hopkins Symptom Checklist-25; HTQ = Harvard Trauma Questionnaire; IES = Impact of Event Scale; IIP = Inventory of Interpersonal Problems; IPT = interpersonal psychotherapy; Lambert's-OQ-IR = Lambert's Outcome Questionnaire(OQ-45.2); MINI = Mini International Neuropsychiatric Interview; TLFB = Timeline Follow-Back; NET = narrative exposure therapy; NMR = Negative Mood Regulation Scale; PCL-C = PTSD Checklist - Civilian Version; PCL-5 = PTSD Checklist for DSM-5; PCL-M = PTSD Checklist - Military Version; PDS = Posttraumatic Diagnostic Scale; PE = prolonged exposure; PTSD = posttraumatic stress disorder; PSS = PTSD Symptom Scale; PSS-I = PTSD Symptom Scale - Interview; PTCI = Posttraumatic Cognitions Inventory; PTO8 = Post-traumatic Stress Treatment Outcome scale; QIDS = Quick Inventory of Depressive Symptomatology; Rosenberg SES = Rosenberg Self-Esteem Scale; SASII = Suicide Attempt Self Injury Interview; STAI = State-Trait Anxiety Inventory; STAI-S = State subscale of the State-Trait Anxiety Inventory; STAIR-PE = skills training in affect and interpersonal regulation- prolonged exposure; TSCS = Tennessee Self-Concept Scale; TSI = Trauma Symptom Inventory.

Appendix III.

References for the studies included in the meta-analysis

- Adenauer, H., Catani, C., Gola, H., Keil, J., Ruf, M., Schauer, M., & Neuner, F. (2011). Narrative exposure therapy for PTSD increases top-down processing of aversive stimuli-evidence from a randomized controlled treatment trial. *BMC Neuroscience*, 12, 127.
- Bichescu, D., Neuner, F., Schauer, M., & Elbert, T. (2007). Narrative exposure therapy for political imprisonment-related chronic post-traumatic stress disorder and depression. *Behaviour Research and Therapy*, 45, 2212-2220.
- Bohus, M., Dyer, A. S., Priebe, K., Krüger, A., Kleindienst, N., Schmahl, C., . . . Steil, R. (2013). Dialectical behaviour therapy for post-traumatic stress disorder after childhood sexual abuse in patients with and without borderline personality disorder: a randomised controlled trial. *Psychotherapy and Psychosomatics*, 82, 221-233.
- Bolton, P., Bass, J. K., Zangana, G. A. S., Kamal, T., Murray, S. M., Kaysen, D., . . . Rosenblum, M. (2014a). A randomized controlled trial of mental health interventions for survivors of systematic violence in Kurdistan, Northern Iraq. *BMC Psychiatry*, 14, 360.
- Bolton, P., Lee, C., Haroz, E. E., Murray, L., Dorsey, S., Robinson, C., . . . Bass, J. (2014b). A transdiagnostic community-based mental health treatment for comorbid disorders: development and outcomes of a randomized controlled trial among Burmese refugees in Thailand. *PLOS Medicine*, 11, e1001757.
- Chard, K. M. (2005). An evaluation of cognitive processing therapy for the treatment of posttraumatic stress disorder related to childhood sexual abuse. *Journal of Consulting and Clinical Psychology*, 73, 965-971.
- Cloitre, M., Koenen, K. C., Cohen, L. R., & Han, H. (2002). Skills training in affective and interpersonal regulation followed by exposure: a phase-based treatment for PTSD related to childhood abuse. *Journal of Consulting and Clinical Psychology*, 70, 1067-1074.
- Cloitre, M., Stovall-McClough, K. C., Noonan, K., Zorbas, P., Cherry, S., Jackson, C. L., . . . Petkova, E. (2010). Treatment for PTSD related to childhood abuse: A randomized controlled trial. *American Journal of Psychiatry*, 167, 915-924.
- Duffy, M., Gillespie, K., & Clark, D. M. (2007). Post-traumatic stress disorder in the context of terrorism and other civil conflict in Northern Ireland: randomised controlled trial. *British Medical Journal*, 334, 1147-1150.
- Edmond, T., Rubin, A., & Wambach, K. G. (1999). The effectiveness of EMDR with adult female survivors of childhood sexual abuse. *Social Work Research*, 23, 103-116.
- Foa, E. B., Hembree, E. A., Cahill, S. P., Rauch, S. A., Riggs, D. S., Feeny, N. C., & Yadin, E. (2005). Randomized trial of prolonged exposure for posttraumatic stress disorder with and without cognitive restructuring: outcome at academic and community clinics. *Journal of Consulting and Clinical Psychology*, 73, 953-964.
- Galovski, T. E., Blain, L. M., Mott, J. M., Elwood, L., & Houle, T. (2012). Manualized therapy for PTSD: Flexing the structure of cognitive processing therapy. *Journal of Consulting and Clinical Psychology*, 80, 968-981.
- Ghafoori, B., Fisher, D., Korosteleva, O., & Hong, M. (2016). A randomized, controlled, pilot study of a single session psychoeducation treatment for urban, culturally diverse, trauma-exposed adults. *The Journal of Nervous and Mental Disease*, 204, 421-430.
- Ghafoori, B., Hansen, M. C., Garibay, E., & Korosteleva, O. (2017). Feasibility of training frontline therapists in prolonged exposure: a randomized controlled pilot study of treatment of complex trauma in diverse victims of crime and violence. *The Journal of Nervous and Mental Disease*, 205, 283-293.
- Harkness, K. L., Bagby, R. M., & Kennedy, S. H. (2012). Childhood maltreatment and differential treatment response and recurrence in adult major depressive disorder. *Journal of Consulting and Clinical Psychology*, 80, 342-353.
- Harned, M. S., Korslund, K. E., & Linehan, M. M. (2014). A pilot randomized controlled trial of Dialectical Behavior Therapy with and without the Dialectical Behavior Therapy Prolonged Exposure protocol for suicidal and self-injuring women with borderline personality disorder and PTSD. *Behaviour Research and Therapy*, 55, 7-17.
- Hensel-Dittmann, D., Schauer, M., Ruf, M., Catani, C., Odenwald, M., Elbert, T., & Neuner, F. (2011). Treatment of traumatized victims of war and torture: a randomized controlled comparison of narrative exposure therapy and stress inoculation training. *Psychotherapy and Psychosomatics*, 80, 345-352.
- Hijazi, A. M., Lumley, M. A., Ziadni, M. S., Haddad, L., Rapport, L. J., & Arnetz, B. B. (2014). Brief narrative exposure therapy for posttraumatic stress in Iraqi refugees: A preliminary randomized clinical trial. *Journal of Traumatic Stress*, 27, 314-322.
- Johnson, D. M., Zlotnick, C., & Perez, S. (2011). Cognitive behavioral treatment of PTSD in residents of battered women's shelters: Results of a randomized clinical trial. *Journal of Consulting and Clinical Psychology*, 79, 542-551.
- Jung, K., & Steil, R. (2013). A randomized controlled trial on cognitive restructuring and imagery modification to reduce the feeling of being contaminated in adult survivors of childhood sexual abuse suffering from posttraumatic stress disorder. *Psychotherapy and Psychosomatics*, 82, 213-220.
- Katz, L. S., Douglas, S., Zaleski, K., Williams, J., Huffman, C., & Cojucar, G. (2014). Comparing holographic reprocessing and prolonged exposure for women veterans with sexual trauma: A pilot randomized trial. *Journal of Contemporary Psychotherapy*,

- 44, 9-19.
- Korte, K., Bountress, K., Tomko, R., Killeen, T., Maria, M. S., & Back, S. (2017). Integrated treatment of PTSD and substance use disorders: the mediating role of PTSD improvement in the reduction of depression. *Journal of Clinical Medicine*, 6, 9.
- Kubany, E. S., Hill, E. E., & Owens, J. A. (2003). Cognitive trauma therapy for battered women with PTSD: Preliminary findings. *Journal of Traumatic Stress*, 16, 81-91.
- McDonagh, A., Friedman, M., McHugo, G., Ford, J., Sengupta, A., Mueser, K., . . . Descamps, M. (2005). Randomized trial of cognitive-behavioral therapy for chronic posttraumatic stress disorder in adult female survivors of childhood sexual abuse. *Journal of Consulting and Clinical Psychology*, 73, 515-524.
- Neuner, F., Kurreck, S., Ruf, M., Odenwald, M., Elbert, T., & Schauer, M. (2010). Can asylum-seekers with posttraumatic stress disorder be successfully treated? A randomized controlled pilot study. *Cognitive Behaviour Therapy*, 39, 81-91.
- Nixon, R. D., Best, T., Wilksch, S. R., Angelakis, S., Beatty, L. J., & Weber, N. (2016). Cognitive processing therapy for the treatment of acute stress disorder following sexual assault: a randomised effectiveness study. *Behaviour Change*, 33, 232-250.
- Pabst, A., Schauer, M., Bernhardt, K., Ruf-Leuschner, M., Goder, R., Elbert, T., . . . Seeck-Hirschner, M. (2014). Evaluation of narrative exposure therapy (NET) for borderline personality disorder with comorbid posttraumatic stress disorder. *Clinical Neuropsychiatry*, 11, 108-117.
- Paivio, S. C., Jarry, J. L., Chagigiorgis, H., Hall, I., & Ralston, M. (2010). Efficacy of two versions of emotion-focused therapy for resolving child abuse trauma. *Psychotherapy Research*, 20, 353-366.
- Pigeon, W. R., May, P. E., Perlis, M. L., Ward, E. A., Lu, N., & Talbot, N. L. (2009). The effect of interpersonal psychotherapy for depression on insomnia symptoms in a cohort of women with sexual abuse histories. *Journal of Traumatic Stress*, 22, 634-638.
- Resick, P. A., Galovski, T. E., Uhlmansiek, M. O. B., Scher, C. D., Clum, G. A., & Young Xu, Y. (2008). A randomized clinical trial to dismantle components of cognitive processing therapy for posttraumatic stress disorder in female victims of interpersonal violence. *Journal of Consulting and Clinical Psychology*, 76, 243-258.
- Resick, P. A., Nishith, P., Weaver, T. L., Astin, M. C., & Feuer, C. A. (2002). A comparison of cognitive-processing therapy with prolonged exposure and a waiting condition for the treatment of chronic posttraumatic stress disorder in female rape victims. *Journal of Consulting and Clinical Psychology*, 70, 867-879.
- Scheck, M. M., Schaeffer, J. A., & Gillette, C. (1998). Brief psychological intervention with traumatized young women: The efficacy of eye movement desensitization and reprocessing. *Journal of Traumatic Stress*, 11, 25-44.
- Schnurr, P. P., Friedman, M. J., Engel, C. C., Foa, E. B., Shea, M. T., Chow, B. K., . . . Bernardy, N. (2007). Cognitive behavioral therapy for posttraumatic stress disorder in women: A randomized controlled trial. *JAMA*, 297, 820-830.
- Stenmark, H., Catani, C., Neuner, F., Elbert, T., & Holen, A. (2013). Treating PTSD in refugees and asylum seekers within the general health care system. A randomized controlled multicenter study. *Behaviour Research and Therapy*, 51, 641-647.
- Sullivan, C. M., & Bybee, D. I. (1999). Reducing violence using community-based advocacy for women with abusive partners. *Journal of Consulting and Clinical Psychology*, 67, 43-53.
- Suris, A., Link-Malcolm, J., Chard, K., Ahn, C., & North, C. (2013). A randomized clinical trial of cognitive processing therapy for veterans with PTSD related to military sexual trauma. *Journal of Traumatic Stress*, 26, 28-37.
- Taft, A. J., Small, R., Hegarty, K. L., Watson, L. F., Gold, L., & Lumley, J. A. (2011). Mothers' AdvocateS In the Community (MOSAIC)-non-professional mentor support to reduce intimate partner violence and depression in mothers: a cluster randomised trial in primary care. *BMC Public Health*, 11, 178.
- Talbot, N. L., Chaudron, L. H., Ward, E. A., Duberstein, P. R., Conwell, Y., O'Hara, M. W., . . . Stuart, S. (2011). A randomized effectiveness trial of interpersonal psychotherapy for depressed women with sexual abuse histories. *Psychiatric Services*, 62, 374-380.
- ter Heide, F. J. J., Mooren, T., Kleijn, W., de Jongh, A., & Kleber, R. (2011). EMDR versus stabilisation in traumatised asylum seekers and refugees: Results of a pilot study. *European Journal of Psychotraumatology*, 2, 5881.
- Tiwari, A., Fong, D. Y. T., Yuen, K. H., Yuk, H., Pang, P., Humphreys, J., & Bullock, L. (2010). Effect of an advocacy intervention on mental health in Chinese women survivors of intimate partner violence: a randomized controlled trial. *JAMA*, 304, 536-543.
- Vitriol, V. G., Ballesteros, S. T., Florenzano, R. U., Weil, K. P., & Benadof, D. F. (2009). Evaluation of an outpatient intervention for women with severe depression and a history of childhood trauma. *Psychiatric Services*, 60, 936-942.
- Weiss, W. M., Murray, L. K., Zangana, G. A. S., Mahmooth, Z., Kay-sen, D., Dorsey, S., . . . Bolton, P. (2015). Community-based mental health treatments for survivors of torture and militant attacks in Southern Iraq: a randomized control trial. *BMC Psychiatry*, 15, 249.

Supplementary Table 1. *Effect Size and Heterogeneity Statistics of Each Comparison on PTSD and Depression*

Treatment	Study name	Comparison	Outcome		Hedges's g		95% CI		Heterogeneity			Heterogeneity		
			PTSD	Depression	PTSD	Depression	Lower limit	Upper limit	Q-value	$df(Q)$	I^2	Q-value	$df(Q)$	I^2
CPT/CT	Bolton 2014a	CPT vs. WL	0.511	0.197	0.824	71.999	0.800	88.889	0.436	0.429	0.116	0.741	86.504	90.752
	Chard 2005	CPT vs. WL	2.687	1.963	3.410					2.317	1.639	2.994		
	Galovski 2012	CPT vs. WL	1.290	0.861	1.719					0.993	0.580	1.407		
	Jung 2013	Cognitive modification+ imagery vs. WL	0.857	0.103	1.610					0.421	-0.306	1.149		
	Kubany 2003	CT vs. WL	3.039	2.030	4.048					2.539	1.617	3.461		
NET	Nixon 2016	CPT vs. TAU	0.078	-0.491	0.647					0.421	-0.154	0.996		
	Resick 2002	CPT vs. WL	1.296	0.882	1.710					0.871	0.476	1.266		
	Suris 2013	CPT vs. present-centered	0.991	0.538	1.444					2.613	2.034	3.193		
	Weiss 2015	CPT vs. WL	0.409	0.116	0.702					0.399	0.106	0.691		
	Random effects		1.155	0.684	1.625					1.172	0.657	1.688		
PE	Adenauer 2011	NET vs. WL	2.939	1.662	4.216	22.568	7.000	68.983	0.352	1.583	0.578	2.588	6.142	7.000
	Bichescu 2007	NET vs. psychoeducation	1.667	0.632	2.701					0.900	-0.028	1.827		
	Hensel-Dittmann 2011	NET vs. stress inoculation	1.045	0.274	1.817					0.653	-0.099	1.406		
	Hijazi 2014	Brief NET vs. WL	0.230	-0.283	0.743					0.231	-0.282	0.744		
	Neuner 2010	NET vs. TAU	1.276	0.531	2.020					0.568	-0.122	1.258		
PE	Pabst 2014	NET vs. TBE	0.205	-0.601	1.011					0.609	-0.215	1.432		
	Stenmark 2013 (refugee)	NET vs. TAU	0.752	0.068	1.436					0.508	-0.163	1.179		
	Stenmark 2013 (asylumseeker)	NET vs. TAU	0.554	-0.354	1.462					0.564	-0.345	1.472		
	Random effects		0.969	0.464	1.474					0.580	0.320	0.841		
	Foa 2005	PE vs. WL	0.811	0.358	1.264	15.666	6.000	61.700	0.077	0.739	0.259	1.218	8.217	6.000
PE	Foa 2005	PE+cognitive restructuring vs. WL	0.715	0.261	1.169					0.614	0.133	1.096		
	Ghafoori 2017	PE vs. person-centered	0.502	0.009	0.995					0.701	0.201	1.201		
	Katz 2014	PE vs. person centered	0.741	-0.111	1.593					0.571	-0.269	1.411		
	McDonagh 2005	PE+cognitive restructuring vs. WL	0.455	-0.091	1.002					0.375	-0.169	0.919		
	Resick 2002	PE vs. WL	1.134	0.729	1.539					0.720	0.330	1.109		
Random effects	Schnurr 2007	PE vs. present-centered	0.269	0.036	0.502					0.229	-0.003	0.462		
	Random effects		0.643	0.371	0.915					0.514	0.319	0.708		

(Continued to the next page)

Supplementary Table 1. Continued

Treatment	Study name	Comparison	Outcome		Hedges's g	95% CI		Heterogeneity			Outcome		Hedges's g	95% CI		Heterogeneity		
			PTSD			Lower limit	Upper limit	Q- value	I ²	df(Q)	Tau ²	Depression		Lower limit	Upper limit	Q- value	I ²	df(Q)
Phased TF-CBT	Bohus 2013	DBT-PTSD vs. TAU			1.343	0.843	1.843	0.050	1.000	0.000	0.000	0.598	0.137	1.059	3.789	1.000	73.609	0.223
	Cloitre 2002	STAIR-PE vs. WL			1.436	0.796	2.076					1.377	0.743	2.011				
	Random effects				1.378	0.984	1.772					0.956	0.195	1.717				
Single TF-CBT	Bolton 2014b	common elements treatment approach vs. WL			1.134	1.360	9.811	35.115	3.000	91.457	0.429				20.101	2.000	90.050	0.497
EMDR	Duffy 2007	CBT vs. WL			0.825	1.355	3.055					0.751	0.225	1.277				
	Korte 2017	Exposure (COPE) vs. CBT for substance			0.704	1.174	2.933					0.515	0.050	0.979				
	Weiss 2015	Common elements treatment approach vs. WL			2.388	2.821	10.795					1.811	1.415	2.207				
	Random effects				1.268	1.944	3.674					1.035	0.194	1.876				
	Edmond 1999	EMDR vs. TAU			0.250	-0.360	0.860	4.239	3.000	29.236	0.050	0.089	-0.519	0.696	1.327	3.000	0.000	0.000
Non-TF CBT	Edmond 1999	EMDR vs. WL			1.065	0.406	1.724					0.382	-0.239	1.003				
	Scheck 1998	EMDR vs. TAU			0.997	0.453	1.541					0.526	0.017	1.034				
	ter Heide 2011	EMDR vs. stabilization			0.718	-0.445	1.881					0.592	-0.557	1.741				
	Random effects				0.767	0.361	1.173					0.374	0.057	0.691				
	Bolton 2014a	behavior activation vs. WL			0.284	-0.020	0.587	0.636	1.000	0.000	0.000	0.321	0.017	0.624	1.418	1.000	29.466	0.017
Present- centered	Johnson 2011	Stabilization (HOPE) vs. usual case management			0.058	-0.405	0.522					0.664	0.187	1.140				
	Random effects				0.216	-0.038	0.470					0.441	0.120	0.762				
	Katz 2014	holographic reprocessing vs. person centered			1.128	0.325	1.931	7.510	4.000	46.736	0.077	0.942	0.156	1.728	2.929	5.000	0.000	0.000
	McDonagh 2005	present-centered vs. WL			0.728	0.134	1.321					0.449	-0.133	1.030				
	Paivio 2010	Emotion-focused (imagi- nal confrontation) vs. EF w/o imaginal confrontation			0.455	-0.204	1.113					0.129	-0.521	0.780				
Random effects	Pigeon 2009	IPT vs. TAU			0.749	0.192	1.306					0.381	-0.087	0.850				
	Talbot 2011	IPT vs. TAU			0.085	-0.332	0.502					0.335	-0.207	0.877				
	Vitriol 2009	psychodynamic vs. TAU			0.562	0.203	0.921					0.555	0.131	0.980				
	Random effects											0.446	0.225	0.667				

Note. COPE = Concurrent Treatment of PTSD and Substance Use Disorders Using Prolonged Exposure; CPT = Cognitive Processing Therapy; CT = Cognitive Therapy; DBT-PTSD = Dialectical Behavior Therapy for Posttraumatic Stress Disorder; EF = emotion-focused; HOPE = Helping to Overcome PTSD through Empowerment; IPT = Interpersonal Psychotherapy; NET = Narrative Exposure Therapy; PE = Prolonged Exposure; PTSD = Posttraumatic Stress Disorder; STAIR-PE = skills training in affect and interpersonal regulation and prolonged exposure; TAU = treatment as usual; TBE = Treatment by Experts for Borderline Personality Disorder; TF-CBT = Trauma-Focused Cognitive Behavioral Therapy; WL = wait list.

Supplementary Table 2. *Effect Size and Heterogeneity Statistics of Each Comparison on Anxiety and Dissociation*

Treatment	Study name	Comparison	Outcome			Hedges's g			95% CI			Heterogeneity			Heterogeneity		
			Anxiety	Dissociation	Outcome	g	Lower limit	Upper limit	Lower limit	Upper limit	Q-value	$df(Q)$	I^2	Tau^2	Q-value	$df(Q)$	I^2
CPT/CT	Bolton 2014a	CPT vs. WL	0.525	0.211	0.839	1.375	1.000	27.282	0.009								
	Chard 2005	CPT vs. WL															
	Galowski 2012	CPT vs. WL															
	Jung 2013	Cognitive modification+ imagery vs. WL															
	Kubany 2003	CT vs. WL															
	Nixon 2016	CPT vs. TAU															
	Resick 2002	CPT vs. WL															
	Suris 2013	CPT vs. present-centered															
	Weiss 2015	CPT vs. WL	0.269	-0.023	0.561												
	Random effects		0.390	0.140	0.641												
NET	Adenauer 2011	NET vs. WL															
	Bichescu 2007	NET vs. psychoeducation															
	Hensel-Dittmann 2011	NET vs. stress inoculation															
	Hijazi 2014	Brief NET vs. WL															
	Neuner 2010	NET vs. TAU															
	Pabst 2014	NET vs. TBE															
	Stenmark 2013 (refugee)	NET vs. TAU															
	Stenmark 2013 (asylumseeker)	NET vs. TAU															
	Random effects																
			0.181	-0.625	0.986												

(Continued to the next page)

Supplementary Table 2. Continued

Treatment	Study name	Comparison	Outcome		Hedges's <i>g</i>	95% CI		Heterogeneity			Outcome			Hedges's <i>g</i>			95% CI			Heterogeneity		
			Anxiety			Lower limit	Upper limit	Q-value	<i>df</i> (Q)	<i>I</i> ²	<i>Tau</i> ²	Dissociation					Lower limit	Upper limit	Q-value	<i>df</i> (Q)	<i>I</i> ²	<i>Tau</i> ²
PE	Foa 2005	PE vs. WL						2.594	3.000	0.000	0.000											
	Foa 2005	PE+cognitive restructuring vs. WL																				
	Ghafoori 2017	PE vs. person-centered			0.287	-0.202	0.775															
	Katz 2014	PE vs. person centered			0.864	0.001	1.726															
Phased TF-CBT	McDonagh 2005	PE + cognitive restructuring vs. WL			0.365	-0.179	0.908															
	Resick 2002	PE vs. WL																				
	Schnurr 2007	PE vs. present-centered			0.170	-0.063	0.402															
	Random effects				0.245	0.055	0.436															
Single TF-CBT	Bohus 2013	DBT-PTSD vs. TAU												0.486	0.028	0.944			1.936	1.000	48.350	0.070
	Cloitre 2002	STAIR-PE vs. WL												1.025	0.419	1.631						
	Random effects													0.718	0.195	1.241						
	Bolton 2014b	Common elements treatment approach vs. WL			0.750	0.968	6.759	13.998	1.000	92.856	0.329											
EMDR	Duffy 2007	CBT vs. WL																				
	Korte 2017	Exposure (COPE) vs. CBT for substance																				
	Weiss 2015	Common elements treatment approach vs. WL			1.592	1.975	8.134															
	Random effects				1.156	1.980	2.748															
EMDR	Edmond 1999	EMDR vs. TAU			0.382	-0.231	0.995	5.615	3.000	46.569	0.108											
	Edmond 1999	EMDR vs. WL			1.369	0.683	2.056															
	Scheck 1998	EMDR vs. TAU			0.622	0.106	1.138															
	ter Heide 2011	EMDR vs. stabilization			1.359	0.091	2.628															
EMDR	Random effects				0.830	0.351	1.309															

(Continued to the next page)

Supplementary Table 2. *Continued*

Treatment	Study name	Comparison	Outcome	Hedges's <i>g</i>	95% CI		Heterogeneity			Outcome	Hedges's <i>g</i>	95% CI		Heterogeneity				
			Anxiety	<i>g</i>	Lower limit	Upper limit	Q-	<i>df</i> (Q)	<i>I</i> ²	Tau ²	Dissociation	<i>g</i>	Lower limit	Upper limit	Q-	<i>df</i> (Q)	<i>I</i> ²	Tau ²
Non-TF CBT	Bolton 2014a	Behavior activation vs. WL		0.513	0.207	0.820												
	Johnson 2011	Stabilization (HOPE) vs. usual case management																
Present-centered	Random effects																	
	Katz 2014	Holographic reprocessing vs. person centered		1.091	0.292	1.890	2.188	2.000	8.609	0.011								
	McDonagh 2005	present-centered vs. WL		0.475	-0.107	1.058					1.081	0.465	1.698					
	Paivio 2010	Emotion-focused (imaginal confrontation) vs. EF w/o imaginal confrontation		0.345	-0.310	1.000												
	Pigeon 2009	IPT vs. TAU																
	Talbot 2011	IPT vs. TAU																
	Vitriol 2009	Psychodynamic vs. TAU																
	Random effects																	
				0.576	0.175	0.978												

Note. COPE = Concurrent Treatment of PTSD and Substance Use Disorders Using Prolonged Exposure; CPT = Cognitive Processing Therapy; CT = Cognitive Therapy; DBT-PTSD = Dialectical Behavior Therapy for Posttraumatic Stress Disorder; EF = emotion-focused; HOPE = Helping to Overcome PTSD through Empowerment; IPT = Interpersonal Psychotherapy; NET = Narrative Exposure Therapy; PE = Prolonged Exposure; PTSD = Posttraumatic Stress Disorder; STAIR-PE = skills training in affect and interpersonal regulation and prolonged exposure; TAU = treatment as usual; TBE = Treatment by Experts for Borderline Personality Disorder; TF-CBT = Trauma-Focused Cognitive Behavioral Therapy; WL = wait list.

Supplementary Table 3. Effect Size and Heterogeneity Statistics of Each Comparison on Negative self-concept and Emotion Regulation

Treatment	Study name	Comparison	Outcome		Heterogeneity			95% CI		Heterogeneity		
			Negative self-concept	Hedges's <i>g</i>	Lower limit	Upper limit	Q-value	<i>df</i> (Q)	<i>I</i> ²	Emotion regulation	Hedges's <i>g</i>	Lower limit
CPT/CT	Bolton 2014a	CPT vs. WL					13.280	2.000	84.939	0.773		
	Chard 2005	CPT vs. WL										
	Galovski 2012	CPT vs. WL										
	Junger 2013	Cognitive modification+ imagery vs. WL		0.453	-0.276	1.182						
	Kubany 2003	CT vs. WL		2.409	1.508	3.310						
	Nixon 2016	CPT vs. TAU		0.594	0.012	1.175						
	Resick 2002	CPT vs. WL										
	Suris 2013	CPT vs. present-centered										
	Weiss 2015	CPT vs. WL										
	Random effects			1.110	0.027	2.193						
	Adenauer 2011	NET vs. WL										
	Bichescu 2007	NET vs. psychoeducation										
NET	Hensel-Dittmann 2011	NET vs. stress inoculation										
	Hijazi 2014	briefNET vs. WL										
	Neuner 2010	NET vs. TAU										
	Pabst 2014	NET vs. TBE										
	Stenmark 2013 (refugee)	NET vs. TAU										
	Stenmark 2013 (asylumseeker)	NET vs. TAU										
	Random effects											
	Foa 2005	PE vs. WL					1.327	1.000	24.662	0.046		
	Foa 2005	PE+cognitive restructuring vs. WL										
	Ghafoori 2017	PE vs. person-centered										
PE	Katz 2014	PE vs. person-centered		1.050	0.169	1.931						
	McDonagh 2005	PE+cognitive restructuring vs. WL		0.441	-0.105	0.986						
	Resick 2002	PE vs. WL										
	Schnurr 2007	PE vs. present-centered										
	Random effects			0.643	0.081	1.205						

(Continued to the next page)

Supplementary Table 3. *Continued*

Treatment	Study name	Comparison	Outcome		95% CI		Heterogeneity		Outcome		95% CI		Heterogeneity	
			Negative self-concept	Hedges's <i>g</i>	Lower limit	Upper limit	Q-value	<i>I</i> ²	Emotion regulation	Hedges's <i>g</i>	Lower limit	Upper limit	Q-value	<i>I</i> ²
Phased TF-CBT	Bohus 2013	DBT-PTSD vs. TAU												
	Cloitre 2002	STAIR-PE vs. WL								1.305	0.677	1.933		
Single TF-CBT	Random effects													
	Bolton 2014b	Common elements treatment approach vs. WL												
	Duffy 2007	CBT vs. WL												
	Korte 2017	Exposure (COPE) vs. CBT for substance												
	Weiss 2015	Common elements treatment approach vs. WL												
EMDR	Random effects													
	Edmond 1999	EMDR vs. TAU												
	Edmond 1999	EMDR vs. WL												
	Scheck 1998	EMDR vs. TAU		0.457	-0.054	0.967								
	ter Heide 2011	EMDR vs. stabilization												
	Random effects													
Non-TF CBT	Bolton 2014a	Behavior activation vs. WL												
	Johnson 2011	Stabilization (HOPE) vs. usual case management												
Present-centered	Random effects													
	Katz 2014	Holographic reprocessing vs. person centered		0.990	0.200	1.780	2.531	2.000	20.985	0.031				
	McDonagh 2005	Present-centered vs. WL		0.678	0.087	1.269								
	Paivio 2010	Emotion-focused (imaginal confrontation) vs. EF w/o imaginal confrontation		0.188	-0.464	0.839								
	Pigeon 2009	IPT vs. TAU												
	Talbot 2011	IPT vs. TAU												
	Vitriol 2009	Psychodynamic vs. TAU												
	Random effects			0.588	0.154	1.021								

Note. COPE = Concurrent Treatment of PTSD and Substance Use Disorders Using Prolonged Exposure; CPT = Cognitive Processing Therapy; CT = Cognitive Therapy; DBT-PTSD = Dialectical Behavior Therapy for Posttraumatic Stress Disorder; EF = emotion-focused; HOPE = Helping to Overcome PTSD through Empowerment; IPT = Interpersonal Psychotherapy; NET = Narrative Exposure Therapy; PE = Prolonged Exposure; PTSD = Posttraumatic Stress Disorder; STAIR-PE = skills training in affect and interpersonal regulation and prolonged exposure; TAU = treatment as usual; TBE = Treatment by Experts for Borderline Personality Disorder; TF-CBT = Trauma-Focused Cognitive Behavioral Therapy; WL = wait list.


Treatment	Study name	Comparison	Outcome Interpersonal problem	Hedges's <i>g</i>	95% CI		Heterogeneity			Outcome Borderline personality	Hedges's <i>g</i>	95% CI		Heterogeneity		
					Lower limit	Upper limit	Q- value	<i>I</i> ²	<i>Tau</i> ²			Lower limit	Upper limit	Q- value	<i>I</i> ²	<i>Tau</i> ²
CPT/CT	Bolton 2014a	CPT vs. WL														
	Chard 2005	CPT vs. WL														
	Galovski 2012	CPT vs. WL														
	Jung 2013	Cognitive modification+ imagery vs. WL														
	Kubany 2003	CT vs. WL														
	Nixon 2016	CPT vs. TAU														
	Resick 2002	CPT vs. WL														
	Suris 2013	CPT vs. present-centered														
	Weiss 2015	CPT vs. WL														
	Random effects															
NET	Adenauer 2011	NET vs. WL														
	Bichescu 2007	NET vs. psychoeducation														
	Hensel-Dittmann 2011	NET vs. stress inoculation														
	Hijazi 2014	brief NET vs. WL														
	Neuner 2010	NET vs. TAU														
	Pabst 2014	NET vs. TBE														
	Stenmark 2013 (refugee)	NET vs. TAU														
	Stenmark 2013 (asylumseeker)	NET vs. TAU														
	Random effects															
	Foa 2005	PE vs. WL														
PE	Foa 2005	PE+cognitive restructuring vs. WL														
	Ghafoori 2017	PE vs. person-centered														
	Katz 2014	PE vs. person centered														
	McDonagh 2005	PE+cognitive restructuring vs. WL														
	Resick 2002	PE vs. WL														
	Schnurr 2007	PE vs. present-centered														
	Random effects															
	Bohus 2013	DBT-PTSD vs. TAU														
	Cloitre 2002	STAIR-PE vs. WL														
	Random effects															
Phased TF-CBT	Bohus 2013	DBT-PTSD vs. TAU														
	Cloitre 2002	STAIR-PE vs. WL														
Random effects																
Phased TF-CBT	Bohus 2013	DBT-PTSD vs. TAU														
	Cloitre 2002	STAIR-PE vs. WL														
Random effects																
Phased TF-CBT	Bohus 2013	DBT-PTSD vs. TAU														
	Cloitre 2002	STAIR-PE vs. WL														
Random effects																
Phased TF-CBT	Bohus 2013	DBT-PTSD vs. TAU														
	Cloitre 2002	STAIR-PE vs. WL														
Random effects																
Phased TF-CBT	Bohus 2013	DBT-PTSD vs. TAU														
	Cloitre 2002	STAIR-PE vs. WL														
Random effects																
Phased TF-CBT	Bohus 2013	DBT-PTSD vs. TAU														
	Cloitre 2002	STAIR-PE vs. WL														
Random effects																
Phased TF-CBT	Bohus 2013	DBT-PTSD vs. TAU														
	Cloitre 2002	STAIR-PE vs. WL														
Random effects																
Phased TF-CBT	Bohus 2013	DBT-PTSD vs. TAU														
	Cloitre 2002	STAIR-PE vs. WL														
Random effects																
Phased TF-CBT	Bohus 2013	DBT-PTSD vs. TAU														
	Cloitre 2002	STAIR-PE vs. WL														
Random effects																
Phased TF-CBT	Bohus 2013	DBT-PTSD vs. TAU														
	Cloitre 2002	STAIR-PE vs. WL														
Random effects																
Phased TF-CBT	Bohus 2013	DBT-PTSD vs. TAU														
	Cloitre 2002	STAIR-PE vs. WL														
Random effects																
Phased TF-CBT	Bohus 2013	DBT-PTSD vs. TAU														
	Cloitre 2002	STAIR-PE vs. WL														
Random effects																
Phased TF-CBT	Bohus 2013	DBT-PTSD vs. TAU														
	Cloitre 2002	STAIR-PE vs. WL														
Random effects																
Phased TF-CBT	Bohus 2013	DBT-PTSD vs. TAU														
	Cloitre 2002	STAIR-PE vs. WL														
Random effects																
Phased TF-CBT	Bohus 2013	DBT-PTSD vs. TAU														
	Cloitre 2002	STAIR-PE vs. WL														
Random effects																
Phased TF-CBT	Bohus 2013	DBT-PTSD vs. TAU														
	Cloitre 2002	STAIR-PE vs. WL														
Random effects																
Phased TF-CBT	Bohus 2013	DBT-PTSD vs. TAU														
	Cloitre 2002	STAIR-PE vs. WL														
Random effects																
Phased TF-CBT	Bohus 2013	DBT-PTSD vs. TAU														
	Cloitre 2002	STAIR-PE vs. WL														
Random effects													</			

(Continued to the next page)


Supplementary Table 4. *Continued*

Treatment	Study name	Comparison	Outcome		Hedges's <i>g</i>		95% CI		Heterogeneity		Outcome		Hedges's <i>g</i>		95% CI		Heterogeneity	
			Interpersonal problem	Borderline personality	<i>g</i>	<i>g</i>	Lower limit	Upper limit	Q- value	<i>I</i> ²	Q- value	Borderline personality	<i>g</i>	<i>g</i>	Lower limit	Upper limit	Q- value	<i>I</i> ²
Single TF-CBT	Bolton 2014b	Common elements treatment approach vs. WL																
	Duffy 2007	CBT vs. WL																
	Korte 2017	Exposure (COPE) vs. CBT for substance																
	Weiss 2015	Common elements treatment approach vs. WL																
EMDR	Random effects																	
	Edmond 1999	EMDR vs. TAU																
	Edmond 1999	EMDR vs. WL																
	Scheck 1998	EMDR vs. TAU																
	ter Heide 2011	EMDR vs. stabilization																
Non-TF CBT	Random effects																	
	Bolton 2014a	Behavior activation vs. WL																
	Johnson 2011	Stabilization (HOPE) vs. usual case management																
Present-centered	Random effects																	
	Katz 2014	Holographic reprocessing vs. person centered							1.402	1.000	28.675	0.031						
	McDonagh 2005	Present-centered vs. WL																
	Paivio 2010	Emotion-focused (imaginal confrontation) vs. EF w/o imaginal confrontation			0.030		-0.621	0.680										
	Pigeon 2009	IPT vs. TAU																
	Talbot 2011	IPT vs. TAU																
	Vitriol 2009	psychodynamic vs. TAU			0.498		0.075	0.921										
	Random effects				0.332		-0.108	0.771										

Note. COPE = Concurrent Treatment of PTSD and Substance Use Disorders Using Prolonged Exposure; CPT = Cognitive Processing Therapy; CT = Cognitive Therapy; DBT-PTSD = Dialectical Behavior Therapy for Posttraumatic Stress Disorder; EF = emotion-focused; HOPE = Helping to Overcome PTSD through Empowerment; IPT = Interpersonal Psychotherapy; NET = Narrative Exposure Therapy; PE = Prolonged Exposure; PTSD = Posttraumatic Stress Disorder; STAIR-PE = skills training in affect and interpersonal regulation and prolonged exposure; TAU = treatment as usual; TBE = Treatment by Experts for Borderline Personality Disorder; TF-CBT = Trauma-Focused Cognitive Behavioral Therapy; WL = wait list.




Supplementary Figure 1. Forest plot of psychological treatment vs TAU/WL comparisons on outcome of PTSD.




Meta Analysis

Supplementary Figure 2. Forest plot of psychological treatment vs TAU/WL comparisons on outcome of depression.




Meta Analysis

Supplementary Figure 3. Forest plot of psychological treatment vs TAU/WL comparisons on outcome of anxiety.



Meta Analysis

Supplementary Figure 4. Forest plot of psychological treatment vs TAU/WL comparisons on outcome of negative self.



Meta Analysis

Supplementary Figure 5. Forest plot of psychological treatment vs TAU/WL comparisons on outcome of dissociation.