

Psychometric Investigation of the Revised Children's Manifest Anxiety Scale, Second Edition, Short Form among the Korean Youth Population

Hayoung Yoon¹ Eun-Ho Lee² Soon-Taeg Hwang³ Sang-Hwang Hong⁴ Ji-Hae Kim^{1†}

¹Department of Psychiatry, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul; ²Depression Center, Department of Psychiatry, Samsung Medical Center, Seoul; ³Department of Psychology, Chungbuk National University, Cheongju;

⁴Department of Education, Chinju National University of Education, Jinju, Korea

The Revised Children's Manifest Anxiety Scale, Second Edition (RCMAS-2) has long been used to measure anxiety levels in youth. Its introduction of a 10-item short form has enriched the efficiency and applicability of the scale; however, more research is warranted to elucidate its psychometric properties. This study aims to investigate the factor structure and the measurement invariance of the RCMAS-2 short form (RCMAS-2 SF) in Korean youth. In total, 1,525 participants from seven different cities of South Korea were included in the analysis (Mean Age = 12.49, *SD* = 2.54). After randomly assigning participants into two groups, we consecutively performed exploratory and confirmatory factor analyses. The results of both analyses demonstrated that pediatric anxiety assessed using the RCMAS-2 SF is composed of two distinct latent factors: physiological anxiety and cognitive anxiety. Furthermore, the results supported strong invariance across gender and age. The RCMAS-2 SF score also showed good indices of internal consistency, test-retest reliability, and concurrent validity. These findings highlight that the scale can be widely used as a time-efficient screening tool that enables valid score comparisons between boys and girls of different ages.

Keywords: anxiety, RCMAS-2, short form, measurement invariance, reliability, validity

Introduction

Childhood and adolescence are the core risk phases for developing anxiety symptoms (Costello et al., 2004; Merikangas et al., 2010). If left untreated, such symptoms persist throughout adulthood, conferring a debilitating threat of secondary conditions such as poor academic achievement, lowered self-esteem, eating disorder, depression, and even attempted suicide (Bittner et al., 2007; Keller et al., 1992; Copeland et al., 2014; Schaumberg et al., 2019; Van Oort

et al., 2009). However, less than one third of children with anxiety symptoms utilize mental health services (Barker et al., 2019; Bienvenu & Ginsburg, 2007). With meta-analytic reviews reporting the global prevalence of pediatric anxiety disorders as high as up to 25%, it has long been a public priority to discover anxious youth in advance and make early intervention to protect them from pathogenic complications (Chisholm et al., 2016). With this regard, attempts have been made to develop sound instruments to measure the severity of childhood anxiety.

Amongst many self-report questionnaires designed to assess anxiety in youth, the Revised Children's Manifest Anxiety Scale, Second Edition (RCMAS-2; Reynolds & Richmond, 2008) is one of the most frequently used instruments (Fenley et al., 2021). It is a 49-item scale that yields the Total Anxiety score (TOT) and three following subscale scores: 1) Physiological Anxiety (PHY), 2) Worry

[†]Correspondence to Ji-Hae Kim, Department of Psychiatry, Samsung Medical Center, Sungkyunkwan University School of Medicine, 81 Irwon-ro, Gangnam-gu, Seoul, Korea; E-mail: jihae0931.kim@samsung.com

Received Dec 1, 2021; Revised Feb 9, 2022; Accepted Mar 8, 2022

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

(WOR), and 3) Social Anxiety (SOC). Several studies have supported its psychometric properties, reporting good internal consistency, test-retest reliability, and construct validity estimates in clinical and nonclinical populations (Ahmad & Mansoor, 2011; Raad, 2013; Mahakwe et al., 2021). It has been translated into diverse languages and is widely used to screen out anxious children or measure therapeutic effects in relieving anxiety (Gaesser & Karan, 2017; McEvoy et al., 2019; Zhou et al., 2019).

One of the most compelling advantages of RCMAS-2 is indeed its 10-item short form (RCMAS-2 SF; Reynolds & Richmond, 2008). The length of a self-report is of paramount importance to its usability (Ahlen et al., 2018; Ebesutani et al., 2012). Short forms provide several advantages over lengthy questionnaires because the latter causes response bias and measurement error by making a respondent more inattentive and careless (Billieux et al., 2012; Li & Lopez, 2007). Some findings also emphasize that too many items lower the completion rate, ultimately resulting in information loss (Rolstad et al., 2011). In light of these circumstances, the short form total anxiety score (SF-TOT) of the RCMAS-2 is suggested as a competent alternative to the original TOT of 49-items, and its correlation coefficient was reported as high as .90 (Reynolds & Richmond, 2008).

Despite its widespread use, research on the RCMAS-2 SF is scarce. There are only three rudimental studies that examined the psychometric properties of the scale. Once the one-factor model was initially suggested and conceptualized as reflecting general manifest anxiety by Lowe (2015a), two confirmatory factor analysis (CFA) studies examined the unidimensionality of the scale (Lowe, 2015b; Lowe & Ang, 2016). The results achieved adequate fit; however, the limited age-range and cultural background of the prior study participants highlight the necessity of an additional examination on the dimensionality underlying the RCMAS-2 SF. Notably, the one-factor model was supported exclusively among American students or English-speaking youth population; thus, it remains unclear whether a consistent factor structure would be observed in cultures where verbal expressions and social acceptance on anxiety symptoms are different.

Culture is indeed a potential candidate that can cause factorial invariance of pediatric anxiety across studies. Few studies have documented such cultural disparities (Chan & Leung, 2015; Vare-

la & Biggs, 2006). For example, data on collectivism-oriented Latin American children showed a different factor structure for the RCMAS than their individualistic European American counterparts. Likewise, a study on Hong Kong adolescents whose culture exerts a high amount of restraint over one's expression of anxiety failed to replicate the original five-factor structure of the Screen for Child Anxiety Related Emotional Disorders (SCARED). The authors argue that such a gap may stem from different social norms and parenting attitudes that wield a strong influence on children's development and expression of anxiety (Pina et al., 2009; Weisz et al., 1987; Valera & Hensley-Maloney, 2009). More specifically, collectivistic cultures that emphasize self-restraint may lead children to express their anxiety in forms of internalizing problem (e.g., shyness, worry) whereas externalizing symptoms (e.g., crying out, anger) may be more linked to individualistic cultures (Weisz et al., 1987; Weitkamp & Seiffge-Krenke, 2019). Another potential suggestion is that children in collectivistic culture may exhibit their anxiety as a somatic symptom because the culture and controlling parents place more importance on the suppression of psychological problems and community harmony than individualistic cultures do (Kim et al., 2019).

Furthermore, a wealth of studies has yielded strong supports for the multi-dimensionality of childhood anxiety. Unlike studies that uniformly propose a single factor underlying the RCMAS-2 SF, no study on the original 49-item RCMAS-2 has claimed to have one factor; instead, researchers have generally suggested a five-factor model with three distinct anxiety subscales and two validity subscales. Studies conducted in Asian countries, in particular, seem to converge into a four-factor model that incorporates two Defensiveness factors, the PHY factor, and one other factor that integrates both the WOR and SOC items (Cha et al., 2020; Raad, 2013; Wu et al., 2016; Zhu & Lowe, 2018). In addition, researches on other questionnaires on childhood anxiety, such as Spence Children's Anxiety Scale (SCAS) and the Youth Anxiety Measure for DSM-5 (YAM-5) have repeatedly supported multidimensional factor structures that reflect complex facets of childhood anxiety (Hale et al., 2011; Simon et al., 2017; Spence et al., 2003). Therefore, it is necessary to examine alternative models other than the one-factor model for the RCMAS-2 SF to reconfirm whether the scale is composed of a single factor that is less supported by the literature.

Meanwhile, the RCMAS-2 SF, is commonly used to compare anxiety levels among youth. Yet, to ensure whether the differences in summary statistics denote true gender or age effects, measurement invariance (MI) must be tested prior to the utilization of the scale (Cheung & Rensvold, 2002; Meredith, 1993; Van de Schoot et al., 2012). Since the response tendency can vary systematically depending on how different groups interpret each item, the confirmation of a unified interpretation by respondents is important. Hence the establishment of MI in RCMAS-2 SF is also needed to argue for age and gender differences in the youth.

Building on the recommendations and limitations aforementioned in the literature, the current study aimed to elucidate the factors underlying pediatric anxiety in Korean community-dwelling sample using the RCMAS-2 SF and to examine if any factors contribute to the measurement invariance of the instrument. Specifically, we firstly examined the factor structure of the RCMAS-2 SF using exploratory factor analysis (EFA) and CFA. Considering the lack of prior studies, we not only examined one factor but also inspected the validity of two and three factor structures. We then investigated the reliability and validity of the scale by calculating Cronbach's alpha, the test-retest reliability index, and correlation coefficients with other measures of childhood psychopathology. Furthermore, we examined the measurement invariance of the RCMAS-2 SF across gender and age.

Method

Participants

The study protocol was approved by the Institutional Board of

Chungbuk National University (CBNU-201708-SB-503-01). The process of translation and validation into Korean has been well documented by Cha et al. (2020). A community sample of 1,565 youth aged 8 to 18 years was recruited from seven different cities in South Korea, hence enabling a nationwide study. All the participants provided written consent after the purpose and ethical considerations of the study were addressed. Among 1,565 students, ten cases with missing demographic data were firstly excluded. Questionnaires were considered incomplete if 1) more than five items ($>10\%$ of total 49 items) in RCMAS-2 or 2) any of the RCMAS-2 SF items were left unanswered. All the incomplete responses were excluded from further analysis. As a result, a total of 1,525 participants ($M=12.49$, $SD=2.54$) with 651 boys (42.7%) and 874 girls (57.3%) were included in the final sample. These participants were then randomly assigned into either the EFA or CFA group. No significant differences were observed in any of the demographic variables between the groups. The statistics are presented in Table 1.

Furthermore, a subset of 191 participants ($M=11.27$, $SD=1.57$) was analyzed in a test-retest analysis because they completed a second administration of the RCMAS-2 two weeks after the first administration. Furthermore, a total of 799 students with a mean age of 10.89 ($SD=1.32$) also completed Spielberger's State-Trait Anxiety Inventory for Children (STAIC) and Children's Depression Inventory 2 (CDI 2) for the test of concurrent validity.

Measures

Revised Children's Manifest Anxiety Scale, Second Edition (RCMAS-2)

The RCMAS-2 is a self-report inventory that assesses anxiety lev-

Table 1. Demographic Characteristics of Participants

| | Total (N= 1,525) | Group1_EFA (n= 785) | Group2_CFA (n= 740) | t/χ^2 | p |
|--|------------------|---------------------|---------------------|------------|------|
| Age, mean (SD) | 12.49 (2.54) | 12.49 (2.51) | 12.50 (2.56) | .062 | .950 |
| Gender, No. (% of Female) | 874 (57.31) | 452 (57.60) | 422 (57.02) | -.218 | .828 |
| School, No. (% of Elementary students) | 929 (60.92) | 477 (60.76) | 452 (61.08) | -.127 | .899 |
| PHY, mean (SD) | 3.21 (2.59) | 3.24 (2.63) | 3.19 (2.54) | -.379 | .704 |
| WOR, mean (SD) | 6.01 (4.34) | 5.99 (4.40) | 6.03 (4.28) | .186 | .852 |
| SOC, mean (SD) | 2.74 (2.82) | 2.77 (2.91) | 2.71 (2.71) | -.397 | .692 |
| TOT, mean (SD) | 11.97 (8.69) | 12.00 (8.93) | 11.93 (8.44) | -.149 | .882 |
| SF-TOT, mean (SD) | 3.24 (2.70) | 3.29 (2.75) | 3.18 (2.64) | -.773 | .440 |

Note. PHY = Physiological Anxiety score; WOR = Worry score; SOC = Social Anxiety score; TOT = Total Anxiety score; SF-TOT = Short Form Total Anxiety score.

* $p < .05$.

Table 2. Factor Solution for K-RCMAS-2 SF Items and Loadings

| Item | Factor | |
|--|--------|-------|
| | PHY | COG |
| 1. Often I feel sick in my stomach | .698 | .002 |
| 2. I am nervous | .505 | .212 |
| 5. I have too many headaches | .743 | -.067 |
| 7. I wake up scared sometimes | .322 | .141 |
| 3. I often worry about something bad happening to me | .427 | .496 |
| 4. I fear other kids will laugh at me in class | -.040 | .944 |
| 6. I worry that others do not like me | .150 | .816 |
| 8. I get nervous around people | .145 | .510 |
| 9. I feel someone will tell me I do things the wrong way | .115 | .772 |
| 10. I fear other people will laugh at me | -.122 | 1.022 |

Note. PHY = physiological anxiety score for short form; COG = cognitive anxiety score for short form.

els in children aged 6-19 years (Reynolds & Richmond, 2008). The scale comprises 49 items to which the child must respond in a yes/no manner. This study used TOT along with three other subscales, namely PHY, WOR, and SOC. Furthermore, we calculated SF-TOT by adding up the number of 'yes' responses for the first ten items in RCMAS-2 as the manual recommends (Reynolds & Richmond, 2008). The contents of each item are listed in Table 2. The internal consistency of the SF-TOT in the Korean normative data was good ($\alpha = .92$). In the current sample, Cronbach's alpha was within the acceptable range ($\alpha = .79$).

Spielberger's State-Trait Anxiety Inventory for Children (STAIC) The State-Trait Anxiety Inventory for Children includes two separate 20-item scales that measure life-long enduring tendencies (trait version) and transitory (state version) level of childhood anxiety, respectively (Cho & Choi, 1989; Spielberger et al., 1973). However, since manifest anxiety of the RCMAS was originally conceptualized as a time-stable anxious disposition that indicate underlying emotional conflict or repressed feelings, the fluctuating nature of the STAIC state version was considered to not correspond to RCMAS-2 SF (Saviola et al., 2020; Schisler et al., 1998). Thus, we decided to use the trait version exclusively as trait anxiety is assumed to be a chronic tendency that determines individual differences in coping with external threats. The items specifically ask about the degree to which anxiety is felt in general and are rated on a 3-point Likert scale (1 to 3). The total score ranges from 20 to 60, with higher scores indicating greater severity of anxiety. The internal con-

sistency of the STAIC in the current sample was good ($\alpha = .91$).

Children's Depression Inventory, Second Edition (CDI 2)

Anxiety and depressive disorders are known as the most typical childhood internalization disorders (Strauss et al., 1988; Chorpita et al., 2000; Garber & Weersing, 2010; Zahn-Waxler et al., 2000). Several recent studies with the latest technologies such as network analysis posit that their boundary is rather vague (Boschloo et al., 2016; McElroy et al., 2018). In similar vein, as the importance of a dimensional approach toward psychopathology is increasingly emphasized, we tried to confirm the concurrent validity of the RCMAS-2 SF using CDI 2 (Kim et al., 2018; Kovacs, 2010) to examine if it can soundly measure children's internalization symptoms. The CDI 2 is a self-report questionnaire that assesses the level of depressive symptoms in children aged 7-17 years. It consists of 28 items; each of which offers three descriptive sentences that represent different levels of symptoms. The total score ranges from 0 to 56 and demonstrates strong internal consistency in the Korean normative data ($\alpha = .85$). In this study, the scores exhibited good reliability ($\alpha = .84$).

Data Analysis

A series of analyses on demographic data are performed in R version 4.0.0 (R Core Team, 2020) using "dplyr" (Wickham et al., 2019). To determine the number of factors to retain in EFA, we conducted a parallel analysis utilizing the "psych" package in advance (Revell, 2015). We consecutively carried out EFA and CFA to randomly assigned groups with the "lavaan" package (Rosseel, 2012). As the scale is an ordered-categorical measure whose values are simultaneously dichotomous yet hierarchical, we applied the weighted least square mean and variance adjusted chi-square (WLSMV) estimator with Geomin oblique rotation for the parameter estimation (Flora & Curran, 2004; Muthén, 1984; Muthén & Muthén, 2012). Since chi-square test of model fit is highly sensitive to the sample size, we used sample size-free indices, namely the comparative fit index (CFI), the Tucker-Lewis index (TLI), and the root mean square error of approximation (RMSEA) values. As in line with common standards, the goodness of fit was evaluated using the following robust criteria: $CFI \geq .95$, $TLI \geq .95$, and $RMSEA \leq .08$ (Hu & Bentler, 1999).

To test the measurement invariance of the scale, three models that gradually increase the invariance stringency, specifically (a) configural, (b) metric, and (c) scalar models, were estimated using the lavaan package (Rosseel, 2012). Each model was compared with its preceding simpler model with the chi-square difference test. However, since it bears the same inherent issue of the sample size-dependent aspect that amplifies the magnitude of negligible model differences, we report chi-square values but make decisions depending on change values in CFI and RMSEA, (Chen, 2007). This implies that the difference in each index lesser than .01, .015, and .03, respectively, was used to reject the more parsimonious model and proceed with the next comparison. As for the test of measurement invariance across age, we dichotomized our participants into two groups: elementary school children (aged 8–13) and secondary school adolescents (aged 14–19). Moreover, we tested partial invariance when significant fit-indices disparity between preceding and subsequent models were found.

Internal consistency was evaluated using Cronbach's alpha to assess the reliability of the scale. The test-retest reliability of the scale was also examined using Pearson correlation. For the test of concurrent validity, the correlation between SF-TOT and other proxies of childhood anxiety and depression severity, namely, STAIC and CDI 2 were investigated.

Results

Factor Analysis

Based on the results of the parallel analysis, which recommended retaining up to three factors, models with one-, two-, and three-factor structures were examined using EFA. According to the robust criteria, the unifactorial model exhibited an unacceptable

Table 3. Results of Exploratory Factor Analysis for K-RCMAS-2 SF

| Model | χ^2 | df | RMSEA (90% CI) | CFI | TLI |
|----------------|----------|----|------------------|------|------|
| 1-factor model | 235.512* | 35 | .085 (.075–.096) | .965 | .955 |
| 2-factor model | 45.667* | 26 | .031 (.015–.046) | .997 | .994 |
| 3-factor model | 26.005* | 18 | .024 (.000–.043) | .999 | .997 |

Note. RCMAS-2 SF = Revised Children's Manifest Anxiety Scale, Second Edition Short Form; *df* = degree of freedom; RMSEA = Root Mean Square Error of Approximation; CI = Confidence Interval; CFI = Comparative Fit Index; TLI = Tucker-Lewis Index.

* $p < .05$.

model fit. By contrast, statistics for the two-factor model suggested a good fit. The fit indices of the three-factor model were also in a good range, but the inspection of the item loadings on each factor was inconsistent with theory and previous suggestions. In particular, item 7 ("I wake up scared sometimes") loaded on WOR but not PHY. Furthermore, Item 9 ("I feel someone will tell me I do things the wrong way") loaded on WOR and SOC simultaneously, making the patterns of item loadings less clear compared to the preceding models. The two-factor model was selected as the best fit concerning the parsimony of the structure and item loadings. Such model was again verified to be appropriate in the results of the CFA performed on separate subgroups. The fit indices for estimated models of the RCMAS-2 SF are shown in Tables 3 and 4.

The standardized factor loadings for two-factor model are demonstrated in Table 2. After carefully examining the semantic contents and cultural backgrounds of the items, we found out that the first factor embraced the same items (items 1, 5, and 7) as the original PHY did. Yet, item 2 ("I am nervous") was newly added to PHY in the current study. This was rather interesting because the item was previously classified as the WOR domain in the literature (Reynolds & Richmond, 2008). Meanwhile, we designated the second factor as Cognitive Anxiety (COG) since it included items representing cognitive concerns about the future and social settings, integrating items originally included in WOR and SOC. The factor correlation was found to be .425, indicating a moderate positive association between PHY and COG. No item exhibited factor loading less than .30. Item 3 ("I often worry about something bad happening to me") had cross-loadings larger than .30 but were kept as a COG item due to structural and cultural considerations.

Measurement Invariance across Gender and Age

All the fit indices of the model fit regarding measurement invariance are demonstrated in Table 5. As shown, the minor fit changes

Table 4. Result of Confirmatory Factor Analysis for K-RCMAS-2 SF

| Model | χ^2 | df | RMSEA (90% CI) | CFI | TLI |
|----------------|----------|----|------------------|------|------|
| 2-factor model | 107.928 | 34 | .054 (.043–.066) | .986 | .982 |

Note. RCMAS-2 SF = Revised Children's Manifest Anxiety Scale, Second Edition Short Form; *df* = degree of freedom; RMSEA = Root Mean Square Error of Approximation; CI = Confidence Interval; CFI = Comparative Fit Index; TLI = Tucker-Lewis Index.

* $p < .05$.

Table 5. Goodness-of-Fit Indices and Model Comparisons for Measurement Invariance across Gender and Age

| | Model | χ^2 | df | CFI | Δ CFI | RMSEA | Δ RMSEA |
|--------|---------------|----------|----|------|--------------|-------|----------------|
| Gender | 1. Configural | 141.808 | 68 | .986 | | .054 | |
| | 2. Metric | 141.579 | 76 | .988 | .002 | .048 | .006 |
| | 3. Scalar | 152.559 | 74 | .985 | .003 | .054 | .006 |
| Age | 1. Configural | 155.457 | 68 | .985 | | .059 | |
| | 2. Metric | 162.214 | 76 | .985 | .000 | .055 | .004 |
| | 3. Scalar | 169.525 | 74 | .983 | .002 | .059 | .004 |

Note. df = degree of freedom; RMSEA = Root Mean Square Error of Approximation; CFI = Comparative Fit Index.

in CFI and RMSEA between configural, metric, and scalar models supported strong invariance across gender. Strong measurement invariance was also observed for age, implying that the factor structure, item loadings, and item thresholds of the RCMAS-2 SF were invariant across boys and girls of divergent ages.

Reliability and Validity of the RCMAS-2 SF

The internal consistency of the SF-TOT measured using Cronbach's alpha was satisfactory ($\alpha = .79$). This was comparable to the original RCMAS-SF in the normative sample ($\alpha = .92$) (Reynolds & Richmond, 2008). The examination of the item-total correlation exhibited that, with a range of .39 to .72, each item similarly contributed to the SF-TOT. Furthermore, the RCMAS-2 SF demonstrated a good two-weeks test-retest reliability, $r = .80$, $p < .05$. We then examined the concurrent validity of the SF-TOT based on the correlation coefficients between the STAIC-T and CDI 2. The results indicated strong correlations, $r = .70$, $p < .05$, $r = .83$, $p < .05$, respectively.

Discussion

This study validated the validation of the 10-item RCMAS-2 SF. Concerning the large body of literature that underscores cross-cultural differences and the complexity of pediatric anxiety, we explored the factor structure of the RCMAS-2 SF to shed light on the question whether the scale is solely composed of a single factor. Our analyses suggest that the two-factor structure is an optimal representation of the items in the RCMAS-2 SF, thus providing a broader understanding of the childhood anxiety and its measurement. Furthermore, the result exhibited satisfactory coefficients of reliability, validity, and measurement invariance across both gen-

der and age.

Most notably, the anxiety of children and adolescents measured by RCMAS-2 SF was not unidimensional. The results highlight that physiological anxiety should be considered as a construct independent of cognitive anxiety that incorporates worry and social concerns. The concept goes along the lines of the traditional information-processing theory of anxiety that proposes physiological arousal and cognitive bias as two principal features of anxious youth (Chorpita et al., 2000; Daleiden & Vasey, 1997; Lau & Waters, 2017). According to the theory, the cognitive component represents the interpretation bias of taking neutral or ambiguous situation as threatening stimuli (Creswell et al., 2005). Physiological component, on the other hand, is reflective of autonomic response of nervous system to threats such as increased blood pressure, elevated body temperature, discomfort within gastrointestinal system, and headache. These two markers of anxiety have been used as a proxy for the measurement of the efficacy of cognitive behavioral therapy on youth anxiety (Krebs et al., 2018). For instance, cognitive restructuring which challenges catastrophic thinking concerning threats and relaxation techniques for physiological arousal are two major non-pharmacological go-to techniques for anxiety relief (Beard et al., 2012; Reuland & Teachman, 2014). As a result, our data provide grounds for the creation of composite PHY and COG subscale scores that can be beneficial in research settings where the evaluation of the therapeutic effect of anxiety-CBT is needed.

Despite the separation of PHY, WOR and SOC were still combined as a single factor. One possible explanation for this convergence may arise from the semantic similarity between the items. For example, item 6 ("I worry that others do not like me") and item 8 ("I get nervous around people") were developed as WOR

items but their contextual meanings are very similar to that of SOC items (e.g., item 10 “I fear other people will laugh at me”). Besides, social anxiety intrinsically holds a worrying-nature because individuals with social anxiety engage in anticipatory concerns and post-event rumination (Clark & Wells, 1995; Rachman et al., 2000; Wells, 1995). In support of this, worry scores are found to be powerless in differentiating children with a generalized anxiety disorder from those with a social anxiety disorder (Whitmore, 2014). Furthermore, more than half of the children and adolescents with generalized anxiety disorder as their primary diagnosis have a comorbid social anxiety disorder (Holmes et al., 2014; Walkup et al., 2008). Hence, it seems logical that WOR and SOC remain as united construct whereas PHY was found to be a separate entity.

The item-level interpretation of the result was also notable. It is interesting that item 2 (“I am nervous”) was previously included in WOR but is now newly incorporated into the PHY construct. Although speculations remain to be examined, this may reflect the tendency of Asian parents who are more likely to accept somatic symptoms, such as stomachache or headache, rather than affective disturbance like anxiety (Essau et al., 2011; Yen et al., 2010). Due to such parenting attitude, Asian students may intrinsically find the source of nervousness in physiological symptoms rather than in cognitive or affective elements. Furthermore, in collectivistic Eastern culture, adolescents are taught to take uncomfortable feelings as temporal hindrance to endure or overcome; thus they may easily think of nervousness as a matter of momentary somatic ordeal without in-depth consideration. Also, item 3 (“I often worry about something bad happening to me”) held cross-loadings on both PHY and COG, indicating the likelihood of Asian students taking ‘something bad’ as physical or somatic symptoms.

Next, we examined the measurement invariance of the total score of the RCMAS-2 SF across gender and age. It is particularly noteworthy that scalar invariance was found in both tests, providing the basis of direct comparison of the SF-TOT scores between boys and girls in different age. This result concurs with the original RCMAS-2 SF study where strong and partial strong invariance across gender and age, respectively, were achieved (Lowe & Ang, 2016). It is also interesting that changes in fit indices of CFI RM-

SEA did not consistently decrease for more restricted models as expected. For example, when testing invariance across gender, metric model with more constraints depicted a better CFI value than a less restricted configural model. However, such pattern supposedly arises from the nature of WLSMV estimation used toward ordered-categorical variables (Bieda et al., 2016; Scholten, Velten, Bieda, Zhang, & Margraf et al., 2017). Our result, in sum, demonstrated that SF-TOT of RCMAS-2 and its underlying meanings can be validly compared across any group of youth.

Reliability analysis yielded acceptable internal consistency with a Cronbach’s alpha of .79. Furthermore, the two-weeks test-retest reliability coefficient was .80, demonstrating that the scale was credible in the time-stable aspect. This is in line with the theoretical assumptions of the RCMAS as manifest anxiety is assumed to be a life-long tendency toward perceived threats. The concurrent validity of the scale was supported by a moderate positive correlation found between the scores of the RCMAS-2 SF, STAI-C, and CDI 2. This proves that the SF-TOT to be a robust instrument that well represents the general psychopathology of the youth.

This study has several limitations that warrant comment. One was the lack of clinical participants in the study. Although our data had a large sample size, the inclusion of the clinical population would have heightened the usability of the scale. However, regarding the literature emphasizing that healthy children can report anxiety symptoms prior to the onset of anxiety disorders, the scale can still be widely used to detect probable pathogenic anxiety early. Another limitation is the absence of the measurement of participants’ socioeconomic status and other background information. More work is required to examine if other factors influence the SF-TOT score. Third, despite a wide range of age groups in our data, children between the ages six to seven (grades 1-2) were not included in the sample due to the possible misunderstanding of the item sentences. Future work may benefit from including younger children who can also suffer from anxiety.

In conclusion, this study provides psychometric evidence for the use of the RCMAS-2 SF. Convincing results of both reliability and validity analyses all demonstrate that it can be used in diverse areas such as research field or therapeutic settings where the measurement of baseline anxiety level is needed. Furthermore, it offers a more accurate understanding of the complex nature of pediatric

anxiety as the unidimensionality of the scale is rejected. The RC-MAS-2 SF score would benefit those who are interested in time-efficient comparisons of the anxiety severity between boys and girls.

Author contributions statement

HHY, a clinical psychology resident in the Department of Psychiatry at Samsung Medical Center, analyzed the data and wrote the manuscript. EHL, a clinical psychologist at the Depression Center at Samsung Medical Center, contributed to the data interpretation. He also led the preparation and correction of the manuscript. STH, a professor from the Department of Psychology at Chungbuk National University, designed the study process in general. He also vastly contributed to data collection the data for this study. SHH, a professor at the Chinju National University of Education, assisted in data collection and critically revised the article. JHK, a supervisor of the Department of Psychiatry at Samsung Medical Center and a professor of Sungkyunkwan University School of Medicine, served as the principal investigator of the research and supervised the research process in general. All the authors provided critical feedback, participated in the revision of the manuscript, and approved the final submission.

References

- Ahlen, J., Vigerland, S., & Ghaderi, A. (2018). Development of the Spence Children's Anxiety Scale-Short version (SCAS-S). *Journal of Psychopathology and Behavioral Assessment*, 40, 288-304.
- Ahmad, R., & Mansoor, I. (2011). What I think and feel: Translation and adaptation of Revised Children's Manifest Anxiety Scale, Second Edition (RCMAS-2) and its reliability assessment. *The International Journal of Educational and Psychological Assessment*, 8, 1-11.
- Barker, M. M., Beresford, B., Bland, M., & Fraser, L. K. (2019). Prevalence and incidence of anxiety and depression among children, adolescents, and young adults with life-limiting conditions: A systematic review and meta-analysis. *JAMA Pediatrics*, 173, 835-844.
- Beard, C., Sawyer, A. T., & Hofmann, S. G. (2012). Efficacy of attention bias modification using threat and appetitive stimuli: A meta-analytic review. *Behavior Therapy*, 43, 724-740.
- Bieda, A., Hirschfeld, G., Schönfeld, P., Brailovskaia, J., Zhang, X. C., & Margraf, J. (2016). Universal happiness? Cross-cultural measurement invariance of scales assessing positive mental health. *Psychological Assessment*, 29, 408-421.
- Bienvenu, O. J., & Ginsburg, G. S. (2007). Prevention of anxiety disorders. *International Review of Psychiatry*, 19, 647-654.
- Billieux, J., Rochat, L., Ceschi, G., Carré, A., Offerlin-Meyer, I., Defeldre, A. C., Khazaal, Y., Besche-Richard, C., Van der Linden, M. (2012). Validation of a short French version of the UPPS-P Impulsive Behavior Scale. *Comprehensive Psychiatry*, 53, 609-615.
- Bittner, A., Egger, H. L., Erkanli, A., Jane Costello, E., Foley, D. L., & Angold, A. (2007). What do childhood anxiety disorders predict? *Journal of Child Psychology and Psychiatry*, 48, 1174-1183.
- Boschloo, L., Schoevers, R. A., van Borkulo, C. D., Borsboom, D., & Oldehinkel, A. J. (2016). The network structure of psychopathology in a community sample of preadolescents. *Journal of Abnormal Psychology*, 125, 599.
- Cha, Y., Lee, E., Hwang, S., Hong, S., & Kim, J. (2020). Psychometric investigation of the Korean Version of the Revised Children's Manifest Anxiety Scale, Second Edition. *Korean Journal of Clinical Psychology*, 39, 203-214.
- Chan, S. M., & Leung, C. H. (2015). Factor structure of the screen for child anxiety-related emotional disorders (SCARED) in a community sample of Hong Kong Chinese adolescents. *Child Psychiatry and Human Development*, 46, 671-682.
- Chen, F. F. (2007). Sensitivity of goodness of fit indexes to lack of measurement invariance. *Structural Equation Modeling*, 14, 464-504.
- Cheung, G. W., & Rensvold, R. B. (2002). Evaluating goodness-of-fit indexes for testing measurement invariance. *Structural Equation Modeling*, 9, 233-255.
- Chisholm, D., Sweeny, K., Sheehan, P., Rasmussen, B., Smit, F., Cuijpers, P., & Saxena, S. (2016). Scaling-up treatment of depression and anxiety: A global return on investment analysis. *The Lancet Psychiatry*, 3, 415-424.
- Cho, S. C., & Choi, J. S. (1989). Development of the Korean form of the state-trait anxiety inventory for children. *Seoul Journal of Psychiatry*, 14, 150-157.
- Chorpita, B. F., Plummer, C. M., & Moffitt, C. E. (2000). Relations of tripartite dimensions of emotion to childhood anxiety and mood disorders. *Journal of Abnormal Child Psychology*, 28, 299-310.
- Chorpita, B. F., Yim, L., Moffitt, C., Umemoto, L. A., & Francis, S. E. (2000). Assessment of symptoms of DSM-IV anxiety and depression in children: A revised child anxiety and depression scale. *Behaviour Research and Therapy*, 38, 835-855.
- Copeland, W. E., Angold, A., Shanahan, L., & Costello, E. J. (2014). Longitudinal patterns of anxiety from childhood to adulthood: the Great Smoky Mountains Study. *Journal of the American Academy of Child and Adolescent Psychiatry*, 53, 21-33.
- Costello, E. J., Egger, H. L., & Angold, A. (2004). Developmental Epidemiology of Anxiety Disorders. In T. H. Ollendick & J. S. March (Eds.), *Phobic and anxiety disorders in children and adolescents: A clinician's guide to effective psychosocial and pharmacological treatments*. New York: Guilford Press.

- macological interventions (pp. 61-91). Oxford University Press.
- Clark, D. M., & Wells, A. (1995). A cognitive model. *Social Phobia: Diagnosis, Assessment, and Treatment*, 69, 1025.
- Creswell, C., Schniering, C. A., & Rapee, R. M. (2005). Threat interpretation in anxious children and their mothers: Comparison with nonclinical children and the effects of treatment. *Behaviour Research and Therapy*, 43, 1375-1381.
- Daleiden, E. L., & Vasey, M. W. (1997). An information-processing perspective on childhood anxiety. *Clinical Psychology Review*, 17, 407-429.
- Ebesutani, C., Regan, J., Smith, A., Reise, S., Higa-McMillan, C., & Chorpita, B. F. (2012). The 10-item positive and negative affect schedule for children, child and parent shortened versions: Application of item response theory for more efficient assessment. *Journal of Psychopathology and Behavioral Assessment*, 34, 191-203.
- Essau, C. A., Ishikawa, S. I., Sasagawa, S., Sato, H., Okajima, I., Otsubi, K., Georgiou, G. A., O'Callaghan, J., & Michie, F. (2011). Anxiety symptoms among adolescents in Japan and England: Their relationship with self-construals and social support. *Depression and Anxiety*, 28, 509-518.
- Fenley, A. R., Langer, D. A., Blake, P. R., & Pincus, D. B. (2021). The relation between risk perception and anxiety in youth: An investigation using a novel developmentally tailored measure. *Journal of Psychopathology and Behavioral Assessment*. <https://doi.org/10.1007/s10862-021-09893-0>
- Flora, D. B., & Curran, P. J. (2004). An empirical evaluation of alternative methods of estimation for confirmatory factor analysis with ordinal data. *Psychological Methods*, 9, 466-491.
- Gaesser, A. H., & Karan, O. C. (2017). A randomized controlled comparison of emotional freedom technique and cognitive-behavioral therapy to reduce adolescent anxiety: A pilot study. *The Journal of Alternative and Complementary Medicine*, 23, 102-108.
- Garber, J., & Weersing, V. R. (2010). Comorbidity of anxiety and depression in youth: implications for treatment and prevention. *Clinical Psychology: Science and Practice*, 17, 293-306.
- Gau, S. S., Chong, M. Y., Chen, T. H., & Cheng, A. T. (2005). A 3-year panel study of mental disorders among adolescents in Taiwan. *The American Journal of Psychiatry*, 162, 1344-1350.
- Hale, W. W., Crocetti, E., Raaijmakers, Q. A., & Meeus, W. H. (2011). A meta-analysis of the cross-cultural psychometric properties of the Screen for Child Anxiety Related Emotional Disorders (SCARED). *Journal of Child Psychology and Psychiatry*, 52, 80-90.
- Holmes, M. C., Donovan, C. L., Farrell, L. J., & March, S. (2014). The efficacy of a group-based, disorder-specific treatment program for childhood GAD—A randomized controlled trial. *Behaviour Research and Therapy*, 61, 122-135.
- Hu, L. T., & Bentler, P. M. (1999). Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Structural Equation Modeling*, 6, 1-55.
- Keller, M. B., Lavori, P. W., Wunder, J., Beardslee, W. R., Schwartz, C. E., & Roth, J. (1992). Chronic course of anxiety disorders in children and adolescents. *Journal of the American Academy of Child and Adolescent Psychiatry*, 31, 595-599.
- Kim, H. J., Lee, E. H., Hwang, S. T., Hong, S. H., & Kim, J. H. (2018). Psychometric properties of the Children's Depression Inventory-2 among a community-based sample of Korean children and adolescents. *Korean Journal of Clinical Psychology*, 37, 177-186.
- Kim, J. H., Tsai, W., Kodish, T., Trung, L. T., Lau, A. S., & Weiss, B. (2019). Cultural variation in temporal associations among somatic complaints, anxiety, and depressive symptoms in adolescence. *Journal of Psychosomatic Research*, 124, 109763.
- Kovacs M. (2010). *Children's depression inventory - second edition (CDI 2) manual*. NY: Multi-Health Services Publishing.
- Krebs, G., Pile, V., Grant, S., Degli Esposti, M., Montgomery, P., & Lau, J. Y. (2018). Research review: Cognitive bias modification of interpretations in youth and its effect on anxiety: A meta-analysis. *Journal of Child Psychology and Psychiatry*, 59, 831-844.
- Lau, J. Y., & Waters, A. M. (2017). Annual research review: An expanded account of information-processing mechanisms in risk for child and adolescent anxiety and depression. *Journal of Child Psychology and Psychiatry*, 58, 387-407.
- Li, H. C. W., & Lopez, V. (2007). Development and validation of a short form of the Chinese version of the State Anxiety Scale for Children. *International Journal of Nursing Studies*, 44, 566-573.
- Lowe, P. A. (2015b). The revised Children's manifest anxiety scale—second edition short form: Examination of the psychometric properties of a brief measure of general anxiety in a sample of children and adolescents. *Journal of Psychoeducational Assessment*, 33, 719-730.
- Lowe, P. A. (2015a). *Reliability and validity of the RCMAS-2 Short Form scores*. Paper presented at the annual meeting of the National Association of School Psychologists, Orlando, FL.
- Lowe, P. A., & Ang, R. P. (2016). Examination of measurement invariance across culture and gender on the RCMAS-2 short form among Singapore and US adolescents. *Journal of Psychoeducational Assessment*, 34, 192-198.
- Mahakwe, G., Johnson, E., Karlsson, K., & Nilsson, S. (2021). A systematic review of self-report instruments for the measurement of anxiety in hospitalized children with cancer. *International Journal of Environmental Research and Public Health*, 18, 1911.
- McElroy, E., Fearon, P., Belsky, J., Fonagy, P., & Patalay, P. (2018). Networks of depression and anxiety symptoms across development. *Journal of the American Academy of Child & Adolescent Psychiatry*, 57, 964-973.
- McEvoy, P. M., Salmon, K., Hyett, M. P., Jose, P. E., Gutenbrunner,

- C., Bryson, K., & Dewhirst, M. (2019). Repetitive negative thinking as a transdiagnostic predictor of depression and anxiety symptoms in adolescents. *Assessment*, 26, 324-335.
- Meredith, W. (1993). Measurement invariance, factor analysis and factorial invariance. *Psychometrika*, 58, 525-543.
- Merikangas, K. R., He, J. P., Burstein, M., Swanson, S. A., Avenevoli, S., Cui, L., Benjet, C., Georgiades, K., & Swendsen, J. (2010). Lifetime prevalence of mental disorders in US adolescents: Results from the National Comorbidity Survey Replication-Adolescent Supplement (NCS-A). *Journal of the American Academy of Child and Adolescent Psychiatry*, 49, 980-989.
- Muthén, B. O. (1984). A general structural equation model with dichotomous, ordered categorical, and continuous latent variable indicators. *Psychometrika*, 49, 115-132.
- Muthén, L. K., & Muthén, B. O. (2012). *Mplus: Statistical analysis with latent variables; User's guide (7th ed.)*. CA: Author.
- Pina, A. A., Little, M., Knight, G. P., & Silverman, W. K. (2009). Cross-ethnic measurement equivalence of the RCMAS in Latino and White youth with anxiety disorders. *Journal of Personality Assessment*, 91, 58-61.
- Raad, J. M. (2013). *Validation of the Revised Children's Manifest Anxiety Scale, (RCMAS-2) scores for children with specific learning disabilities* (Doctoral dissertation). <http://guides.lib.ku.edu/etd>
- Rachman, S., Grüter-Andrew, J., & Shafran, R. (2000). Post-event processing in social anxiety. *Behaviour Research and Therapy*, 38, 611-617.
- R Core Team (2020). *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. <https://www.R-project.org>
- Reuland, M. M., & Teachman, B. A. (2014). Interpretation bias modification for youth and their parents: A novel treatment for early adolescent social anxiety. *Journal of Anxiety Disorders*, 28, 851-864.
- Revelle, W. (2015). *Psych: Procedures for personality and psychological research*. <http://personality-project.org/r>
- Reynolds, C. R., & Richmond, B. O. (1985). *Revised children's manifest anxiety scale (RCMAS) Manual*. CA: Western Psychological Services.
- Reynolds, C. R., & Richmond, B. O. (2008). *The Revised Children's Manifest Anxiety Scale, Second Edition (RCMAS-2)*. Los Angeles, CA: Western Psychological Services.
- Rolstad, S., Adler, J., & Rydén, A. (2011). Response burden and questionnaire length: Is shorter better? A review and meta-analysis. *Value in Health*, 14, 1101-1108.
- Rosseel, Y. (2012). Lavaan: An R package for structural equation modeling and more. Version 0.5-12 (BETA). *Journal of Statistical Software*, 48, 1-36.
- Saviola, F., Pappaianni, E., Monti, A., Grecucci, A., Jovicich, J., & De Pisapia, N. (2020). Trait and state anxiety are mapped differently in the human brain. *Scientific Reports*, 10, 1-11.
- Schaumberg, K., Zerwas, S., Goodman, E., Yilmaz, Z., Bulik, C. M., & Micali, N. (2019). Anxiety disorder symptoms at age 10 predict eating disorder symptoms and diagnoses in adolescence. *Journal of Child Psychology and Psychiatry*, 60, 686-696.
- Schisler, T., Lander, J., & Fowler-Kerry, S. (1998). Assessing children's state anxiety. *Journal of Pain and Symptom Management*, 16, 80-86.
- Scholten, S., Velten, J., Bieda, A., Zhang, X. C., & Margraf, J. (2017). Testing measurement invariance of the Depression, Anxiety, and Stress Scales (DASS-21) across four countries. *Psychological Assessment*, 29, 1376.
- Simon, E., Bos, A. E., Verboon, P., Smeekens, S., & Muris, P. (2017). Psychometric properties of the Youth Anxiety Measure for DSM-5 (YAM-5) in a community sample. *Personality and Individual Differences*, 116, 258-264.
- Spence, S. H., Barrett, P. M., & Turner, C. M. (2003). Psychometric properties of the Spence Children's Anxiety Scale with young adolescents. *Journal of Anxiety Disorders*, 17, 605-625.
- Spielberger, C. D., Edwards, C. D., Lushene, R. E., Montuori, J., & Platzek, D. (1973). *State-trait anxiety inventory for children*. Palo Alto, CA: Consulting Psychologists Press.
- Strauss, C. C., Last, C. G., Hersen, M., & Kazdin, A. E. (1988). Association between anxiety and depression in children and adolescents with anxiety disorders. *Journal of Abnormal Child Psychology*, 16, 57-68.
- Van Oort, F. V. A., Greaves-Lord, K., Verhulst, F. C., Ormel, J., & Huizink, A. C. (2009). The developmental course of anxiety symptoms during adolescence: The TRAILS study. *Journal of Child Psychology and Psychiatry*, 50, 1209-1217.
- Van de Schoot, R., Lugtig, P., & Hox, J. (2012). A checklist for testing measurement invariance. *European Journal of Developmental Psychology*, 9, 486-492.
- Varela, R. E., & Hensley-Maloney, L. (2009). The influence of culture on anxiety in Latino youth: A review. *Clinical Child and Family Psychology Review*, 12, 217-233.
- Varela, R. E., & Biggs, B. K. (2006). Reliability and validity of the Revised Children's Manifest Anxiety Scale (RCMAS) across samples of Mexican, Mexican American, and European American children: A preliminary investigation. *Anxiety, Stress and Coping: An International Journal*, 19, 67-80.
- Walkup, J. T., Albano, A. M., Piacentini, J., Birmaher, B., Compton, S. N., Sherrill, J. T., Ginsburg, G. S., Rynn, M. A., McCracken, J., Waslick, B., Iyengar, S., March, J. S., & Kendall, P. C. (2008). Cognitive behavioral therapy, sertraline, or a combination in childhood anxiety. *New England Journal of Medicine*, 359, 2753-2766.
- Weems, C. F., Graham, R. A., Scott, B. G., Banks, D. M., & Russell, J. D. (2013). Suppressor effects and age differences in the expres-

- sion of anxious emotion. *Personality and Individual Differences*, 55, 283-287.
- Weisz, J. R., Suwanlert, S., Chaiyasit, W., Weiss, B., Achenbach, T. M., & Walter, B. R. (1987). Epidemiology of behavioral and emotional problems among Thai and American children: Parent reports for ages 6 to 11. *Journal of the American Academy of Child and Adolescent Psychiatry*, 26, 890-897.
- Weitkamp, K., & Seiffge-Krenke, I. (2019). The association between parental rearing dimensions and adolescent psychopathology: A cross-cultural study. *Journal of Youth and Adolescence*, 48, 469-483.
- Wells, A. (1995). Meta-cognition and worry: A cognitive model of generalized anxiety disorder. *Behavioural and Cognitive Psychotherapy*, 23, 301-320.
- Whitmore, M. J., Kim-Spoon, J., & Ollendick, T. H. (2014). Generalized anxiety disorder and social anxiety disorder in youth: Are they distinguishable? *Child Psychiatry and Human Development*, 45, 456-463.
- Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L. D. A., François, R., Grolemund, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T. L., Miller, E., Bache, S. M., Muller, K., Ooms, J., Robinson, D., Seidel, D. P., . . . & Yutani, H. (2019). Welcome to the Tidyverse. *Journal of Open Source Software*, 4, 1686.
- Wu, L. M., Liu, Y., Chen, H. M., Tseng, H. C., & Lin, W. T. (2016). Psychometric properties of the RCMAS-2 in pediatric cancer patients. *European Journal of Oncology Nursing*, 20, 36-41.
- Yen, C. F., Ko, C. H., Wu, Y. Y., Yen, J. Y., Hsu, F. C., & Yang, P. (2010). Normative data on anxiety symptoms on the Multidimensional Anxiety Scale for Children in Taiwanese children and adolescents: Differences in sex, age, and residence and comparison with an American sample. *Child Psychiatry and Human Development*, 41, 614-623.
- Zahn-Waxler, C., Klimes-Dougan, B., & Slattery, M. J. (2000). Internalizing problems of childhood and adolescence: Prospects, pitfalls, and progress in understanding the development of anxiety and depression. *Development and Psychopathology*, 12, 443-466.
- Zhou, X., Zhang, Y., Furukawa, T. A., Cuijpers, P., Pu, J., Weisz, J. R., Yang, L., Hetrick, S. E., Giocane, C. D., Cohen, D., James, A. C., Yuan, S., Whittington, C., Jiang, X., Teng, T., Cipriani, A., & Xie, P. (2019). Different types and acceptability of psychotherapies for acute anxiety disorders in children and adolescents: A network meta-analysis. *JAMA Psychiatry*, 76, 41-50.
- Zhu, Q., & Lowe, P. A. (2018). Examination of the psychometric properties of the Revised Children's Manifest Anxiety Scale-Second Edition scores among Chinese secondary school students. *Journal of Psychoeducational Assessment*, 36, 725-735.