



# First Report of the Leopard Slug (*Limax maximus*), a New Alien Species in Republic of Korea

Kibeom Park<sup>1</sup>, Jihee Kim<sup>2</sup>, Su-gon Park<sup>3</sup>, Chang-gon Mun<sup>4</sup>, Su-hwan Kim<sup>4</sup>, Youngho Cho<sup>4\*</sup>,  
Youngjun Park<sup>4\*</sup>

<sup>1</sup>Invirustech, Gwang-ju, Republic of Korea

<sup>2</sup>Research Institute of Polar, Incheon, Republic of Korea

<sup>3</sup>Yeungnam University, Gyeongsan, Republic of Korea

<sup>4</sup>National Institute of Ecology, Seoecheon, Republic of Korea

## ABSTRACT

On August 21, 2023, the National Institute of Ecology reported the sighting of an invasive slug resembling *Limax maximus* Linnaeus, 1758. near Bambat Tree Frog Park, Suwon-si, Korea. This slug is known for its aggressiveness. Specimens were collected around the park and from nearby farms. Through barcoding analysis and sequence comparison, it was identified as *L. maximus*, confirming its presence in Republic of Korea, alongside two previously identified *Limax* species, *Limax flavus* Linnaeus, 1758 and *Lehmannia marginatus* O. F. Müller, 1774. This study represents the first documented report of *L. maximus* in the country.

**Keywords:** Invasive alien species, Slugs, *Limax maximus*, Republic of Korea

## Introduction

*Limax maximus* (Gastropoda: Limacidae) is recognized as one of the world's most aggressively invasive terrestrial mollusks (Barker & McGhie, 1984; Barker, 1999). Originating from Europe, North Africa, and Asia, *L. maximus* has become an invasive species. It has successfully colonized various countries and islands, including, but not limited to, North America, South America, Africa, the Pacific Islands, Australia, Hawaii, New Zealand, and Japan (Barker

& McGhie, 1984; Roth & Sadeghian, 2006; Joe & Daehler, 2008; Hasegawa *et al.*, 2009). *L. maximus* exhibits omnivorous tendencies, with a diet comprising dead plant matter and fungi. It is considered to have an impact on plant diversity through selective feeding (Buschmann *et al.*, 2005; Strauss *et al.*, 2009). Notably, this species is regarded as a major agricultural pest in Florida, USA, particularly on cucumber, sweet potato, lettuce, Chinese cabbage, and horticulture plants, likely favoring tender, leafy vegetables, and other soft-stemmed plants (Stange *et al.*, 2009). Furthermore, in regions such as Hawaii and New Zealand, *L. maximus* and other slugs have been reported to be direct causes of local plant extinctions (Joe & Daehler, 2008).

Two other slug species in the genus *Limax*, namely *L. flavus* and *L. marginatus*, have been unintentionally introduced to the Republic of Korea (Lee *et al.*, 2010). However, to date, there is no documented evidence of established *L. maximus* populations in the Republic of Korea (CABI Digital Library, n.d.; GBIF Secretariat, 2023; National Institute

Received February 16, 2024; Revised March 19, 2024;  
Accepted March 20, 2024

\*Corresponding author: Youngho Cho  
e-mail themoth@nie.re.kr

https://orcid.org/0000-0001-5672-5121

Youngjun Park

e-mail aquatic@nie.re.kr

https://orcid.org/0000-0003-3916-5796



This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

of Biological Resources (NIBR), n.d.). The first documented occurrence of *L. maximus* in Japan was reported in 2006 (Hasegawa *et al.*, 2009). Since then, it has spread to Hokkaido and mainland Japan (Morii & Nakano, 2017). This implies that *L. maximus* could potentially thrive in all regions of Republic of Korea, given that Japan and Korea have similar latitude and environmental conditions.

This study aimed to provide the first confirmed reports of *L. maximus* in Republic of Korea and furnish ecological and molecular data that could aid in the effective management of indigenous natural ecosystems following its introduction. Specifically, this research focused on barcoding analysis for the identification and classification of *L. maximus*, offering fundamental information necessary for understanding its potential impact within ecosystems and for the development of management strategies.

## Materials and Methods

### Site description


Suwon-si is a special case city covering an area of 121 km<sup>2</sup> in the inland region of southwestern Gyeonggi-do, Republic of Korea. It is characterized by a gentle slope running from northeast to southwest. The topography to the south predominantly consists of plains, lacking high or rugged mountains in the vicinity. The area experiences a continental climate with an average annual temperature of 12.5°C, hot summers, and relatively cold winters according to data from [www.weather.go.kr](http://www.weather.go.kr). The survey was conducted three times daily and nightly between August and September 2023, focusing around Bambat Tree Frog Park in Suwon (N37°18'25.46", E126°58'16.1"), where the initial sighting of *L. maximus* occurred. The survey area characterized by forested landscapes has hobby farms, agricultural farms, and trails. It is nestled by mountains connecting to Deokseong Mountain (160 m above sea level), with the Bamkkot Village apartment complex within a 100 m radius. This proximity makes it a favored spot for hiking and nature exploration (Fig. 1).

### Sample collection and preservation

Slugs were collected alive, one at a time. They were immediately submerged in 70% ethanol and transported to the laboratory. Upon arrival, tissues were carefully excised using sterilized surgical scalpels for further analysis.

### DNA sequence analysis

A 50 mg tissue specimen was excised from a *L. maximus* slug sample using a sterilized knife and transferred into a 3 mm ceramic bead tube. To enhance DNA extraction efficiency from the slug sample, a modification was made to the Clear-S™ Quick DNA extraction kit's manual (IVT3002, Invirustech, Republic of Korea), which was then



**Fig. 1.** Survey site and capture locations of *Limax maximus* at the Bambat Tree Frog Park, Suwon-si, Republic of Korea.

used for the extraction process. In short, 210 µL of HBS buffer and 10 µL of TCEP-HCl were added to the tube containing the 50 mg sample. Using a homogenizer (MB-24YS, Biobase, China), the sample was repeatedly crushed three times at 50 Hz for 30 s. To the crushed sample, 20 µL of Proteinase K (20 mg/mL) was added. The mixture was then incubated at 55°C for 30 min. Following this, 490 µL of DLD buffer was added. After thorough mixing, the mixture was loaded into a spin column. Subsequent steps were executed in accordance with the kit's manual. DNA extraction results were assessed based on absorbance, with the OD260/280 value measuring 2.075. The yield was determined to be 0.55 µg.

Following DNA extraction, a polymerase chain reaction (PCR) was carried out to amplify the mitochondrial COI region with a length of approximately 1340 bp of the slug. The PCR process consisted of two stages. In the initial stage, primers mtCOI-1F-54 (5'-tttcaacaaycataargatattgg-3') and mtCOI-1R-53 (5'-aayaccaatagaaattatagcataaa-3') were used to amplify the first fragment. The second stage used mtCOI-2F (5'-ttagrcggggcaattactatrc-3') and mtCOI-2R (5'-cgaaaaca-gatattaacgaaccat-3') to amplify the second fragment. Each primer was designed with reference to the slug COI universal primer developed by Nitz *et al.* (2009). EzPCR™ Basic 5X Master mix was used to carry out the PCR process, starting with an initial denaturation step at 92°C for 4 min, followed by 40 cycles of amplification at 92°C for 1 min, 50°C for 1 min, and 72°C for 1 min. Following the completion of the PCR cycling, a final extension step was carried out at 72°C for 5 min. Then 2 µL of amplicon was mixed with 0.5X TBE buffer and subjected to 1.5% agarose gel electrophoresis at 150 V for 30 min. A Clear-S™ PCR/Gel DNA fragment purification kit (IVT3005, Invirustech, Republic of Korea) was used for sequence confirmation and purification of the PCR product. The purified DNA was then sent to Bionics, Inc. (Daejeon,

Republic of Korea) for Sanger sequencing. After obtaining sequences for both the first and second fragments, a consensus sequence was obtained using CAP3, yielding a mtCOI DNA sequence of 1381 bp in length.

Following electrophoresis of PCR amplicons on a 1.5% TBE agarose gel for DNA sequence analysis, lane 1 was loaded with a 100 bp DNA ladder, serving as a reference for molecular size. Lane 2 was loaded with PCR amplicons using the mtCOI-1F-54 and mtCOI-1R-53 primer set 1, and lane 3 was loaded with PCR amplicons using the mtCOI-2F and mtCOI-2R primer set 2. Electrophoresis revealed a distinct band around the target size of 830 bp in lane 2, accompanied by another band near the target size of 800 bp (Fig. 2a). These amplicons were then sequenced and assembled using the CAP3 software. Approximately 1381 bp of the slug mtCOI gene sequence was acquired for phylogenetic analysis, which was aligned using ClustalO. Following alignment, regions with gaps in the bilateral sequence were trimmed, resulting in a final gene sequence of 1231 bp, which was used for subsequent alignment and phylogenetic analysis. Using the program MEGA-X (Kumar *et al.*, 2018), a phylogenetic tree was constructed using the maximum likelihood method based on the Kimura 2-parameter model (Kimura, 1980; Kumar *et al.*, 2018). This process was repeated 100 times to generate a bootstrap consensus tree, clustering taxa based on maximum similarity.

## Results and Discussion

Through daily and nightly surveys, a total of 132 slugs were collected from the survey area. They exhibited a tendency to hide in various microhabitats in and around the farm environment, often under awnings and fences during the daytime. Conversely, a large number of individuals were collected along trails leading to nearby fields and near gullies during nocturnal surveys (Table 1). Additionally, these slugs showed increased activity levels during cloudy and rainy weather conditions, resulting in a notable increase in the number of specimens collected during these weather conditions and at night, consistent with their habits elsewhere (Bailey, 1981). In total, 97 (73.5%) slugs were collected along the trail connecting the farmland with the forested region, while 24 (18.2%) slugs were collected from within farms and fences, 15 (11.4%) slugs from the forested area, and two (1.5%) slugs from grassy plains (Table 1). These results indicate that the habitat range of *L. maximus* in Republic of Korea is quite broad, including gardens, parks, and greenhouses, consistent with a previous report (Weigand, 2014).

*L. maximus*, which is of the same genus as the domestically recorded species *L. flavus* and *L. marginata* in Korea, stands out due to its size and coloration. The largest *L. m-*

*aximus* specimen collected during this study measured 150 mm, showcasing no significant size deviation from specimens reported in Japan (Morii & Nakano, 2017). Unlike the domestic species, *L. maximus* exhibited a larger size, with a body color ranging from yellowish-white to grey and brown. This species can be distinguished by a distinct irregular pattern of brown spots and blotches extending from the head to the tail, including the mantle (Fig. 2b), making it larger than the domestic record species. In comparison, *L. flavus*, known as medium-sized slugs with an average body length of 130 mm, display a yellowish body adorned with subtle brown spots. Meanwhile, *L. marginatus* categorized as small slugs with an average body length of 50 mm, feature two distinct black stripes running along their body, distinguishing them from *L. maximus* (Kwon *et al.*, 2001).

Results revealed that *Limax maximus* S. Korea (2023), which represented the first discovery of this species in South Korea as identified in this survey, formed a phylogenetic clade with other *L. maximus* (Nitz *et al.*, 2009) (Fig. 3). In a BLASTn analysis, genes closely matched with those of the *L. maximus* isolate L1185 COI gene (FJ606469.1), sharing 99.7% sequence similarities and a query cover of 100%. This evidence, along with their appearance, confirmed that these slugs were *L. maximus*. Originally, *L. maximus* was primarily found in various forested areas across Europe.


**Table 1.** Coordinates of survey points and capture information of *Limax maximus* around the Bambat Tree Frog Park, Suwon-si, Republic of Korea

| Date | Coordinate |           | No. of individuals | Capture location | Time  |
|------|------------|-----------|--------------------|------------------|-------|
|      | Longitude  | Latitude  |                    |                  |       |
| 8/22 | 126.971382 | 37.310421 | 1                  | Forest           | Day   |
|      | 126.970886 | 37.311977 | 6                  | Fence            | Day   |
|      | 126.971099 | 37.310493 | 5                  | Farm             | Day   |
| 9/19 | 126.971328 | 37.31483  | 1                  | Farm             | Day   |
|      | 126.970520 | 37.313110 | 6                  | Farm             | Day   |
|      | 126.969718 | 37.313632 | 3                  | Farm             | Day   |
|      | 126.970146 | 37.313018 | 3                  | Farm             | Day   |
|      | 126.97026  | 37.312988 | 2                  | Grassland        | Day   |
|      | 126.97113  | 37.309745 | 74                 | Walking trails   | Night |
| 9/25 | 126.970909 | 37.311908 |                    |                  |       |
|      | 126.97113  | 37.309745 | 23                 | Walking trails   | Night |
|      | 126.970909 | 37.311908 |                    |                  |       |
|      | 126.971549 | 37.310539 | 8                  | Forest           | Night |





**Fig. 2.** (a) Results of slug universal mtCOI amplification; (b) General appearance of *Limax maximus*.



**Fig. 3.** Results of phylogenetic and sequence similarity molecular analysis of *Limax maximus*. Numbers on nodes indicate bootstrap support values.

However, its habitats have gradually expanded into anthropogenic environments, including urban and rural gardens, abandoned structures, parks, and greenhouses (Weigand, 2014). In Japan, this species was initially documented in 2012 within a remote natural forest in the Maruyama Forest Park in Sapporo, Hokkaido. Remarkably, within just six years, *L. maximus* was recorded in an area approximately 100 km from Sapporo (Morii & Nakano, 2017). Since then, this species has spread further into neighboring regions, including the Honshu, Kanto, and Chugoku prefectures (GBIF Secretariat, 2023). Given that Republic of Korea and Japan share similar latitude, it is reasonable to assume that *L. maximus* has the potential to inhabit the landscape of the Republic of Korea. As *L. maximus* has been classified as an invasive species in South Africa and Hawaii, USA (CABI Digital Library, n.d.) with potential to prey on and consume native slug species, it is important to consistently monitor the spread of this

species (Byers, 2000; Paustian & Barbosa, 2012; Rollo & Wellington, 1979). As highlighted above, distinct characteristics of *L. maximus*, such as its size and appearance, make it easy to distinguish it from native slug species and other *Limax* species in Republic of Korea. Thus, the use of citizen science, as has been successfully used in Japan (Morii & Nakano, 2017), could be an effective way to manage the establishment and spread of *L. maximus*.

### Conflict of Interest

The author declares that they have no competing interests.

### Acknowledgments

This work was first initiated through a tip from an influencer (TV Biological Encyclopedia). This work was

supported by a grant of “Nationwide Survey of Non-native Species in Republic of Korea (NIE-A-2024-12)” funded by the National Institute of Ecology, Republic of Korea.

## References

- Bailey, S.E.R. (1981). Circannual and circadian rhythms in the snail *Helix aspersa* Müller and the photoperiodic control of annual activity and reproduction. *Journal of Comparative Physiology*, 142, 89–94. <https://doi.org/10.1007/BF00605480>
- Barker, G.M. (1999). Naturalised Terrestrial Stylommatophora (Mollusca: Gastropoda). *Fauna of New Zealand*, 38, 1–254. <https://doi.org/10.7931/J2/FNZ.38>
- Barker, G.M. and McGhie, R.A. (1984). The biology of introduced slugs (Pulmonata) in New Zealand. Introduction and notes on *Limax maximus*. *New Zealand Entomologist*, 8, 106–111. <https://doi.org/10.1080/00779962.1984.9722482>
- Buschmann, H., Keller, M., Porret, N., Dietz, H., and Edwards, P.J. (2005). The effect of slug grazing on vegetation development and plant species diversity in an experimental grassland. *Functional Ecology*, 19, 291–298. <https://doi.org/10.1111/j.1365-2435.2005.00960.x>
- Byers, J.E. (2000). Competition Between Two Estuarine Snails: Implications for Invasions of Exotic Species. *Ecology*, 81, 1225–1239. [https://doi.org/10.1890/0012-9658\(2000\)081\[1225:CBTESI\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2000)081[1225:CBTESI]2.0.CO;2)
- CABI Digital Library. (2024). CABI Compendium. Wallingford, UK: CAB International. Retrieved January 14, 2024 from <https://www.cabidigitallibrary.org>
- GBIF Secretariat. (2023). *Limax maximus* Linnaeus, 1758 in GBIF Backbone Taxonomy. Checklist dataset. <https://doi.org/10.15468/39omei>. Retrieved March 13, 2024 from <https://www.gbif.org/dataset/d7dddbf4-2cf0-4f39-9b2a-bb099caae36c>
- Hasegawa, K., Fukuda, H., and Ishikawa, J. (2009). Discovery of naturalized populations of the introduced slug, *Limax maximus* in Tsuchiura City, Ibaraki Prefecture, Japan. *Chiribotan*, 39, 101–105.
- Joe, S.M., and Daehler, C.C. (2008). Invasive slugs as underappreciated obstacles to rare plant restoration: evidence from the Hawaiian Islands. *Biological Invasions*, 10, 245–255. <https://doi.org/10.1007/s10530-007-9126-9>
- Kimura, M. (1980). A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution*, 16, 111–120. <https://doi.org/10.1007/bf01731581>
- Kumar, S., Stecher, G., Li, M., Knyaz, C., and Tamura, K. (2018). MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms. *Molecular Biology and Evolution*, 35, 1547–1549. <https://doi.org/10.1093/molbev/msy096>
- Lee, J.S., Lee, Y.S., and Min, D.K. (2010) Introduced molluscan species to Republic of Korea. *Korean Journal of Malacology*, 26, 45–49.
- Morii, Y., and Nakano, T. (2017). Citizen science reveals the present range and a potential native predator of the invasive slug *Limax maximus* Linnaeus, 1758 in Hokkaido, Japan. *BioInvasions Records*, 6, 181–186. <https://doi.org/10.3391/bir.2017.6.3.01>
- National Institute of Biological Resources. (2024). NIBR Home Page. Retrieved January 9, 2024 from [www.species.nibr.go.kr](http://www.species.nibr.go.kr)
- Nitz, B., Heim, R., Schnepf, U.E., Hyman, I., and Haszprunar, G. (2009). Towards a new standard in slug species descriptions: the case of *Limax sarnensis* Heim & Nitz n. sp. (Pulmonata: Limacidae) from the Western Central Alps. *Journal of Molluscan Studies*, 75, 279–294. <https://doi.org/10.1093/mollus/eyp030>
- Paustian, M.E. and Barbosa, P. (2011). Overlap of food and microhabitat preferences among some native and nonnative slugs in mid-Atlantic forests of eastern North America. *Journal of Molluscan Studies*, 78, 92–99. <https://doi.org/10.1093/mollus/eyr041>
- Rollo, C.D. and Wellington, W.G. (1979). Intra- and inter-specific agonistic behavior among terrestrial slugs (Pulmonata: Stylommatophora). *Canadian Journal of Zoology*, 57, 846–855. <https://doi.org/10.1139/z79-104>
- Roth, B., and Sadeghian, P. (2006). *Checklist of the Land Snails and Slugs of California*. Santa Barbara, CA: Santa Barbara Museum of Natural History Contributions in Science.
- Shellfish Museum. (2001). *Korean Mollusks with Color Illustration (in Korean)*. Hangeul Book Publishing.
- Stange, L.A., Deisler, J.E., and Fasulo, T. (2002). *The Slugs of Florida (Gastropoda: Pulmonata). Featured Creatures*. Entomology and Nematology Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, EENY-087.
- Strauss, S.Y., Stanton, M.L., Emery, N.C., Bradley, C.A., Carleton, A., Dittich-Reed, D.R., *et al.* (2009). Cryptic seedling herbivory by nocturnal introduced generalists impacts survival, performance of native and exotic plants. *Ecology*, 90, 419–429. <https://doi.org/10.1890/07-1533.1>
- Weigand, A.M. (2014). Next Stop: Underground. Variable degrees and variety of reasons for cave penetration in terrestrial gastropods. *Acta Carsologica* 43, 175–183. <https://doi.org/10.3986/ac.v43i1.585>