



Planning Non-Invasive Conservation Genetic Experiments Based on Factors Affecting DNA Amplification Using Fecal Samples of Korean Long-Tailed Goral (*Naemorhedus caudatus*)

Baek-Jun Kim*

National Institute of Ecology, Seochon, Republic of Korea

ABSTRACT

In this review, we compared the success rates of DNA amplification and introduced the efficient non-invasive sampling of fecal samples collected from captive and wild Korean long-tailed gorals (*Naemorhedus caudatus*) by referring to previous non-invasive studies, including three important references (Kim *et al.*, 2008; Kim, 2021; Kim, 2022). A large difference in PCR success rates in the captive and wild populations was observed for mitochondrial (100 and 70.0%), sex-linked (44.4 and 20.8%), and microsatellite markers (73.9 and 34.8%, respectively). Out of the three types of genetic markers, the mitochondrial marker showed the highest success rate, followed by microsatellite and sex-linked markers. In addition, we estimated two factors that affected the PCR success, including the length of the amplified fragments (long, medium, and short) and the type of primer (universal and specific) in fecal samples from a captive population. The length of the PCR fragment was inversely proportional to the PCR success (5.3, 44.4, and 55.6% for long, medium, and short fragments, respectively), and the specific primer set (100%) was more efficient than the universal primer set (60.0%). This review is fundamental but would be greatly helpful for new non-invasive conservation genetic studies, particularly those that use fecal samples from captive and wild populations of rare endangered species. We recommend beginning conservation genetic experiments using mitochondrial markers and then nuclear markers, such as microsatellite and sex-linked markers, to save time, costs, and labor.


Keywords: Goral, Microsatellite, Mitochondrial cytochrome *b*, Non-invasive sampling, Sex-linked gene

Introduction

Feces is an attractive source of samples for the conservation genetic studies of free-ranging animals because of the ease of sampling and unprecedented sample size without

any need to capture or observe the animals (Taberlet and Luikart, 1999; Wehausen *et al.*, 2004). A number of studies have been conducted using these non-invasive samples in a variety of mammalian species, such as primates (baboon, bonobo, langur monkey, and orangutan), carnivores (badger, brown bear, canids, cougar, lynx, mustelids, otter, and seals), herbivores (bats, bighorn sheep, elephant, red deer, rabbit, reindeer, and rhinoceros), sirenians (dugong), cetaceans (dolphin), and marsupials (quokka and wombat) in the wild (Waits and Paetkau, 2005).

Received July 10, 2024; Revised July 22, 2024;
Accepted July 23, 2024

*Corresponding author: Baek-Jun Kim
e-mail naturalist71@nie.re.kr
 <https://orcid.org/0000-0002-8207-1931>



This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Many non-invasive sampling studies were aimed at examining various factors affecting DNA extraction yield, such as temperature during collection (Nsubuga *et al.*, 2004), the condition of the samples (Lucchini *et al.*, 2002), and the sampling season (Lucchini *et al.*, 2002), and comparing the efficiency of the sample storage methods (Nsubuga *et al.*, 2004) and DNA extraction protocols (Nsubuga *et al.*, 2004; Wehausen *et al.*, 2004). Also, different fecal pellet materials (outer, inner, and whole pellet) and different amounts of fecal pellet materials have been used to evaluate the rate of DNA amplification success (Wehausen *et al.*, 2004). However, non-invasive studies that quantitatively compared different DNA amplification success of different markers, including studies that compared DNA amplification success between different types of primer set or lengths of the amplified fragments, have rarely been conducted in wildlife populations (Taberlet and Luikart, 1999; Byrne *et al.*, 2021). Addressing these issues seems fundamental but is essential to planning a new research project related to wildlife conservation genetics.

In this review, we selected the long-tailed goral (*Naemorhedus caudatus*) as a target species for our analyses. The endangered long-tailed goral is a member of the genus *Naemorhedus*, tribe Rupicaprini, and subfamily Caprinae (Min *et al.*, 2004; An, 2006; Kim, 2021 and 2022). The goral species is registered in the International Union for Conservation of Nature Red List of Threatened Species and Convention on International Trade in Endangered Species Appendices I (Lee *et al.*, 2019) and is distributed across northeast Asia, including Russia, China, and Korea (Lee *et al.*, 2019). Their population size was presumed to be about 800 individuals in South Korea after a massive decline due to illegal hunting, over-exploitation, habitat destruction, and habitat fragmentation (Ministry of Environment 2002; Yang, 2002; Kim, 2021 and 2022). The population size of gorals has increased recently due to policies protecting the endangered species and their habitats, albeit slowly (Lee *et al.*, 2019).

The goal of this review was to provide researchers with efficient guidelines for initiating a new non-invasive study of a variety of wildlife species, particularly endangered species.

Review of Previous Non-Invasive Studies

In this review, we referred to previous studies using non-invasive fecal sampling. In particular, three critical goral papers for mitochondrial DNA (Kim, 2021), microsatellite DNA (Kim, 2022), and sex-linked DNA marker (Kim *et al.*, 2008) were reanalyzed. For this, we calculated the percentage of successful DNA amplifications for each factor (type of genetic marker, length of PCR fragments, and type of primer) in the references. In sequence, we

compared the rate of fecal DNA amplification success between three different types of markers using fecal samples obtained from captive and wild goral populations. We also compared whether different types of primer sets and lengths of the amplified fragments could affect the DNA amplification success rate in captive gorals.

Previous experimental studies reporting differences in fecal DNA amplification success rate between different types of genetic markers (e.g., mitochondrial, microsatellite, and sex-linked), lengths of the amplified fragments (e.g., long, medium, and short), or types of primer sets (e.g., universal and specific) are rare (Byrne *et al.*, 2021). Such a study would be a very useful starting point for conservation genetic experiments. The present review quantified and then compared different success rate of DNA amplification depending on various factors, including three markers, three lengths of amplified fragments, and two primer sets.

DNA Amplification Success of Three Different Markers

Three important papers reported that mitochondrial markers showed the highest percentage of DNA amplification success in both captive and wild goral populations, followed by microsatellite and sex-linked markers (Fig. 1). Large differences were seen between the two goral populations in the percentage of DNA amplification success for the three genetic markers. The captive population (24/24 = 100%) showed a higher mitochondrial marker PCR success rate (partial cytochrome *b* amplified by a primer set, NCF and NCR; Kim, 2021) than the wild population (28/40 = 70.0%). For microsatellite markers ($n = 6$ for each population; SY12A, SY50, SY58, SY129, SY17, and SY48), the captive population showed a 73.9% (17/23) PCR success rate but the wild population showed 34.8% (8/23; Kim, 2022). For the sex-linked marker (*AMELX/Y* amplified by a primer set, SE47 and SE48) between the two populations, the captive population (8/18 = 44.4%) showed a higher percentage of DNA amplification success than the wild population (5/24 = 21%) as microsatellite and mitochondrial markers (Kim *et al.*, 2008).

Feces in wild populations are more easily influenced by various factors (e.g., heat, desiccation, and decomposition; Chame, 2003) than in captive populations, such as those in zoos or farms. Therefore, researchers can collect and use fresher feces from captive populations than from wild populations. More degraded samples are found in wild than in captive populations. These factors might affect the DNA amplification success rate of the fecal samples. Non-invasive samples contain mitochondrial DNA (e.g., cytochrome *b* gene and D-loop region) and nuclear DNA (e.g., microsatellites and sex-linked genes). Hundreds




Fig. 1. Percentages (%) of successful DNA amplification of the captive and wild populations for three different genetic markers.




Fig. 2. Percentages (%) of successful DNA amplifications of a sex-linked gene in the captive population. Long: long-length amplified fragment; Medium: medium-length amplified fragment; Short: short-length amplified fragment.

to thousands of copies of mitochondrial DNA are found in a cell, whereas only two copies of nuclear DNA are found in most cells (Birky *et al.*, 1989). Thus, the rates of successful mitochondrial DNA extraction and amplification are higher than those of nuclear DNA (Frantzen *et al.*, 1998; Kohn *et al.*, 1999; Poole *et al.*, 2001; Lucchini *et al.*, 2002; Waits, 2004; Waits and Paetkau, 2005).

DNA Amplification Success According to Different Lengths of Amplified Fragments and Types of Primers

Regarding the percentage of DNA amplification success of sex-linked genes in the captive population according to different lengths of amplified fragments (e.g., long: *ZFX/Y* amplified by LGL331/335; and medium and short: *AMELX/Y* amplified by SE47/48 and SE47/53, respectively), the short-length fragment (10/18 = 55.6%) showed a higher value than the medium- (8/18 = 44.4%) and long-length fragments (1/19 = 5.3%; Fig. 2). In particular, the smallest value was below 10%, which is very low. Finally, two types of mitochondrial primers (e.g., specific: cytochrome *b* amplified by NCF/NCR; universal: cytochrome *b* amplified by L14841/H15149) showed different percentages of DNA amplification success in the captive population (Fig. 3). The goral-specific primers (100%) had much higher success rates than the universal primers (60.0%), although the length of the fragment amplified by NCF/NCR (378 bp) was slightly longer than that amplified by L14841/H15149 (376 bp, including primer regions).




Fig. 3. Percentages (%) of successful DNA amplifications of a mitochondrial gene in the captive population. Universal: universal primers; Specific: specific primers.

Both the length of the amplified fragments and the type of primer set could be important factors affecting successful DNA amplification in non-invasive studies. Short fragments will be amplified better from low quantities of degraded DNA (Frantzen *et al.*, 1998; Murphy *et al.*, 2000; Roon *et al.*, 2003). Sexing studies have been performed using various markers in mammals, such as *SRY*, *ZFX/Y*, and *AMELX/Y*. Among them, the *AMELX/Y* gene was

frequently applied because of its short-length fragment (Yamamoto *et al.*, 2002). The ZFX/Y primer set showed the lowest rate of successful sex-linked DNA amplification compared to other primer sets for AMELX/Y (SE47/48 and SE47/53). Waits and Paetkau (2005) previously reported that short target sequences are generally amplified more robustly. For species identification, two types of primers can be used in non-invasive studies, such as species-specific primers (Dalen *et al.*, 2004) and universal primers (Kocher *et al.*, 1989), designed using conserved regions in various species. The latter primers could be less efficient for DNA amplification success than the former primers due to a few nucleotides unmatched among different species.

Recommendation for Non-Invasive Conservation Genetic Experiments

In conclusion, the additional time, costs, and labor involved in working with fecal DNA should be considered in advance. This review showed that mitochondrial markers in both the captive and wild goral populations showed the highest percentage of success, followed by microsatellite and sex-linked markers. Fecal DNA amplification success rate was much higher in the captive than in the wild population due to relatively low DNA degradation. The DNA amplification success rate was much higher for shorter amplified fragments and a specific primer set than for longer amplified fragments and a non-specific primer set, respectively. Better results might be obtained by identifying and selecting the optimal amplified fragment length and type of primer set. This could be helpful as a first step for planning non-invasive genetic studies. We recommend beginning genetic experiments using mitochondrial markers and then using nuclear markers, such as microsatellite and sex-linked markers, to save time, costs, and labor. Also, we suggest that it is better to use fecal samples collected from a captive population in a pilot experiment and then use those collected from a wild population to decrease unnecessary experimental effort and increase experimental efficiency. Finally, it is expected that using short DNA markers rather than long DNA markers and specific markers rather than universal markers will yield better results in non-invasive conservation genetic studies.

Author Contributions

The author confirms sole responsibility for the following: study conception and design, data collection, analysis and interpretation of results, and manuscript preparation.

Conflict of Interest

The authors declare that they have no competing interests.

Acknowledgments

This work was supported in part by the National Institute of Ecology, funded by the Ministry of Environment (MOE) of the Republic of Korea (No. NIE-B-2024-18). It was also partially supported by the Research Institute for Veterinary Science and the Brain Korea 21 Program for Veterinary Science, Seoul National University. We thank Professor H. Lee and Professor Y. J. Won for helping with conducting previous Korean goral studies and Dr. H. J. Bae for giving comments for this review paper.

References

- An, J.H. (2006). Development and characterization of microsatellite markers for endangered Korean goral (*Nemorhaedus caudatus raddeanus*) and its molecular phylogenetic status. Doctoral dissertation. Seoul National University, Seoul.
- Birky, C., Fuerst, P., and Maruyama, T. (1989). Organelle gene diversity under migration, mutation and drift: Equilibrium expectations, approach to equilibrium, effects of heteroplasmic cells, and comparison to nuclear genes. *Genetics*, 121, 613-627. <https://doi.org/10.1093/genetics/121.3.613> [PMID: 2714640 PMID: PMC1203645]
- Byrne, M.S., Peralta, D.M., Ibañez, E.A., Nardelli, M., and Túnez, J.I. (2021). Non-invasive sampling techniques applied to conservation genetic studies in mammals. In M. Nardelli, and J.I. Túnez (Eds.), *Molecular Ecology and Conservation Genetics of Neotropical Mammals* (pp. 63-83). Springer https://doi.org/10.1007/978-3-030-65606-5_4
- Chame, M. (2003). Terrestrial mammal feces: A morphometric summary and description. *Mem Inst Oswaldo Cruz, Rio de Janeiro*, 98, 71-94. <https://doi.org/10.1590/s0074-02762003000900014> [PMID: 12687767]
- Dalen, L., Gotherstrom, A., and Angerbjorn, A. (2004). Identifying species from pieces of feces. *Conservation Genetics*, 5, 109-111.
- Frantzen, M.A.J., Silk, J.B., Ferguson, J.W., Wayne, R.K., and Kohn, M.H. (1998). Empirical evaluation of preservation methods for fecal DNA. *Molecular Ecology*, 7, 1423-1428. <https://doi.org/10.1046/j.1365-294x.1998.00449.x> [PMID: 9787450]
- Kim, B.J. (2021). Simple assessment of taxonomic status and genetic diversity of Korean long-tailed goral (*Naemorhedus caudatus*) based on partial mitochondrial Cytochrome *b* gene using non-invasive fecal samples. *Proceedings of the National Institute of Ecology*, 2, 32-41. <https://doi.org/10.22920/PNIE.2021.2.1.32>
- Kim, B.J. (2022). Microsatellite markers for non-invasive examination of individual identity, genetic variation, and population differentiation in two populations of Korean long-tailed goral (*Naemorhedus caudatus*). *Proceedings of the National Institute of Ecology*, 3, 191-198. <https://doi.org/10.22920/PNIE.2022.3.4.191>
- Kim, B.J., Lee, Y.S., An, J., Park, H.C., Okumura, H., Lee, H., *et al.* (2008). Species and sex identification of the Korean goral (*Nemorhaedus caudatus*) by molecular analysis of

- non-invasive samples. *Molecules and Cell*, 26, 314-318. [https://doi.org/10.1016/S1016-8478\(23\)14001-5](https://doi.org/10.1016/S1016-8478(23)14001-5)
- Kocher, T.D., Thomas, W.K., Meyer, A., Edwards, S.V., Paabo S, Villablanca, F.X. *et al.* (1989). Dynamics of mitochondrial DNA evolution in animals: Amplification and sequencing with conserved primers. *Proceeding of the National Academic of Science USA*, 86, 196-200. <https://doi.org/10.1073/pnas.86.16.6196> [PMID: 2762322 PMCID: PMC297804]
- Kohn, M.H., York, E.C., Kamradt, D.A., Haught, G., Sauvajot, R.M., and Wayne, R.K. (1999). Estimating population size by genotyping faeces. *Proceedings of the Royal Society of London Series B*, 266, 657-663. <https://doi.org/10.1098/rspb.1999.0686> [PMID: 10331287 PMCID: PMC1689828]
- Lee, S., Kim, B.J., and Bhang, K.J. (2019). Habitat analysis of endangered Korean long-tailed goral (*Naemorhedus caudatus raddeanus*) with weather forecasting model. *Sustainability*, 11, 6086. <https://doi.org/10.3390/su11216086>
- Lucchini, V., Fabbri, E., Marucco, F., Ricci, S., Boitani, L., and Randi, E. (2002). Noninvasive molecular tracking of colonizing wolf (*Canis lupus*) packs in the Western Italian Alps. *Molecular Ecology*, 11, 857-868. <https://doi.org/10.1046/j.1365-294x.2002.01489.x> [PMID: 11975702]
- Min, M.S., Okumura, H., Jo, D.J., An, J.H., Kim, K.S., Kim, C.B., *et al.* (2004). Molecular phylogenetic status of the Korean goral and Japanese serow based on partial sequences of the mitochondrial cytochrome *b* gene. *Molecules and Cells*, 17, 365-372. [https://doi.org/10.1016/S1016-8478\(23\)13052-4](https://doi.org/10.1016/S1016-8478(23)13052-4) [PMID: 15179056]
- Ministry of Environment (2002). New Technique for the Restoration of Endangered Species in Korea, Gwacheon: Ministry of Environment.
- Murphy, M.A., Waits, L.P., and Kendall, K.C. (2000). Quantitative evaluation of fecal drying methods for brown bear DNA analysis. *Wildlife Society Bulletin*, 28, 951-957. <https://www.jstor.org/stable/3783853>
- Nsubuga, A.M., Robbins, M.M., Roeder, A.D., Morin, P.A., Boesch, C., and Vigilant, L. (2004). Factors affecting the amount of genomic DNA extracted from ape faeces and the identification of an improved sample storage method. *Molecular Ecology*, 13, 2089-2094. <https://doi.org/10.1111/j.1365-294X.2004.02207.x> [PMID: 15189228]
- Poole, K.G., Mowat, G., and Fear, D.A. (2001). DNA-based population estimate for grizzly bears *Ursus arctos* in Northeastern British Columbia, Canada. *Wildlife Biology*, 7, 105-115. <https://doi.org/10.2981/wlb.2001.014>
- Roon, D.A., Waits, L.P., and Kendall, K.C. (2003). A quantitative evaluation of two methods for preserving hair samples. *Molecular Ecology Notes*, 3, 163-166. <https://doi.org/10.1046/j.1471-8286.2003.00358.x>
- Taberlet, P., and Luikart, G. (1999). Non-invasive genetic sampling and individual identification. *Biological Journal of the Linnean Society*, 68, 41-55. <https://doi.org/10.1111/j.1095-8312.1999.tb01157.x>
- Waits, L.P. (2004). Using noninvasive genetic sampling to detect and estimate abundance of rare wildlife species. In W.L. Thomas (Ed.), *Sampling rare or elusive species: concepts, designs and techniques for estimating population parameters*. (pp. 211-228). Island Press.
- Waits, L.P., and Paetkau, D. (2005). Noninvasive genetic sampling tools for wildlife biologists: A review of applications and recommendations for accurate data collection. *Journal of Wildlife Management*, 69, 1419-1433. [https://doi.org/10.2193/0022-541X\(2005\)69\[1419:NGSTFW\]2.0.CO;2](https://doi.org/10.2193/0022-541X(2005)69[1419:NGSTFW]2.0.CO;2)
- Wehausen, J.D., Ramey, II R.R., and Epps, C.W. (2004). Experiments in DNA extraction and PCR amplification from bighorn sheep feces: The importance of DNA extraction method. *Journal of Heredity*, 95, 503-509. <https://doi.org/10.1093/jhered/esh068> [PMID: 15475396]
- Yamamoto, K., Tsubota, T., Komatsu, T., Katayama, A., Murase T., Kita, I., and Kudo, T. (2002). Sex identification of Japanese black bear, *Ursus thibetanus japonicus*, by PCR based on amelogenin gene. *Journal of Veterinary Medical Science*, 64, 505-508. <https://doi.org/10.1292/jvms.64.505> [PMID: 12130835]
- Yang, B.G. (2002). Systematics, ecology and current population status of the goral, *Naemorhedus caudatus*, in Korea. Doctoral dissertation. Chungbuk National University, Cheongju.